



HAL
open science

Integration of retroviruses: a fine balance between efficiency and danger.

Alain Fischer, Marina Cavazzana-Calvo

► **To cite this version:**

Alain Fischer, Marina Cavazzana-Calvo. Integration of retroviruses: a fine balance between efficiency and danger.. PLoS Medicine, 2005, 2, pp.e10. 10.1371/journal.pmed.0020010 . inserm-00080151

HAL Id: inserm-00080151

<https://inserm.hal.science/inserm-00080151>

Submitted on 14 Jun 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Perspectives

Integration of Retroviruses: A Fine Balance between Efficiency and Danger

Alain Fischer*, Marina Cavazzana-Calvo

One of the key characteristics of retroviruses is their ability to integrate a copy of the DNA reverse-transcribed from their viral RNA genome into host cell DNA. This integration is mediated by a preintegration complex (PIC) comprising viral DNA, reverse transcriptase, and integrase, as well as poorly characterized host proteins [1]. It is this property that makes retroviruses good vectors for the transfer of therapeutic genes, and many such vectors have been made.

Stages in Vector Development

Early vectors included those derived from murine oncoretroviruses (RVs) such as murine leukemia virus (MLV). A limitation of these viruses was that their PIC requires that the nuclear membrane is dissolved so that the PIC can come in direct contact with host cell DNA; hence, there is efficient integration only in dividing cells. Later

The Perspectives section is for experts to discuss the clinical practice or public health implications of a published study that is freely available online.

vectors were based on lentiviruses (LVs) such as human immunodeficiency virus and simian immunodeficiency virus (SIV), whose PIC can penetrate into the nucleus so that integration can occur in nondividing cells that are in the G1 phase of the cell cycle.

Insertional Mutagenesis

MLV-derived vectors have been used with success to achieve sustained correction of two forms of severe combined immunodeficiency (SCID)—SCID-X1 (Gamma-c deficiency) [2] and adenosine deaminase deficiency [3]. In both cases, hematopoietic progenitors were infected out of the patient's body with nonreplicative MLV-derived vectors. However, integration carries the risk of insertional mutagenesis. This mutagenesis has been demonstrated in the chicken by using replicative RVs. RV integration close to protooncogenes has been shown to induce their activation, leading to tumorigenesis. Nonreplicative MLV vectors have also been reported to induce insertional mutagenesis in a murine model [4] and, more worryingly, in two patients from the

SCID-X1 trial [5]. In both instances, it is suggested that cooperation between vector-associated transgene expression (dLNGFR in one case and common Gamma chain in the other) and long

Citation: Fischer A, Cavazzana-Calvo M (2005) Integration of retroviruses: A fine balance between efficiency and danger. *PLoS Med* 2(1): e10.

Copyright: © 2005 Fischer and Cavazzana-Calvo. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abbreviations: LV, lentivirus; MLV, murine leukemia virus; PIC, preintegration complex; RV, murine oncoretrovirus; SCID, severe combined immunodeficiency; SIV, simian immunodeficiency virus

Alain Fischer is Professor of Pediatric Immunology and Head of the Pediatric Immunology Department and the Inserm Research Unit 429 at Necker University Hospital, Paris, France. Marina Cavazzana-Calvo is Professor of Hematology and Head of the Biotherapy Department at Necker University Hospital, Paris, France.

*To whom correspondence should be addressed. E-mail: alain.fischer@nck.ap-hop-paris.fr

Competing Interests: The authors declare that they have no competing interests.

DOI: 10.1371/journal.pmed.0020010


terminal repeat–driven enhancement of protooncogene expression (*evl-1* and *LMO-2*, respectively) was responsible for aberrant clonal proliferation. No such events have been reported yet in the use of LV vectors in experimental settings.

Sites of Retroviral Integration

It was initially believed that integration of retroviruses occurred randomly, but the advent of technology allowing the assessment of RV or LV integration into host cell genomes has led to a reassessment of this assumption. Using a combination of ligation-mediated polymerase chain reactions and sequencing of amplified integration sites (unique sequences made from the viral long terminal repeat and the host-genome-associated sequence), it is possible to determine all the integration sites that can be found in a given transduced cell population (or its progeny). Exact mapping can be done back to the human (or relevant animal) genome database. However, even this methodology may not detect all of the integration sites present in a given transduced cell population.

Key papers have determined the “rules of the game” for RV and LV integration into a variety of cell lines [6,7,8]; these rules seem to differ between the virus types. In both cases, however, the integration pattern is not random. RV integration tends to be close to transcription start sites of active genes—close enough to regulatory sequences to potentially exert a long terminal repeat–mediated enhancer effect [7,8]. By contrast, LVs integrate mostly in transcription units, with a preference for actively transcribed genes, but do not target the region downstream of transcription start sites [6,8]. These data indicate that there are virus-specific PIC-associated determinants that cause specific targeting with the host cell genome. However, much remains to be done to identify viral factors and host ligands involved in these interactions.

Data from these pioneering papers were obtained by in vitro infection of mature cells or cell lines with the relevant RV or LV [6,7,8]. However, it is possible that the pattern of integration might differ in other cell subsets, particularly immature cells such as hematopoietic progenitors. In



DOI: 10.1371/journal.pbio.0020423.g002

Figure 1. Distribution of MLV and SIV Integration Sites within a 60-kb Window Centered on Transcription Start Sites

The vertical arrow points to 0 kb. Each gray bar corresponds to the percentage of SIV integration sites within a 5-kb interval, and black bars correspond to the percentages of MLV integration sites in a 5-kb interval. The distribution of a set of 65,000 in silico-generated random integration sites is represented by the dashed line. (Source: [9].)

fact, as shown by Mitchell et al. [8], infection of mature cells of different tissue types leads to a partially distinct pattern of integration sites related to the set of genes transcribed in these different cell types [8].

In Vivo Models

In a paper published in last month’s *PLoS Biology*, Hematti et al. [9] took the in vivo analysis further by analyzing the pattern of integration sites in cells derived from simian hematopoietic progenitor cells transduced either with a RV (MLV) or a LV (SIV) vector and transplanted into monkeys. This experimental setting is the closest possible to human trials. The results essentially confirm the nonrandom insertion pattern of both types of vectors as shown by the analysis of cell lines transduced in vitro. The analysis showed the Gaussian distribution of insertions centered on the transcription start site—thus very close to regulatory elements—of RV (Figure 1), and the preference of LV for the transcription units, with a concentration of integrations into some gene-dense regions [9]. A study such as this is therefore a useful piece of preclinical work that will help the interpretation of the analysis of clinical samples.

Similar data were also obtained by Laufs et al. in the analysis of a set of RV integration sites into human hematopoietic progenitors xenotransplanted into immunodeficient mice [10]. It thus appears that the overall distinct pattern of RV and LV integration could be independent of transduced cell types (immature versus differentiated, and tissue type). However, the targeted genes could differ considerably depending on the set of genes expressed in the target cell. Several parameters could potentially influence the transcription profile in a clinical setting, including stage of cell maturity, tissue type, ex vivo transduction culture conditions, patient age [5], underlying genetic disease, and any modification of the vector.

Designing Future Vectors

It will thus be essential to build a free, accessible database incorporating all relevant information gathered from both experimental and clinical settings. In this respect, information gathered from the SCID-X1 and adenosine deaminase deficiency trials is awaited with great interest. Combining all available information will be the only way to determine the frequency of

insertions that have a potential to induce activation of a protooncogene (a risk primarily associated with the use of RV) or to induce disruption of a regulatory gene (a risk primarily associated with the use of LV). Studies of relevant gene expression activation or suppression should therefore be carried out in parallel.

It is only from these multiple analyses, including a careful comparison of the oncogenic potential of vectors in relevant animal models, that a precise assessment of the risk associated with use of retroviruses for gene therapy will come. These data will thus be the basis for objective

comparison between technologies and for the design of safer vectors. ■

References

1. Coffin J, Hugues S, Varmus H (1997) *Retroviruses*. New York: Cold Spring Harbor Laboratory Press. 843 p.
2. Cavazzana-Calvo M, Hacein-Bey S, de Saint Basile G, Gross F, Yvon E, et al. (2000) Gene therapy of human severe combined immunodeficiency (SCID)-X1 disease. *Science* 288: 669–672.
3. Aiuti A, Slavin S, Aker M, Ficara F, Deola S, et al. (2002) Correction of ADA-SCID by stem cell gene therapy combined with nonmyeloablative conditioning. *Science* 296: 2410–2413.
4. Li Z, Dullmann J, Schiedlmeier B, Schmidt M, von Kalle C, et al. (2002) Murine leukemia induced by retroviral gene marking. *Science* 296: 497.
5. Hacein-Bey-Abina S, Von Kalle C, Schmidt M, McCormack MP, Wulfraat N, et al. (2003) LMO2-associated clonal T cell proliferation in two patients after gene therapy for SCID-X1. *Science* 302: 415–419.
6. Schroder AR, Shinn P, Chen H, Berry C, Ecker JR, et al. (2002) HIV-1 integration in the human genome favors active genes and local hotspots. *Cell* 110: 521–529.
7. Wu X, Li Y, Crise B, Burgess SM (2003) Transcription start regions in the human genome are favored targets for MLV integration. *Science* 300: 1749–1751.
8. Mitchell RS, Beitzel BF, Schroder AR, Shinn P, Chen H, et al. (2004) Retroviral DNA integration: ASLV, HIV, and MLV show distinct target site preferences. *PLoS Biol* 2: e234.
9. Hematti P, Hong BK, Ferguson C, Adler R, Hanawa H, et al. (2004) Distinct genomic integration of MLV and SIV vectors in primate hematopoietic stem and progenitor cells. *PLoS Biol* 2: e423.
10. Laufs S, Gentner B, Nagy KZ, Jauch A, Benner A, et al. (2003) Retroviral vector integration occurs in preferred genomic targets of human bone marrow-repopulating cells. *Blood* 101: 2191–2198.