



HAL
open science

Effect of TNF and LTA polymorphisms on biological markers of response to oxidative stimuli in coal miners: a model of gene-environment interaction. Tumour necrosis factor and lymphotoxin alpha.

Rachel Nadif, Anne Jedlicka, Margaret Mintz, Jean-Pierre Bertrand, Steven R. Kleeberger, Francine Kauffmann

► To cite this version:

Rachel Nadif, Anne Jedlicka, Margaret Mintz, Jean-Pierre Bertrand, Steven R. Kleeberger, et al.. Effect of TNF and LTA polymorphisms on biological markers of response to oxidative stimuli in coal miners: a model of gene-environment interaction. Tumour necrosis factor and lymphotoxin alpha.. Journal of Medical Genetics, 2003, 40, pp.96-103. <inserm-00086489>

HAL Id: inserm-00086489

<https://inserm.hal.science/inserm-00086489v1>

Submitted on 30 Jan 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Role of *TNF* and *LTA* polymorphisms on biological markers of response to oxidative stimuli in coal miners: a model of gene-environment interaction

Rachel Nadif,¹ Anne Jedlicka,² Margaret Mintz,² Jean-Pierre Bertrand,³ Steven Kleeberger^{2,4},
Francine Kauffmann¹

¹INSERM U 472-IFR69, Villejuif, France

²Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD USA

³Service Medical HBL, Freyming- Merlebach, France

⁴ Present affiliation: Laboratory of Pulmonary Pathobiology, National Institute of Environmental Health Sciences, Research Triangle Park, NC USA

Address for reprints:

Rachel NADIF, PhD
Institut National de la Santé et de la Recherche Médicale
Epidémiologie et Biostatistique U 472
16 avenue Paul Vaillant Couturier
94807 Villejuif cedex, France

Phone number: 33 1 45 59 51 89

Fax number: 33 1 45 59 51 69

E-mail: nadif@vjf.inserm.fr

Corresponding author for the paper:

Rachel NADIF, PhD
Institut National de la Santé et de la Recherche Médicale
Epidémiologie et Biostatistique U 472
16 avenue Paul Vaillant Couturier
94807 Villejuif cedex, France

Phone number: 33 1 45 59 51 89

Fax number: 33 1 45 59 51 69

E-mail: nadif@vjf.inserm.fr

Running title: *TNF*, *LTA* and response to oxidative stress

Abstract

Introduction - Interaction between genetic background and oxidative environmental stimuli in the pathogenesis of human lung disease has been largely unexplored.

Methods - A prospective epidemiologic study was undertaken in 253 coal miners.

Intermediate quantitative phenotypes of response to oxidant exposure, including erythrocyte glutathione peroxidase (GSH-Px) and catalase activities were studied. Oxidant exposures studied were smoking habits and cumulative dust exposure assessed by job history and ambient measures. Disease phenotypes included subclinical computed tomography score at the first survey and X-ray profusion grades twice, 5 years apart, to assess established coal workers' pneumoconiosis (CWP). Miners were genotyped for common functional polymorphisms in the gene for tumor necrosis factor α (*TNF*) and lymphotoxin α (*LTA*), two proinflammatory cytokines that have been implicated in the pathogenesis of chronic lung diseases.

Results - Regarding gene-environment interaction on intermediate phenotypes, results showed interaction of a promoter polymorphism at the -308 position in *TNF* with occupational exposure on erythrocyte GSH-Px activity with a significant association in those with high exposure ($p = 0.003$) whereas no association was observed among those with low exposure (interaction $p = 0.06$). Regarding gene-intermediate phenotype interaction on clinical outcome, results showed an association of CWP prevalence with a *NcoI* polymorphism in *LTA* in those with low catalase activity ($p = 0.05$) whereas no association was observed in those with high activity (interaction $p = 0.03$). No other significant association was observed.

Conclusion - Results suggest that interactions of genetic background with environmental exposure and intermediate response phenotypes are important components in the pathogenesis of CWP.

Key words – *TNF*; *LTA*; oxidative stress; interaction, pneumoconiosis.

Introduction

Genetic factors that modify the response to oxidative environmental stimuli in humans are largely unknown. The overproduction of reactive oxygen species (ROS) due to major environmental factors such as cigarette smoking and chronic exposure to particles induces chronic airway inflammation [1,2,3]. Inflammation is fundamental to the pathogenesis of many chronic lung diseases including asthma, chronic obstructive pulmonary disease (COPD), and pneumoconiosis/fibrosis [4,5]. Coal workers' pneumoconiosis (CWP) is an inflammatory lung disease caused by chronic inhalation of coal dust, and ROS are thought to be important in disease pathogenesis. The source of ROS is proposed to derive directly from the coal dust, and indirectly from activated inflammatory cells such as alveolar macrophages and polymorphonuclear leukocytes [6].

The specific mechanisms of oxidative injury are becoming increasingly clear. ROS and specific pro-inflammatory cytokines such as tumor necrosis factor alpha (TNF- α) activate transcription factors including nuclear factor- κ B (NF- κ B) via specific intracellular signaling pathways. Translocation of transcription factors to the nucleus from the cytoplasm subsequently initiates transcription in genes with multiple inflammatory functions, including cytokines (TNF, IL-4, IL10, etc.), and antioxidant enzymes such as glutathione peroxidase, superoxide dismutases and catalase [7,8].

Interestingly, only some individuals similarly exposed to environmental oxidative stimuli (e.g. cigarette smoke and particles) develop lung disease, and the determinants of susceptibility are poorly understood. Intrinsic factors such as genetic background may be important determinants of inter-individual susceptibility to disease pathogenesis. A number of candidate genes for susceptibility to chronic pulmonary diseases have been investigated. The -308 G/A SNP in the promoter region of the gene (*TNF*) for *TNF- α* has been studied frequently. *TNF* is located in tandem with the lymphotoxin- α (*LTA*) gene (*LTA*) in the class III region of the 3.6 Mb human major histocompatibility complex (MHC) region on chromosome 6p21.31 [9], and the *TNF* gene superfamily plays an important role in inflammation. The -308 promoter polymorphism has been associated with subphenotypes of

asthma, COPD, fibrosis and CWP in some studies [10,11,12,13,14] but not in others [15,16]. Other genetic factors of CWP have also been studied [17]. Similarly, a functional *NcoI* RFLP polymorphism in *LTA* was associated with sarcoidosis in the study of Yamaguchi et al. [18] but it has not been replicated. Discordant results may be related to differences in population characteristics. Differential effects of the same environmental exposure on individuals with different genotypes may change the magnitude or the direction of the associations but unfortunately, none of the studies mentioned above have considered the effects of gene-environment interactions in the disease etiology or included biological markers of the response to oxidative stress.

The overall hypothesis of the present study was that *TNF* -308 and *LTA NcoI* polymorphisms modify the pulmonary responses to oxidants in coal miners differently exposed to cigarette smoke and coal mine dusts. To test this hypothesis, genotype-environment interactions on intermediate quantitative phenotypes of response to oxidative stress were specifically investigated. Further, genotype-intermediate phenotype interactions on disease phenotypes were studied. The response to oxidative stimuli (coal dust, cigarette smoke) was assessed by biological intermediate quantitative phenotypes (enzyme activities), and detailed information on coal workers' pneumoconiosis (assessed from computed tomography and twice by chest radiography 5 years apart). A major strength of this study is the quantitative assessment of environmental exposure to oxidative stimuli that permits evaluation of gene-environment interaction.

Subjects and Methods

Subjects studied took part in a longitudinal epidemiological study, including quantitative environmental measurements, biological assessment of response to oxidative stress and radiological assessment of coal workers' pneumoconiosis (CWP), including subclinical phenotypes.

Study sample

The population studied consisted of 259 unrelated coal miners recruited through a standardized protocol [19] in a French coal mine (Houillères du Bassin de Lorraine, in north east France) based on a selection contrasted by exposure and disease status. The sample included 240 active coal miners aged 34-50 years in 1990, including: 80 subjects heavily exposed to underground coal dusts (≥ 10 years at the coal face) with chest X-ray classified 0/1 or 1/0; 80 subjects exposed to underground coal dusts with normal chest X-ray classified 0/0; and 80 subjects slightly exposed with normal chest X-ray. Also included were 19 retired miners with CWP, *i.e.* chest X-ray classified 1/1 or over in 1994 and an evolution of two stages of X-ray grade within the past 10 years. The active coal miners were examined three times (1990, 1994, and 1999) and the retired miners twice (1994 and 1999). Most of the analysis has been performed on the active miners using 1994 and 1999 data. The retired miners with established disease have been included for detailed analysis in relation to disease. Out of the 240 active miners recruited in 1990, 234 took part in the 1994 survey and 227 in 1999. Out of the 19 retired miners recruited in 1994, 18 were reexamined in 1999. Based on available data, comparison of those lost to follow-up and the remainder did not show difference regarding exposure, genotype and disease.

Environment

Low or high current exposure based on job description and cumulative exposure assessed through job history and dust measurements at various sites of the mine were recorded. High current exposure refers to miners working at the coal face, mining, stope or

drift advance, and low exposure to those working at ventilation maintenance, pumping, haulage, shaft, stock equipment, or safety. Cumulative personal exposure to dust was estimated from each person's job history and from dust measurements at various sites of the coal mine according to the calculation previously described [20]. The estimates were summations of each dust measurement (mg/m^3) for the respective time spent in each job. Estimated cumulative exposures to dust were calculated until 1999 and expressed as $\text{mg}/\text{m}^3 \times \text{yr}$. Most of the analysis is based on high and low cumulative exposure using the same threshold ($>71\text{mg}/\text{m}^3 \times \text{yr}$) as previously [21]. Detailed smoking habits were recorded by standardized questionnaire.

Radiographic examination

Chest X-rays and computed tomography (CT) scans were performed to evaluate disease status. International Labour Office (ILO) classification in 12 profusion grades (0/-, 0/0, 0/1, 1/0, 1/1, 1/2, 2/1, 2/2, 3/1, 3/2, 3/3, 3/+) was used at the 3 examinations for all the miners [22]. Pneumoconiosis is defined by a grade $\geq 1/1$. Five categories were used in analyses: 0/- or 0/0, 0/1, 1/0, 1/1 or 1/2, and 2/1 or over. No subject was grade 3/1, 3/2 or 3/3 and only one subject was grade 3/+. CT scans were performed twice for all active miners at recruitment, *i.e.* 240 according to standardized methods [23,24] and interpreted by two experienced radiologists blinded to individual exposure and X-ray findings. Analyses were based on the profusion score already described [25] computed as the average over six lung zones (upper, middle, lower parts of both lungs) of profusion score rated from 0 to 3 (absence, rare, intermediate, high profusion). The study was approved by the appropriate ethical committee and written consent was obtained from all subjects.

Assessment of response to oxidative stimuli

Total plasma antioxidant concentration (ascorbate, protein thiols, bilirubin, urate and α tocopherol), plasma selenium concentration, $\text{Cu}^{++}/\text{Zn}^{++}$ superoxide dismutase ($\text{Cu}^{++}/\text{Zn}^{++}$ SOD), glutathione peroxidase (GSH-Px), and catalase activities were determined at the 1994

survey by methods previously described [21,26]. Activities were measured in the two weeks following storage and expressed as U/g Hb (Cu⁺⁺/Zn⁺⁺ SOD, GSH-Px) or k/g Hb (catalase) in erythrocytes, and U/L (GSH-Px) in plasma. All samples were analyzed in duplicate or triplicate and the precision (coefficient of variation) was <10%. Measures were available for 227 miners who did not differ regarding exposure or disease from those without.

Genotyping

Genomic DNA was isolated from 300 µL aliquots of stored blood utilizing the Wizard Genomic DNA Purification Kit from Promega Corporation (Madison, WI). The manufacturer's recommended protocol was modified slightly as follows: 1) centrifugation was performed at 14,000 x g; 2) incubation and centrifugation periods were doubled; 3) rehydration in TE was achieved with gentle rotation at room temperature for one week. Amplification for both polymorphisms was performed utilizing our previously optimized conditions, which include: 60 ng DNA, 1.5 mM MgCl₂, 0.4 µM forward and reverse primers, 200 µM dNTPs and 0.5 U Taq polymerase in a 25 µL reaction volume.

For determination of the A→G polymorphism at position -308 within *TNF*, we synthesized primers according to Zhu et al. [27]. Briefly, the forward primer for amplification was designed with a mismatch that inserts a *StyI* restriction site into the wild-type allele. Thirty-five cycles of amplification with an annealing temperature of 59°C were performed. PCR products were subsequently digested with *StyI* at 37°C for 2 hours. Fragments were resolved on 3% NuSieve/TBE gels stained with ethidium bromide. The resultant product sizes were 20 and 123 bp for wild-type alleles (allele 1), 143 bp for mutant alleles (allele 2), and 20, 123 and 143 bp for heterozygote alleles.

The *NcoI* RFLP within the first intron of *LTA* was analyzed using the primer design and temperature parameters of Stuber et al. [28]. Briefly, 37 cycles of 95°C denaturation for 30 seconds, 68°C annealing for 30 seconds, and 74°C extension for 42 seconds were performed. PCR products were digested at 37°C for 1 hour with *NcoI*. Fragments were resolved on 2% agarose gels stained with ethidium bromide. The resultant product sizes were

196 and 586 bp for *LTA* mutant alleles (allele 1), 782 bp for *LTA* wild-type alleles (allele 2), and 196, 586, and 782 bp for heterozygote alleles. Genotypes were determined for 231 miners. Comparisons of genotyped miners with those not ($n = 22$) did not show differences regarding exposure and disease.

Statistical methods

Standard statistical tests were used including χ^2 (or Fisher exact test when appropriate), analysis of variance, and multiple regression analysis. The interaction between genetic polymorphisms with exposure to oxidative stress and health indicators was statistically tested. Significance was assessed at the 5% two sided level. All analyses were performed with the SAS statistical software [29].

Results

Genotype distributions and environmental exposures to oxidative stress

Genotypes for the *TNF* -308 and *LTA NcoI* polymorphisms were successfully determined for 212 and 210 active coal miners respectively, and for the 19 retired miners with CWP. As expected, strong linkage disequilibrium was observed between *TNF* -308 and *LTA NcoI* genotypes (data not shown). No difference in genotype distributions was observed according to the geographic origin of the miners (France, other European countries and North Africa). Genotype distributions for the *TNF* -308 and *LTA NcoI* polymorphisms fit predictions for Hardy-Weinberg equilibrium.

The mean age of active miners was 47 years, and 55 years for retired miners. Half of the active miners were current smokers, 28% were currently exposed to coal dust and 4% had CWP in 1994. Thirteen percent (13%) of those reexamined 5 years later (1999) had CWP. In the small group of retired miners, none was currently exposed, approximately half were ex-smokers, and all had CWP in 1994. Age, body mass index, smoking, and current exposure to dust were unrelated to *TNF* -308 and *LTA NcoI* genotypes (Table 1). Cumulative dust exposure was also unrelated to *TNF* -308 genotype. A relation of borderline significance between individuals homozygous or heterozygous for *LTA NcoI* allele 1 with cumulative exposure was observed, which remained after adjustment for age ($p = 0.07$).

Stages of disease

TNF -308 genotypes were unrelated to 1994 X-ray grade or to the CT-scan micronodule score with 1999 X-ray grade (Table 2). In contrast, CT-scan micronodule score significantly varied according to *LTA NcoI* genotypes, even though *LTA NcoI* genotype was unrelated to 1994 X-ray grade (Table 2). A statistically significant relation of *LTA NcoI* genotypes with 1999 X-ray grade was found. In 1999, individuals homozygous for *LTA NcoI* allele 1 had the lowest prevalence of normal X-ray grade (35.3%) and individuals with X-ray grade 2/1 or over were only found in individuals homozygous for allele 2 both in 1994 and in 1999. The single subject with 3/+ X-ray grade was heterozygous for *TNF* -308 and *LTA NcoI*.

Markers of response to oxidative stress

Overall, individuals homozygous or heterozygous for *TNF* -308 allele 2 had significantly lower GSH-Px activity and higher erythrocyte catalase activity (association was statistically significant after exclusion of two outliers) (Table 3). No association was found between biological markers and *LTA NcoI* genotypes.

Gene-environment interaction on GSH-Px activity.

We previously found, with the same study sample, a borderline significant association of erythrocyte GSH-Px activity and a significant association of catalase activity with occupational exposure to coal dust [21]. In the present study, erythrocyte GSH-Px activity was significantly lower in individuals homozygous or heterozygous for *TNF* -308 allele 2, and we hypothesized that environmental exposure to oxidants (cigarette smoke, dust exposure) interacts with *TNF* -308 genotype to determine the capacity of GSH-Px to respond to oxidative stress. Therefore, we studied the association of erythrocyte GSH-Px activity with smoking habits and occupational exposure to dust according to *TNF* -308 genotypes.

No significant interaction was found between *TNF* -308 genotypes with smoking habits on erythrocyte GSH-Px activity. Nevertheless, among the miners homozygous or heterozygous for *TNF* -308 allele 2, mean (\pm SD) erythrocyte GSH-Px activity was negatively and significantly related to smoking habits (44.2 ± 10.8 U/g Hb in non smokers, 36.7 ± 12.6 U/g Hb in ex-smokers and 34.7 ± 11.4 U/g Hb in smokers, $p = 0.05$ trend test). Among the miners homozygous for *TNF* -308 allele 1, mean GSH-Px activity was unrelated to smoking habits (43.3 ± 12.7 U/g Hb in non smokers, 39.1 ± 15.6 U/g Hb in ex-smokers and 42.5 ± 12.6 U/g Hb in smokers, $p = 0.27$). Among miners genotyped for *TNF* -308, a borderline significant association was found between smoking habits and mean estimated cumulative exposure to dust (58.9 ± 46.0 mg/m³ x yr in non smokers, 66.5 ± 45.8 mg/m³ x yr in ex-smokers and 51.1 ± 35.9 mg/m³ x yr in smokers, $p = 0.06$).

A significant interaction between cumulative dust exposure and *TNF* -308 genotypes was found (Figure 1). In miners with low cumulative dust exposure, no significant difference was

found in erythrocyte GSH-Px activity between miners homozygous or heterozygous for *TNF* -308 allele 2 and those homozygous for *TNF* -308 allele 1. In contrast, erythrocyte GSH-Px activity was significantly lower in miners homozygous or heterozygous for *TNF* -308 allele 2 and with high cumulative dust exposure. A multivariate linear regression model of GSH-Px activity (dependent variable) on cumulative exposure (low vs. high) and *TNF* -308 (11 vs. 12 or 22) was used to test statistically the interaction (i.e. the heterogeneity of the association of GSH-Px activity to *TNF* -308) of exposure x *TNF* -308, and found a p value of 0.06 (0.05 in a model also including smoking) for the interaction. In those with high cumulative dust exposure, differences according to *TNF* -308 genotypes were similar when considering separately those with and without CWP in 1994. In miners without CWP, mean (\pm SEM) GSH-Px activity was significantly lower in miners homozygous or heterozygous for *TNF* -308 allele 2 (36.2 ± 2.8 ; n = 17) compared with those homozygous for allele 1 (44.6 ± 2.0 ; n = 43); p = 0.02). In the small group of miners with CWP, mean GSH-Px activity was also lower in miners homozygous or heterozygous for *TNF* -308 allele 2 (23.4 ± 5.7 ; n = 3) vs. those homozygous for allele 1 (52.1 ± 11.4 ; n = 3; p = 0.09). Five-year CWP incidence between *TNF* -308 genotypes in miners with low cumulative dust exposure was similar in miners homozygous for *TNF* -308 allele 1 and in those homozygous or heterozygous for *TNF* -308 allele 2, whereas in those with high cumulative dust exposure, more than two fold (although not statistically significant) higher CWP incidence was observed. No interaction was found between *LTA NcoI* genotypes with environmental exposure to oxidants on erythrocyte GSH-Px or catalase activities.

Gene-intermediate phenotype interaction on prevalence of CWP

We also previously showed [21] that erythrocyte catalase activity was positively and significantly associated with 1994 X-ray grade. We therefore hypothesized that an interaction exists between *LTA NcoI* genotypes and erythrocyte catalase activity on disease phenotypes. To test this hypothesis, we studied the association of *LTA NcoI* genotypes with CT-scan score or CWP prevalence according to erythrocyte catalase activity. Miners were classified

according to low or high catalase activity, with a threshold based on the third tertile. Due to the small number of miners with CWP homozygous for the *LTA NcoI* allele 1, miners homozygous or heterozygous for the *LTA NcoI* allele 1 were combined and compared with miners homozygous for *LTA NcoI* allele 2.

Significant interactions between catalase activity and *LTA NcoI* genotypes on CWP prevalence in 1994 and 1999 were found (Figure 2). In miners with low catalase activity, the CWP prevalence was significantly higher in miners homozygous for the *LTA NcoI* allele 2 whereas in miners with high catalase activity, the CWP prevalence was lower in miners homozygous for the *LTA NcoI* allele 2. At both surveys, odds ratios were greater than 1 in those with low catalase activity, i.e. *LTA NcoI* 22 genotype increased the risk of pneumoconiosis, but odds ratios were lower than 1 in those with high catalase activity. No interaction was found between *LTA NcoI* genotypes with catalase activity on CT-scan score, or between *TNF* -308 genotypes with catalase activity on disease phenotypes.

Discussion

In this genetic epidemiologic follow-up study, we tested the general hypothesis that *TNF* -308 and *LTA NcoI* polymorphisms modify pulmonary responses to oxidants in coal miners differently exposed to cigarette smoke and coal mine dusts. Results indicated significant gene-environment interaction on intermediate phenotypes (response to oxidative stimuli), and gene-intermediate phenotype interaction on clinical disease (coal worker's pneumoconiosis). An interaction of *TNF* -308 genotype with quantitatively assessed occupational exposure on erythrocyte GSH-Px activity was observed, with a significant association in those with high exposure whereas no association was found among those with low exposure. Further, assessment of the interaction of gene to biological response to oxidative stress on clinical phenotype identified a significant association of CWP prevalence with *LTA Nco I* in those with low catalase activity whereas no association was observed in those with high (*a priori* protective) activity.

Assessing gene-environment interaction has proven to be difficult, and few clear examples have been evidenced and replicated. Problems often encountered are the lack of specific hypotheses (candidate interactions), insufficient characterization of exposure, and the availability of sufficiently low level (closer to the gene than clinical disease) phenotype markers. Unraveling biochemical and physiological hierarchies leading from genes to clinical endpoints in complex diseases need new strategies [30]. This study attempted to incorporate a conceptual temporal sequence [31] from environment and genes towards disease. In such an overall framework of etiological factors to severe disease, it is possible to distinguish 1) internal dose (inhaled coal dust), 2) early biologic effect (low-level intermediate phenotype [30], such as enzymatic response to oxidative stimuli, 3) altered structure/function (CT score), and 4) clinical (X-ray confirmed CWP) and severe (nodular form) disease. We presently attempted to apply this framework to study gene-environment interaction and gene-intermediate phenotype interaction in the pathogenesis of CWP.

Strengths of the study were the contrasted exposure by design in the study sample, the availability of quantitative objective measures of the coal mine dust, the availability of various quantitative phenotypes to assess response to oxidative stress, subclinical computed tomography score, and two assessments of CWP by standardized X-ray five years apart. Limitations of the study were the relatively small sample size that precludes detailed analyses addressing simultaneously the various steps of exposure, as well as intermediate and clinical phenotypes. Another limitation is the particular type of the population studied (coal miners), but it does permit study of a homogeneous group of subjects. The window of opportunity for evaluation of the role of single or multiple genes in disease pathology may depend on the relative timing of the environmental stimuli, but also the stage of the disease. Furthermore, genetic background may play a key role in the occurrence and/or the severity of the disease [32,33]. In the present study, it was not possible to address the role of *TNF* and *LTA* in the pathogenesis from established CWP towards the most severe forms, but continued follow-up of the study sample will allow us to address that question later.

Another important consideration in a study of gene-environment interaction and disease pathogenesis is the complex quantitative nature of the disease under study. Almost certainly, multiple genes contribute to CWP susceptibility. However, due to the small size of the study population, the investigation was limited to those genes with relatively common polymorphisms with known roles in chronic pulmonary diseases (see below). That is, the study population does not have sufficient statistical strength to evaluate potentially important rare gene polymorphisms. The *TNF* -308 and *LTA NcoI* polymorphisms are relatively common in European populations (approx 14% and 30%, respectively, in the present study), and each has demonstrated roles in chronic pulmonary diseases and responses to oxidant stimuli. For example, the *TNF* -308 polymorphism has been associated with chronic bronchitis [10], fibrosing alveolitis [11], asthma [34], and a higher risk to poor prognosis in COPD [35]. *TNF- α* has also been demonstrated to be an important determinant of susceptibility to lung inflammation and injury induced by the environmental oxidant ozone (O_3) in animal models [36]. Similarly, Shimura et al. [37] reported that *LTA NcoI* allele 2 was less prevalent in 135 lung cancer patients as compared to 165 normal healthy subjects. The

authors also reported that among patients with stages III and IV lung cancer (n = 31), those homozygous for the *LTA NcoI* allele 1 (n = 9) had a two fold greater survival rate 5 years after operation than in patients homozygous or heterozygous for the *LTA NcoI* allele 2, whereas no difference was found among patients with stages I and II. Campbell et al. [38] found a significant protection of individuals homozygous for *LTA NcoI* allele 1 against recurrent wheeze and a borderline protection against asthma in a random cohort of 224 subjects. Further, Stuber et al. [28] reported a significant association of *LTA NcoI* allele 2 with worse outcome among 40 patients with severe sepsis. On the contrary, Yamaguchi et al. [18] found that *LTA NcoI* allele 1 is associated with a prolonged clinical course of sarcoidosis in 110 Japanese patients and other studies reported no association with disease [34,39]. Ethnic difference or the fact that some of these patients showed mild to moderate disease manifestations may partly explain these discrepancies.

Several arguments support the hypothesis that markers of the response to oxidative stress are relevant intermediate phenotypes for CWP. It is well known that CWP is related to increased amounts of hydrogen peroxide [40,41] and catalase converts hydrogen peroxide to water and oxygen. We previously found that catalase activity was related to pneumoconiosis and was higher in miners with a CT micronodule score ≥ 4 [21]. In a 5-year follow-up study conducted in 66 coal miners, incidence or disease progression was associated with the combination of initial levels of various antioxidant enzyme activities [42]. Likewise, a potential role of *TNF* genotype on GSH-Px response to oxidative stimuli is consistent with previous studies. *TNF- α* is known to generate ROS by leakage from the mitochondrial electron transport chain [43], and modulates glutathione levels (GSH), one key component for the GSH-Px activity. Moreover the gene for glutathione peroxidase is induced in response to *TNF- α* in lung cells [44]. This may have important implications for the oxidant/antioxidant balance in inflammatory lung diseases. Overall, due to the systemic nature of the response to oxidative stress studied, the framework applied in this study is of general interest for other diseases that depend on the same pathways.

The present study demonstrated an interaction of the *TNF* -308 genotype with high coal dust exposure on GSH-Px activity. As the less common allele 2 is associated with higher basal and induced expression of TNF- α [45], and TNF- α and ROS cooperate to activate transcription factors [46], we may postulate that TNF- α modulates the GSH-Px activity through the regulation of the GSH levels. In the presence of chronic oxidative stress due to high cumulative dust exposure, antioxidant defenses may be overwhelmed leading to alteration of the antioxidant enzymes as lower erythrocyte Cu⁺⁺/Zn⁺⁺ SOD activity, as we observed previously [21], and down regulation of GSH-Px [47]. Erythrocyte GSH-Px activity plays a major role in ROS metabolism in most tissues and has been shown to control NF- κ B transcription factor activation [48]. As plasma GSH-Px originates from the kidney [49], it plays mostly a role in response to acute oxidative stress which is consistent with our observation that only erythrocyte GSH-Px response depends on cumulative exposure.

Our results support and extend findings from a Belgian study in which *TNF* -308 allele 2 was more prevalent in 18 ex-coal miners with X-ray grade 0/1 or over (14 of whom with X-ray 1/1 or over) as compared to 60 ex-coal miners with normal X-ray (0/0) and 56 controls [13]. Previously, they reported that spontaneous and coal-dust-stimulated TNF release from peripheral blood monocytes was increased in coal miners with early CWP [50]. Furthermore, they found that miners who showed abnormally high release of TNF stimulated by coal dust had an increased risk of progression in CWP over 5 years [51]. Similarly, *TNF* -308 allele 2 was found to be more prevalent in 80 patients who had CWP than in 54 healthy controls, with a two times higher prevalence in those with large opacities as compared to those with simple CWP [14]. An autopsy study, in which the severity of fibrosis was anatomically assessed, showed a significant prevalence of allele 2 restricted to moderate fibrosis in 325 former coal miners as compared to 164 controls [52]. In contrast, the association of allele 2 with disease was restricted to severe silicosis (9 cases) in 241 South African gold miners [33]. In none of these studies was the analysis stratified according to dust exposure to assess gene-environment interaction. Limited information is available on association of *TNF* with the first stages of the disease. No association in the present study between *TNF* -308 genotypes and CT-score was found, but it is worth noting that although it was not statistically significant, a

two fold increase in five-year CWP incidence was observed in miners homozygous or heterozygous for *TNF* -308 allele 2 with high cumulative dust exposure.

Overall, our findings and previous results on the severity of various diseases (in particular, silicosis) implicate *TNF* -308 at various stages of disease progression, but further work is needed to clarify mechanisms and interaction with other genes. Another important consideration is that CWP, similar to other complex diseases, likely involves numerous genes acting at each step. While good rationale is provided for the study of *TNF* and *LTA*, other genes implicated in the development of oxidative stress such as antioxidant enzyme genes and transcription factors that modulate their expression may/should be investigated. Further, well-designed studies that take advantage of emerging expression array and proteomics technologies in human subjects and animal models exposed to coal dust, or oxidant mixtures, should yield insight to other important candidate susceptibility genes.

In conclusion, this is the first demonstration of the involvement of functional polymorphisms of two candidate genes *TNF* and *LTA* in the control of the physiological/biochemical responses to quantitatively assessed oxidative stimuli. The study design described in this study enabled investigation of environment- and disease-interactions, and provides a better understanding of the etiology of the chronic inflammatory lung diseases such as pneumoconiosis.

Acknowledgments

The paper is dedicated to the late Quang-Thieu Pham who set up the coal miner study. We thank the medical staff and all the French coal mine workers who made this study possible. We also thank A. Berthelin, M. Dusch and G. Hellier for their skilful technical assistance. This research was in part, supported by grant numbers 7280.03.001 and 7280.03.035 from the European Community of Steel and Coal, National Institutes of Health grant ES-09606, and Environmental Protection Agency grant R-825815.

References

1. Malech HL, Gallin JI. Neutrophils in human diseases. *N Engl J Med* 1987;317:687-694
2. Pryor WA, Stone K. Oxidants in cigarette smoke: radicals, hydrogen peroxides, peroxyxynitrate, and peroxyxynitrite. *Ann N Y Acad Sci* 1993;686:12-28
3. Petit JC, Boettner JC. Evidence of H₂O₂ formation in the early stage of coal oxidation. In: Charcosset H (ed) *Advanced methodologies in coal characterisation*. Elsevier, 1990;253-266.
4. Fujimura N. Pathology and pathophysiology of pneumoconiosis. *Curr Opin Pulm Med* 2000;6:140-144
5. Jeffery PK. Remodeling in asthma and chronic obstructive lung disease. *Am J Respir Crit Care Med* 2001;164:S28-S38
6. Schins RPF, Borm PJA. Mechanisms and mediators in coal dust induced toxicity: a review. *Ann Occup Hyg* 1999;43:7-33.
7. Barnes PJ, Karin M. Nuclear factor- κ B—a pivotal transcription factor in chronic inflammatory diseases. *N Engl J Med* 1997;336:1066-1071
8. Rahman I, MacNee W. Regulation of redox glutathione levels and gene transcription in lung inflammation: therapeutic approaches. *Free Radic Biol Med* 2000;28:1405-1420
9. Carroll MC, Katzman P, Alicot EM, Koller BH, Geraghty DE, Orr HT, Strominger JL, Spies T. Linkage map of the human major histocompatibility complex including the tumour necrosis factor genes. *Proc Natl Acad Sci USA* 1987;84:8535-8539
10. Huang SL, Su CH, Chang SC. Tumor necrosis factor- α gene polymorphism in chronic bronchitis. *Am J Respir Crit Care Med* 1997;156:1436-1439
11. Whyte M, Hubbard R, Meliconi R, Whidborne M, Eaton V, Bingle C, Timms J, Duff G, Facchini A, Pacilli A, Fabbri M, Hall I, Britton J, Johnston I, Di Giovine F. Increased risk of fibrosing alveolitis associated with interleukin-1 receptor antagonist

- and tumor necrosis factor-alpha gene polymorphisms. *Am J Respir Crit Care Med* 2000;162:755-758
12. Winchester EC, Millwood IY, Rand L, Penny MA, Kessler AM. Association of the TNF- α -308 (G \rightarrow A) polymorphism with self-reported history of childhood asthma. *Hum Genet* 2000;107:591-596
 13. Zhai R, Jetten M, Schins RPF, Franssen H, Borm PJA. Polymorphisms in the promoter of the tumor necrosis factor- α gene in coal miners. *Am J Ind Med* 1998;34:318-324
 14. Kim KA, Cho YY, Cho JS, Yang KH, Lee WK, Lee KH, Kim YS, Lim Y. Tumor necrosis factor-alpha gene promoter polymorphism in coal workers' pneumoconiosis. *Mol Cell Biochem* 2002;234-235:205-209
 15. Higham MA, Pride NB, Alikhan A, Morrell NW. Tumor necrosis factor- α gene promoter polymorphism in chronic obstructive pulmonary disease. *Eur Respir J* 2000;15:281-284
 16. Pantelidis P, Fanning GC, Wells AU, Welsh KI, Du Bois RM. Analysis of tumor necrosis factor-alpha, lymphotoxin-alpha, tumor necrosis factor receptor II, and interleukin-6 polymorphisms in patients with idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2001;163:1432-1436
 17. Zhai R, Liu G, Ge X, Yang C, Huang C, Wu C, Christiani DC. Genetic polymorphisms of MnSOD, GSTM1, GSTT1, and OGG1 in coal workers' pneumoconiosis. *J Occup Environ Med* 2002;44:372-377
 18. Yamaguchi E, Itoh A, Hizawa N, Kawakami Y. The gene polymorphism of tumor necrosis factor- β , but not that of tumor necrosis factor- α , is associated with the prognosis of sarcoidosis. *Chest* 2001;119:753-761
 19. Pham QT, Bourgard E, Chau N, Willim G, Megherbi SE, Teculescu D, Bohadana A, Bertrand JP. Forced oscillation technique (FOT): a new tool for epidemiology of occupational lung diseases? *Eur Respir J* 1995;8:1307-1313
 20. Attfield MD, Moring K. The derivation of estimated dust exposures for U.S. coal miners working before 1970. *Am Ind Hyg Assoc J* 1992;53:248-255

21. Nadif R, Bourgkard E, Dusch M, Bernadac P, Bertrand JP, Mur JM, Pham QT. Relations between occupational exposure to coal mine dusts, erythrocyte catalase and Cu⁺⁺/Zn⁺⁺ superoxide dismutase activities, and the severity of coal worker's pneumoconiosis. *Occup Environ Med* 1998;55:533-540
22. International Labour Office (ILO). Guideline for the use of ILO international classification of radiographs of pneumoconiosis. ILO division of Occupational Safety and Health Sciences, Geneva, 1980;publication 22
23. Bernadac P, Hennequin L, Bertal A, Jaafar S. Radiologie et pneumoconioses. *Encycl Med Chir Radiodiagnostic - Cœur – Poumon* 1997;32-450-A-10:20p. Elsevier, Paris
24. Kleinerman J, Green F, Harley RA, Lapp L, Laqueur W, Naeye RL, Pratt P, Taylor G, Wiot J, Wyatt J. Pathology standards for coal worker's pneumoconiosis: report of the pneumoconiosis committee of the College of American Pathologists to the National Institute for Occupational Safety and Health. *Arch Pathol Lab Med* 1979;103:375-432
25. Bourgkard E, Bernadac P, Chau N, Bertrand JP, Teculescu D, Pham QT. Can the evolution to pneumoconiosis be suspected in coal miners ? A longitudinal study. *Am J Respir Crit Care Med* 1998;158:504-509
26. Nadif R, Oryszczyn MP, Fradier-Dusch M, Hellier G, Bertrand JP, Pham QT, Kauffmann F. Cross sectional and longitudinal study on selenium, glutathione peroxidase, smoking, and occupational exposure in coal miners. *Occup Environ Med* 2001;58:239-245
27. Zhu S, Chan-Yeung M, Becker AB, Dimich-Ward H, Ferguson AC, Manfreda J, Watson WTA, Pare PD, Sandford AJ. Polymorphism of the IL-4, TNF α , and Fc ϵ RI β genes and the risk of allergic disorders in at-risk infants. *Am J Respir Crit Care Med* 2000;161:1655-1659
28. Stuber F, Petersen M, Bokelman F, Schade U. A genomic polymorphism within the tumor necrosis factor locus influences plasma tumor necrosis factor-alpha concentrations and outcome of patients with severe sepsis. *Crit Care Med* 1996;24:373-375
29. SAS Institute. SAS release 6.12, 1997 Cary, NC

30. Schork NJ. Genetics of complex disease. Approaches, problems and solutions. *Am J Respir Crit Care Med* 1997;156:S103-S109
31. Schulte PA. A conceptual framework for the validation and use of biologic markers. *Environ Res* 1989;48:129-144
32. Pin I, Siroux V, Cans C, Kauffmann F, Maccario J, Pison C, Dizier MH. Familial resemblance of asthma severity in the EGEA study. *Am J Respir Crit Care Med* 2002;165:185-189
33. Corbett EL, Mozzato-Chamay N, Butterworth AE, De Cock KM, Williams BG, Churchyard GJ, Conway DJ. Polymorphisms in the tumor necrosis factor- α gene promoter may predispose to severe silicosis in black south African miners. *Am J Respir Crit Care Med* 2002;165:690-693
34. Witte JS, Palmer LJ, O'Connor RD, Hopkins PJ, Hall LM. Relation between tumour necrosis factor polymorphism TNFalpha-308 and risk of asthma. *Eur J Hum Genet* 2002;10:82-85
35. Keatings VM, Cave SJ, Henry MJ, Morgan K, O'Connor CM, FitzGerald MX, Kalsheker N. A polymorphism in the tumor necrosis factor-alpha gene promoter region may predispose to a poor prognosis in COPD. *Chest* 2000;118:971-975
36. Kleeberger SR, Levitt RC, Zhang L, Longphre M, Harkema J, Jedlicka A, Eleff SM, DiSilvestre D, Holroyd KJ. Linkage analysis of susceptibility to ozone-induced lung inflammation in inbred mice. *Nat Genet* 1997;17:475-478
37. Shimura T, Hagihara M, Takebe K, Munkhbat B, Odaka T, Kato H, Nagamachi Y, Tsuji K. The study of tumor necrosis factor beta gene polymorphism in lung cancer patients. *Cancer* 1994;73:1184-1188
38. Campbell DA, Li Kam Wa E, Britton J, Holgate ST, Markham AF, Morrison JFJ. Polymorphism at the tumour necrosis factor locus and asthma. In: Hall IP (ed) *Genetics of asthma and atopy*. Monogr Allergy. Karger, Basel, 1996;33:125-137
39. Patuzzo C, Gile LS, Zorzetto M, Trabetto E, Malerba G, Pignatti PF, Luisetti M. Tumor necrosis factor gene complex in COPD and disseminated bronchiectasis. *Chest* 2000;117:1353-1358

40. Rom WN, Bitterman PB, Rennard SI, Cantin A, Crystal RG. Characterization of the lower respiratory tract inflammation of nonsmoking individuals with interstitial lung disease associated with chronic inhalation of inorganic dusts. *Am Rev Respir Dis* 1987;136:1429-1434
41. Dalal NS, Suryan MM, Vallyathan V, Green FHY, Jafari B, Wheeler R. Detection of reactive free radicals in fresh coal mine dust and their implication for pulmonary injury. *Ann Occup Hyg* 1989;33:79-84
42. Schins RPF, Keman S, Borm PJA. Blood antioxidant status in coal dust-induced respiratory disorders: a longitudinal evaluation of multiple biomarkers. *Biomarkers* 1997;2:45-50
43. Schmidt KN, Amstad P, Cerruti P, Baeuerle PA. The roles of hydrogen peroxide and superoxide as messengers in the activation of transcription factor NF-kappa B. *Chem Biol* 1995;2:13-22
44. Rahman I, Antonicelli F, MacNee W. Molecular mechanism of the regulation of glutathione synthesis by tumor necrosis factor- α and dexamethasone in human alveolar epithelial cells. *J Biol Chem* 1999;274:5088-5096
45. Wilson AG, Symons JA, McDowell TL, McDewitt HO, Duff GW. Effects of a polymorphism in the human tumor necrosis factor alpha promoter on transcription activation. *Proc Natl Acad Sci USA* 1997;94:3195-3199
46. Janssen-Heininger YMW, Macara I, Mossman T. Cooperativity between oxidants and tumor necrosis factor in the activation of nuclear factor (NF)- κ B. Requirement of Ras/mitogen-activated protein kinases in the activation of NF- κ B by oxidants. *Am J Respir Cell Mol Biol* 1999;20:942-952
47. Fujii J, Taniguchi N. Down regulation of superoxide dismutases and glutathione peroxidase by reactive oxygen and nitrogen species. *Free Rad Res* 1999;31:301-308
48. Brigelius-Flohé R, Maurer S, Lotzer K, Bol GB, Kallionpaa H, Letholainen P, Viita H, Yla-Herttuala S. Overexpression of PHGPx inhibits hydroperoxide-induced oxidation, NF κ B activation and apoptosis and affects oxLDL-mediated proliferation of rabbit aortic smooth muscle cells. *Atherosclerosis* 2000;152:307-316

49. Harris ED. Regulation of antioxidant enzymes. *Faseb J* 1992;6:2675-2683
50. Borm PJA, Palmén N, Engelen JJM, Buurman WA. Spontaneous and stimulated release of tumor necrosis factor- α (TNF) from blood monocytes of miners with coal workers' pneumoconiosis. *Am Rev Respir Dis* 1988;138:1589-1594
51. Schins RPF, Borm PJA. Epidemiological evaluation of release of monocyte TNF- α as an exposure and effect marker in pneumoconiosis: a five year follow up study of coal workers. *Occup Environ Med* 1995;52:441-450
52. Yucesoy B, Vallyathan V, Landsittel DP, Sharp DS, Weston A, Burleson GR, Simeonova P, McKinstry M, Luster MI. Association of tumor necrosis factor- α and interleukine-1 gene polymorphisms with silicosis. *Toxicol Appl Pharmacol* 2001;172:75-82

Table 1. *TNF* -308 and *LTA NcoI* polymorphisms and environmental exposure to oxidative stress in coal miners.

	<i>TNF</i> -308					<i>LTA NcoI</i>					P Value
	11	12	22	12 or 22	P Value ^a	11	12	22	P Value		
n	158	52	2	54		18	91	101			
Age (years, mean \pm SD)	46.5 \pm 3.5	47.1 \pm 3.5	46.5 \pm 2.1	47.1 \pm 3.5	0.3	46.1 \pm 3.3	47.2 \pm 3.4	46.4 \pm 3.5	0.16		
Body mass index (mean \pm SD)	27.1 \pm 3.7	27.6 \pm 3.1	29.2 \pm 3.6	27.6 \pm 3.1	0.3	27.6 \pm 3.1	27.2 \pm 3.1	27.1 \pm 4.0	0.9		
	26.8 \pm 3.2 ^b				0.12			26.8 \pm 3.2 ^b	0.5		
Smoking habits (%):											
Non smokers	20.3	17.3	50.0	18.5		11.1	24.2	17.8			
Ex-smokers	29.1	32.7	0	31.5	0.9	27.8	31.9	28.7	0.5		
Current smokers	50.6	50.0	50.0	50.0		61.1	43.9	53.5			
Pack-years (mean \pm SD)	17.0 \pm 14.9	17.7 \pm 15.2	0.1 \pm .14	17.0 \pm 15.3	1.00	14.4 \pm 9.3	17.0 \pm 17.7	17.4 \pm 13.1	0.7		
Current exposure (%):											
None (retirement)	24.7	21.1	50.0	22.2		16.7	26.4	23.8			
Low	48.7	48.1	50.0	48.2	0.9	55.6	47.2	48.5	0.9		
High	26.6	30.8	0	29.6		27.8	26.4	27.7			
Cumulative dust exposure (mg/m ³ x yr, mean \pm SD)	55.3 \pm 38.8	63.4 \pm 48.9	49.0 \pm 48.1	62.9 \pm 48.5	0.3	73.6 \pm 35.6	60.7 \pm 49.9	51.9 \pm 32.7	0.08		

^a: 12 or 22 vs. 11. ^b: excluding two outliers (43.6 and 45.7 kg/m²).

Table 2. *TNF* -308 and *LTA NcoI* polymorphisms and stages of pneumoconiosis in coal miners.

	<i>TNF</i> -308				<i>LTA NcoI</i>				P Value
	11	12	22	12 or 22	11	12	22	P Value	
1994 X-ray grade, n	158	52	2	54	18	91	101		
0/0, %	72.8	69.2	50.0	68.5	55.6	69.2	76.2		
0/1, %	15.2	13.5	50.0	14.8	22.2	17.6	11.9		
1/0, %	8.2	11.5	0.0	11.1	16.7	11.0	5.9	0.8	0.3
1/1 or 1/2, %	3.2	5.8	0.0	5.6	5.5	2.2	5.0		
2/1 or over, %	0.6	0.0	0.0	0.0	0.0	0.0	1.0		
1994 CT-scan micronodule score, n	157	52	2	54	18	90	101		
(mean \pm SD)	2.15 \pm 2.84	1.67 \pm 2.09	1.00 \pm 1.41	1.65 \pm 2.07	2.78 \pm 2.56	1.49 \pm 1.91	2.39 \pm 3.17	0.16	0.03
1999 X-ray grade, n	154	49	1	50	17	86	99		
0/0, %	68.2	61.2	100	62.0	35.3	65.1	72.7		
0/1, %	8.4	8.2	0.0	8.0	17.7	10.5	5.1		
1/0, %	10.4	12.2	0.0	12.0	17.6	15.1	6.1	0.9	0.007
1/1 or 1/2, %	11.0	16.3	0.0	16.0	29.4	9.3	12.1		
2/1 or over, %	2.0	2.1	0.0	2.0	0.0	0.0	4.0		

^a: 12 or 22 vs. 11.

Table 3. *TNF* -308 and *LTA NcoI* polymorphisms and biological markers of response to oxidative stress in coal miners in 1994.

	<i>TNF</i> -308			<i>LTA NcoI</i>			P Value
	11	12	22	11	12	22	
n	149	52	1	16	89	95	
In erythrocytes:							
Catalase (k/g Hb)	95 ± 32 92 ± 23 ^b	102 ± 23	116	99 ± 15	101 ± 37 97 ± 25 ^b	92 ± 23	0.09 0.2
Cu ⁺⁺ /Zn ⁺⁺ SOD (U/g Hb)	145 ± 31	147 ± 34	120	149 ± 40	148 ± 32	143 ± 31	0.4
GSH-Px (U/g Hb)	42.0 ± 13.6	36.5 ± 12.2	36.9	41.3 ± 10.9	39.7 ± 14.0	41.1 ± 13.5	0.8
In plasma:							
GSH-Px (U/L)	716 ± 226	681 ± 215	1202	760 ± 209	708 ± 234	697 ± 218	0.6
Total antioxidant concentration (mmol/L)	1.36 ± 0.20 1.35 ± 0.19 ^c	1.34 ± 0.16	1.54	1.40 ± 0.15	1.34 ± 0.17	1.36 ± 0.21 1.35 ± 0.19 ^c	0.4 0.4
Se concentration (ng/L)	66.3 ± 19.3	61.2 ± 18.8	69.8	67.5 ± 23.5	64.5 ± 18.5	64.7 ± 19.3	0.8

Results are expressed as mean ± SD.

^a: 12 or 22 vs. 11.

^b: excluding two outliers. Catalase activities were 254 and 303 k/g Hb and were aged 47 and 49 yrs. Both outliers had normal X-rays at both surveys and low current exposure; one was a smoker, one an ex-smoker.


^c: excluding one outlier (2.29 mmol/L, 47 yrs, low current exposure, normal X-ray both surveys, ex-smoker).

Figure 1 . Associations of *TNF* –308 genotypes to quantitative phenotype of oxidative stress and pneumoconiosis incidence according to cumulative occupational exposure.


Cumulative coal dust exposure. Low: $\leq 71 \text{ mg.m}^3 \times \text{yr}$; High: $> 71 \text{ mg.m}^3 \times \text{yr}$ based on job history and ambient measurements in the coal mine. Numbers of miner (all active in 1994) in each group are figured below each bar. Five-year incidence refers to the proportion of those with CWP in 1999 among those without in 1994. OR: odds ratio [95% confidence interval].

Figure 2. Associations of *LTA NcoI* genotypes to pneumoconiosis prevalence according to the level of quantitative phenotype of oxidative stress

Erythrocyte catalase activity in 1994. Low: $\leq 107 \text{ k/g Hb}$; high: $> 107 \text{ k/g Hb}$ (highest tertile). Numbers of miners (either active or retired in 1994) in each group are figured below each bar. OR : odds ratio [95% confidence interval]. Breslow and Day test for interaction refers to the heterogeneity of odds ratios according to catalase activity.



Nadif et al. Figure 1.



Nadif et al. Figure 2.