



## Metabolic syndrome in a sub-Saharan African setting: Central obesity may be the key determinant.

Leopold Fezeu, Beverley Balkau, André-Pascal Kengne, Eugène Sobngwi,  
Jean-Claude Mbanya

### ► To cite this version:

Leopold Fezeu, Beverley Balkau, André-Pascal Kengne, Eugène Sobngwi, Jean-Claude Mbanya.  
Metabolic syndrome in a sub-Saharan African setting: Central obesity may be the key determinant.: Metabolic syndrome in Sub-Saharan Africa. Atherosclerosis, 2007, 193 (1), pp.70-76.  
10.1016/j.atherosclerosis.2006.08.037 . inserm-00128606

HAL Id: inserm-00128606

<https://inserm.hal.science/inserm-00128606>

Submitted on 6 Jun 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Table 1: Age adjusted, gender specific rural - urban characteristics of the study population

Clinical and biological variables	WOMEN			MEN		
	Rural	Urban	p	Rural	Urban	P
N	374	513		252	414	
Age (years)	46 (13)	37 (9)	< 0.001	46 (14)	38 (9)	< 0.001
Anthropometric parameters						
Body mass index (kg/m <sup>2</sup> )	22.1 (0.2)	26.4 (0.2)	< 0.001	21.7 (0.2)	25.1 (0.2)	< 0.001
Waist circumference (cm)	81.2 (0.5)	81.3 (0.4)	0.9	79.6 (0.5)	84.4 (0.4)	< 0.001
Hip circumference (cm)	91.2 (0.5)	103.8 (0.4)	< 0.001	89.2 (0.5)	98.7 (0.4)	< 0.001
Waist hip ratio	0.90 (0.004)	0.78 (0.004)	< 0.001	0.90 (0.004)	0.85 (0.003)	< 0.001
Blood Pressure (mmHg)						
Systolic	112.4 (0.9)	114.8 (0.7)	0.007	118.1 (1.1)	123.3 (0.9)	< 0.001
Diastolic	69.9 (0.6)	74.7 (0.5)	< 0.001	72.6 (0.9)	81.8 (0.7)	< 0.001
Plasma glucose (mmol/l)						
Fasting	4.1 (0.1)	4.2 (0.1)	0.5	4.1 (0.1)	4.3 (0.1)	0.09
2 hour post load	5.1 (0.1)	4.9 (0.1)	0.2	5.2 (0.1)	5.2 (0.1)	0.8
Insulin (μU/ml)						
Fasting*	2.9 (1.0)	5.4 (1.0)	< 0.001	1.9 (1.0)	4.5 (1.0)	< 0.001
2 hour post load*	10.5 (1.0)	24.2 (1.0)	< 0.001	8.2 (1.0)	21.8 (1.0)	< 0.001
Blood lipids						
Total cholesterol (mmol/l)	2.6 (0.04)	3.6 (0.03)	< 0.001	2.5 (0.1)	3.5 (0.04)	< 0.001
Triglycerides (mmol/l)*	0.48 (1.0)	0.47 (1.0)	0.2	0.48 (1.0)	0.53 (1.0)	0.001
HOMA-IR*	0.09 (1.0)	0.20 (1.0)	< 0.001	0.06 (1.0)	0.10 (1.0)	< 0.001

\* Variables logarithmically transformed before comparisons; non transformed values are expressed for easier interpretation.  
Data are means (standard errors of the mean).

Table 2: Gender specific rural - urban characteristics (unadjusted prevalences and age adjusted odds ratios) for components of the metabolic syndrome:

Components of the metabolic syndrome	WOMEN						MEN					
	Unadjusted prevalences			Age adjusted ORs			Unadjusted prevalences			Age adjusted ORs		
	Rural	Urban	P	Rural	Urban	Rural	Urban	P	Rural	Urban	Rural	Urban
<u>Central obesity or obesity</u>												
WHR (M>0.90/W>0.85)	73.5	13.5	<0.001	1	0.07 (0.05-0.1)*	35.7	13.3	<0.001	1	0.4 (0.3 - 0.7)*		
BMI $\geq$ 30 kg/m <sup>2</sup>	2.4	20.5	<0.001	1	17 (8 - 37)*	1.2	10.9	<0.001	1	21 (6 - 74)*		
WHO definition	74.1	29.6	<0.001	1	0.2 (0.15 - 0.3)*	35.7	20.3	<0.001	1	0.8 (0.5 - 1.1)		
IDF definition	58.6	49.5	0.008	1	1.0 (0.8 - 1.4)	7.5	12.8	0.03	1	3.2 (1.7 - 6.2)†		
NCEP-ATP III Definition	22.2	23.8	0.6	1	1.4 (0.97 - 2.0)	2.4	4.4	0.2	1	4.0 (1.3 - 11)†		
<u>High blood Pressure</u>												
WHO definition	9.4	8.7	0.7	1	2.0 (1.1 - 3.4)†	14.3	21.6	0.02	1	2.9 (1.7 - 4.7)*		
NCEP-ATP III definition	18.3	15.7	0.3	1	1.6 (1.1 - 2.5)†	29.1	35.3	0.09	1	2.0 (1.4 - 3.0)*		
IDF definition	19.0	18.5	0.9	1	2.0 (1.3 - 3.0)*	29.4	37.7	0.03	1	2.3 (1.5 - 3.3)*		
<u>High glucose</u>												
WHO definition	6.4	3.2	0.02	1	0.7 (0.6 - 1.5)	11.5	6.0	0.02	1	0.8 (0.4 - 1.4)		
NCEP-ATP III definition	1.1	1.2	0.9	1	2.1 (0.5 - 8.8)	1.2	1.2	1.0	1	1.6 (0.3 - 8)		
IDF definition	1.3	2.5	0.2	1	2.8 (0.9 - 8.7)	1.6	2.4	0.6	1	2.2 (0.6 - 7.9)		
<u>High lipids</u>												
High triglycerides	0.3	0.4	1.0	1	1.1 (0.1 - 13)	0.4	1.4	0.3	1	3.6 (0.4 - 32)		
High total cholesterol	0.3	2.7	0.005	1	13.0 (2 - 110)†	0.0	2.9	0.006	---	---	---	
$\geq$ 4 <sup>th</sup> quartile HOMA-IR (%)	13.2	37.8	<0.001	1	4.1 (2.8 - 5.9)*	5.8	31.7	<0.001	1	12 (5 - 23)*		
IR and/or IGR (WHO)	19.7	38.9	<0.001	1	2.7 (2.0 - 3.8)*	17.1	33.4	<0.001	1	3.7 (2.7 - 3.8)*		
<u>Prevalence of metabolic syndrome</u>												
WHO definition	1.8	5.9	0.002	1	6.4 (2.5 - 16)†	1.9	7.3	0.001	1	13 (4 - 39)†		
NCEP-ATP III definition	0.0	0.2	1.0	1	---	0.0	0.5	0.5	1	---		
IDF definition	0.3	1.5	0.09	1	11 (1.3-100)‡	0.0	1.2	0.08	1	---		

P for difference: †&lt; 0.05 ‡&lt; 0.01 \*&lt;0.001. IR: insulin resistance; IGR: impaired glucose regulation (impaired glucose tolerance + impaired fasting glycaemia + diabetes)

**Table 3:** Standardised beta coefficients of the relation between each component of the metabolic syndrome and 1) waist circumference 2) HOMA-IR after adjustments, and adjusted R<sup>2</sup> associated with the introduction of these terms in the basic models

Model I†	Waist circumference			HOMA-IR*		
	Standardised β	p	Δ Adjusted R <sup>2‡</sup>	Standardised β	p	Δ Adjusted R <sup>2§</sup>
Systolic blood pressure (mmHg)	0.25	< 0.001	+ 5.7 %	0.10	< 0.001	+ 0.72 %
Diastolic blood pressure (mmHg)	0.26	< 0.001	+ 6.3 %	0.07	0.006	+0.37 %
Fasting plasma glucose (mmol/l)	0.02	0.4	- 0.01 %	0.32	< 0.001	+ 8.00 %
2 hours plasma glucose (mmol/l)	0.045	0.08	+ 0.13 %	0.25	< 0.001	+ 4.78 %
Total cholesterol (mmol/l)	0.12	< 0.001	+ 1.4 %	0.06	0.02	+ 0.23 %
Total triglycerides (mmol/l)*	0.16	< 0.001	+ 4.1 %	0.12	< 0.01	+ 2.04 %
Model II						
Systolic blood pressure (mmHg)	0.18	< 0.001	+ 1.16 %	0.04	0.2	+ 0.05 %
Diastolic blood pressure (mmHg)	0.16	< 0.001	+ 0.94 %	0.001	1.0	- 0.06 %
Fasting plasma glucose (mmol/l)	- 0.008	0.8	- 0.06 %	0.33	< 0.001	+ 8.09 %
2 hours plasma glucose (mmol/l)	- 0.013	0.7	- 0.06 %	0.23	< 0.001	+ 4.30 %
Total cholesterol (mmol/l)	- 0.006	0.87	- 0.05 %	0.013	0.6	- 0.04 %
Total triglycerides (mmol/l)*	0.15	< 0.001	+ 1.78 %	0.09	< 0.001	+ 1.13 %

\* Variables logarithmically transformed before comparisons

Variation of adjusted R<sup>2</sup> when waist circumference (‡) or HOMA-IR (§) is introduced in the model.

† Model adjusted for age, place of residence, and sex

|| Model adjusted for age, place of residence, sex and BMI