



HAL
open science

In vitro glycosidized low-density lipoproteins and low-density lipoproteins isolated from type 2 diabetic patients activate platelets via p38 mitogen-activated protein kinase.

Catherine Calzada, Laurent Coulon, Déborah Halimi, Elodie Le Coquil, Valérie Pruneta-Deloche, Philippe Moulin, Gabriel Ponsin, Evelyne Véricel, Michel Lagarde

► To cite this version:

Catherine Calzada, Laurent Coulon, Déborah Halimi, Elodie Le Coquil, Valérie Pruneta-Deloche, et al.. In vitro glycosidized low-density lipoproteins and low-density lipoproteins isolated from type 2 diabetic patients activate platelets via p38 mitogen-activated protein kinase.. *Journal of Clinical Endocrinology and Metabolism*, 2007, 92 (5), pp.1961-4. <10.1210/jc.2006-2045>. <inserm-00267256>

HAL Id: inserm-00267256

<https://inserm.hal.science/inserm-00267256v1>

Submitted on 26 Mar 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

***In vitro* Glycooxidized Low-Density Lipoproteins and Low-Density Lipoproteins Isolated from Type 2 Diabetic Patients Activate Platelets via p38 MAPK**

Catherine Calzada, Laurent Coulon, Déborah Halimi, Elodie Le Coquil, Valérie Pruneta-Deloché, Philippe Moulin, Gabriel Ponsin, Evelyne Véricel, and Michel Lagarde

INSERM, U870, F-69008 ; INSA-Lyon, RMND, F-69621 ; INRA, U1235, F-69008 ; Univ Lyon 1, F-69003 ; Hospices civils de Lyon, F-69003, Lyon, France

Abbreviated title: Glycooxidized LDL activate platelets

Key terms : Platelets, LDL, lipid peroxidation, type 2 diabetes, oxidative stress

Word count : text 1793, abstract 204, 2 figures, 20 references

Disclosure statement : The authors have nothing to declare.

Corresponding author :

E-mail : Catherine.Calzada@insa-lyon.fr

Dr Catherine Calzada

INSERM UMR 870 /INSA / INRA UMR 1235/ UCBL / HCL / IMBL

Régulations métaboliques, Nutrition et Diabète

Bât. Louis Pasteur, INSA-Lyon

20 av. Albert Einstein

69621 Villeurbanne, France

Tel : (33) 4 72 43 84 79

Fax : (33) 4 72 43 85 24

Structured abstract

Context: Platelet hyperactivation contributes to the increased risk for atherothrombosis in type 2 diabetes and is associated with oxidative stress. Plasma low-density lipoproteins (LDL) are exposed to both hyperglycemia and oxidative stress and their role in platelet activation remains to be ascertained.

Objective: The aim of this study was to investigate the effects of LDL modified by both glycation and oxidation *in vitro* or *in vivo* on platelet arachidonic acid (AA) signaling cascade. The activation of platelet p38 MAPK, the stress kinase responsible for the activation of cytosolic phospholipase A₂, and the concentration of thromboxane B₂, the stable catabolite of the pro-aggregatory AA metabolite thromboxane A₂, were assessed.

Results: Firstly, *in vitro* glycoxidized LDL increased the phosphorylation of platelet p38 MAPK as well as the concentration of thromboxane B₂. Secondly, LDL isolated from plasma of poorly controlled type 2 diabetic patients stimulated both platelet p38 MAPK phosphorylation and thromboxane B₂ production and possessed high levels of malondialdehyde but normal alpha-tocopherol concentrations. By contrast, LDL from sex- and age-matched healthy volunteers had no activating effects on platelets.

Conclusions: Our results indicate that LDL modified by glycooxidation may play an important contributing role in platelet hyperactivation observed in type 2 diabetes via activation of p38 MAPK.

Abbreviations: AA, arachidonic acid; BHT, butylated hydroxytoluene; EDTA, ethylene diamine tetraacetic acid; MDA, malondialdehyde; p38 MAPK, p38 mitogen-activated protein kinase; TBARS, thiobarbituric acid reactive substances; TxA₂, thromboxane A₂; TxB₂, thromboxane B₂.

Introduction

Oxidative stress has been identified as one of the factors closely associated with platelet hyperactivation in type 2 diabetic patients (1) even in the absence of any vascular complications (2). It remains to be determined whether oxidative stress is inherent to platelets and/or is a consequence of circulating factors that could influence platelet function. Among them, low-density lipoproteins (LDL) are submitted to both glycation and oxidation in diabetes (3). In this context, the aims of our study were: 1) to determine the effect of *in vitro* glycooxidized LDL on platelets and compared it with either control, glycated or oxidized LDL. 2) to investigate the effect of LDL isolated from plasma of type 2 diabetic patients compared to healthy volunteers on platelets. Anti-/pro-oxidant status of LDL was assessed and the activation of platelet p38 mitogen-activated protein kinase (p38 MAPK) as well as the formation of thromboxane B₂ (TxB₂) were determined.

Subjects and Methods

In vitro experiments

Low-density lipoprotein isolation from healthy subjects

LDL were isolated from plasma by density gradient ultracentrifugation (density 1.019-1.063 g.ml⁻¹) (4). The concentration of protein was estimated using the Bradford assay (5). LDL were stored at 4°C in the dark under nitrogen and used within 2 days after preparation.

In vitro low-density lipoprotein modification

LDL (3.5 mg protein/ml) in phosphate buffered saline (PBS, pH 7.2) were all incubated at 37°C in the dark, under nitrogen in the presence of sodium azide (1.5 mmol/liter) :

- Control LDL were prepared by incubating LDL with ethylenediamine tetraacetic acid (EDTA) (1 mmol/liter) and butylated hydroxytoluene (BHT) (5 µmol/liter) for 5 days.

- Glycated LDL were LDL incubated with 50 mmol/liter D-glucose for 5 days in the presence of EDTA (1 mmol/liter) and BHT (5 μ mol/liter).

- Oxidized LDL were prepared by incubating LDL for 5 days and then treating them with 1 μ mol/liter CuSO_4 for 1 hour at 37°C.

- Glycoxidized LDL consisted of LDL incubated with 50 mmol/liter D-glucose for 5 days and treating them with 1 μ mol/liter CuSO_4 for 1 hour at 37°C.

LDL preparations were then dialyzed against PBS before their interaction with platelets.

LDL characterization

LDL concentrations of α -tocopherol were determined by reversed-phase HPLC (6). Briefly, LDL samples, containing tocol (6-hydroxy-2-methyl-2-phytylchroman) and δ -tocopherol as internal standards, were extracted with 4 volumes of hexane following the addition of 1 volume of ethanol. Tocopherol isomers were separated onto a Nucleosil C_{18} column, 5 μ m (4 x 150 mm) and detected fluorimetrically (excitation 295nm, emission 340nm).

Overall lipid peroxidation was evaluated by quantitation of malondialdehyde (MDA) by reversed-phase HPLC (7). LDL samples were mixed with thiobarbituric acid (TBA) (10 mmol/liter) and acetic acid in the presence of BHT (5 mmol/liter) and incubated at 95°C for 1 hour. TBA-MDA adduct was extracted with ethyle acetate, separated on a Nucleosil C_{18} column 5 μ m (4.6 x 250 mm) and detected fluorimetrically (excitation 515nm, emission 553nm).

The degree of glycation was determined by the trinitrobenzene sulfonic acid assay (8). LDL glycation was expressed as the percentage of relative reduction of the detected amino groups of lysine of modified LDL compared with control LDL.

Diabetic and control subjects

Ten type 2 diabetic patients (5 men and 5 women, aged 58 ± 2 yr) from the Department of Endocrinology and Metabolic Diseases were matched for sex and age to ten healthy subjects (5 men and 5 women, aged 54 ± 2 yr). Exclusion criteria for diabetic patients were smoking, antioxidant/vitamin supplementation, anti-aggregating drugs, insulin treatment. 6/10 were on metformin or sulfamides, 3/10 on glitazones and 7/10 took lipid-lowering drugs (statins). The patients had poorly controlled diabetes (fasting glycemia : 11.9 ± 2.0 mmol/liter ; glycated hemoglobin HbA_{1C} : $8.8 \pm 0.6\%$). They had mild dyslipidemia with mild hypertriglyceridemia (triglycerides : 2.0 ± 0.2 mmol/liter), normal LDL-cholesterol (2.7 ± 0.3 mmol/liter) and normal HDL-cholesterol (1.3 ± 0.1 mmol/liter). Control subjects were in good health as assessed by medical history and exclusion criteria were any pathology including diabetes and anti-aggregatory drugs. Written informed consent was obtained from all participants.

Interaction between platelets and LDL

Platelet isolation and incubation with LDL

Blood was collected at the regional blood center from healthy volunteers who had not ingested any aspirin or other non-steroidal anti-inflammatory drugs in the previous 10 days. Platelets were prepared (9) and incubated in the presence or absence of LDL (500 μ g/ml) for 2 hours at 37°C.

Platelet p38 MAPK activation

Platelets were lysed (10), proteins (25 μ g) were denatured for 10 min at 100°C, electrophoresed in 12% Tris-HCl polyacrylamide gels at 25 mA for 135 min and transferred to nitrocellulose membranes (100V, 30 min). The membranes were incubated with either 1/1000 anti-p38 MAPK or anti-phospho-p38 MAPK polyclonal antibodies (Cell Signaling Technologies, Beverly, MA, USA), washed and incubated with 1/2000 goat anti-rabbit horseradish peroxidase conjugate for 1 h. p38 MAPK and phospho-p38 MAPK were

visualized by enhanced chemiluminescence and bands were quantified by densitometry with an ImageMaster VDS-CL camera (Amersham Biosciences, Buckinghamshire, UK).

Platelet thromboxane B₂ measurement

Platelet TxB₂ was quantified by enzyme immunoassay (Amersham Biosciences, Buckinghamshire, UK). Coefficient of variation was lower than 10%.

Statistical analysis

Results are expressed as the mean \pm SD. Comparisons between groups were performed using a Wilcoxon test. Statistical significance was established at $p < 0.05$.

Results

Effects of in vitro modified LDL isolated from healthy subjects on platelets

Glycoxidized LDL were compared with LDL incubated solely with glucose or CuSO₄ to differentiate the specific effects of glycation or oxidation from combined effects. The incubation of LDL with 50 mmol/liter glucose had no significant effects on LDL α -tocopherol (7.70 ± 1.84 nmol/mg protein in glycated LDL vs 8.53 ± 2.27 nmol/mg protein in control LDL, n=8) and MDA level compared to control LDL (0.19 ± 0.16 nmol/mg protein vs 0.16 ± 0.17 nmol/mg protein, respectively, n=5). As expected, the addition of CuSO₄ to LDL resulted in a decreased α -tocopherol (6.73 ± 3.27 nmol/mg protein) and an increased MDA levels (0.27 ± 0.26 nmol/mg protein) compared to control LDL. Glycooxidation of LDL neither led to significant additional decreases of α -tocopherol (7.03 ± 2.34 nmol/mg protein) nor to further increases of peroxide levels (0.27 ± 0.33 nmol/mg protein) compared to oxidized LDL. Their percentage of glycation was $34 \pm 6\%$, similar to that measured in glycated LDL ($35 \pm 9\%$).

The incubation of platelets with control LDL or glycated LDL had no significant effect on the activation of p38 MAPK (Fig. 1A) and TxB₂ concentration (Fig. 1B) as compared to platelets alone. The addition of oxidized LDL to platelets increased p38 MAPK phosphorylation by 2-fold and TxB₂ concentration LDL by 69%. Finally, glycoxidized LDL had the most pronounced effect on platelet p38 MAPK phosphorylation (3-fold increase) and TxB₂ level (2.4-fold increase) compared to platelets alone.

Effect of LDL isolated from type 2 diabetic patients on platelets

MDA levels increased by 6-fold in LDL from type 2 diabetic patients compared to LDL from control subjects (1.74 ± 2.58 nmol/mg protein vs 0.29 ± 0.29 nmol/mg protein, respectively, n=10) whereas LDL α -tocopherol concentrations did not differ between patients and control subjects (10.25 ± 1.94 nmol/mg protein vs 10.26 ± 3.25 nmol/mg protein in LDL, respectively, n=10). The incubation of platelets with LDL from diabetic patients resulted in a 2.2-fold increase of phosphorylated p38 MAPK amount (Fig. 2A) and a 2-fold increased basal concentration of TxB₂ as compared to platelets (578 ± 164 pmol/10⁹ platelets vs 267 ± 22 pmol/10⁹ platelets) (Fig. 2B). It is worth noting that neither significant phosphorylation of p38 MAPK nor increase of TxB₂ concentration were observed in platelets incubated with LDL from healthy volunteers.

Discussion

Our results show that LDL modified by glycooxidation *in vitro* or *in vivo* activate platelets whereas control or glycated LDL have no effect. Previous studies on the interaction between platelets and LDL have shown that LDL do not induce platelet activation *per se* (11) but may increase the sensitivity of platelets to different agonists (12) *via* an increased phosphorylation of p38 MAPK (13). Such a discrepancy between results could be ascribed to differences in the

degree of LDL oxidation. We show that the incubation of LDL with glucose did not change their anti-/pro-oxidant status corroborating that glycation alone is insufficient to promote LDL oxidation (14). Moreover, glycated LDL had no stimulating effects on platelets. Although one study reported that glycated LDL caused an increased platelet aggregating response to ADP, the presence of high levels of TBARS in those LDL suggests that the effects were likely due to the combination of LDL glycation and oxidation (15). We also show that copper-oxidized LDL increased p38 MAPK phosphorylation by 2-fold and TxB₂ concentration by 69% confirming that LDL defined as minimally or mildly oxidized may activate platelets (12,16). Finally, glycoxidized LDL, reflecting at best the state that may occur in LDL from diabetic patients, were the most effective triggers of platelet AA signaling cascade. Although the presence of glucose has been shown to accelerate copper-induced LDL oxidation (17), we did not observe any significant differences between oxidized LDL and glycoxidized LDL in terms of vitamin E and MDA levels. This could be related to the copper-LDL ratios used which are much lower in our study than in the above published studies, as suggested by Knott *et al.* (14). Supporting our *in vitro* results, we present new data indicating that LDL isolated from poorly controlled type 2 diabetic patients increase platelet p38 MAPK phosphorylation and TxB₂ formation whereas those isolated from healthy control subjects do not. It has been shown that LDL isolated from type 1 diabetic patients increased platelet aggregation and TxB₂ release (18). Concerning the agents in LDL from diabetic patients responsible for platelet activation in our experiments, their identification seem rather difficult since LDL represent a heterogenous mixture of particles modified to different degrees of glycation and oxidation. However, an important feature is that LDL from selected patients possessed high concentrations of MDA which supports data reporting increased levels of plasma lipid peroxides in type 2 diabetic patients (19). No modification of vitamin E was observed ruling out an involvement of this antioxidant. Moreover, LDL isolated from selected diabetic

patients can be defined as glycoxidized LDL since the percent of HbA_{1C} (8.8%) is known to be correlated with the percent of glycated LDL (20). In conclusion, LDL from type 2 diabetic patients – as well as glycoxidized LDL *in vitro* – activate platelet AA signaling cascade. The effect of LDL appears to be related to the combination of hyperglycemia and lipid peroxidation independently of vitamin E status. Thus, it suggests that glycoxidized LDL may act as one of the triggers of platelet activation occurring in type 2 diabetes.

Acknowledgements

This work was supported by INSERM and Agence Nationale de la Recherche ANR 2005 "Cardiovasculaire, Obésité et Diabète". We gratefully thank nurses for expert blood drawing assistance. Catherine Calzada is supported by CNRS.

References

1. **De Cristofaro R, Rocca B, Vitacolonna E, Falco A, Marchesani P, Ciabattoni G, Landolfi R, Patrono C, Davi G** 2003 Lipid and protein oxidation contribute to a prothrombotic state in patients with type 2 diabetes mellitus. *J Thromb Haemost* 1:250-256
2. **Véricel E, Januel C, Carreras M, Moulin P, Lagarde M** 2004 Diabetic patients without vascular complications display enhanced basal platelet activation and decreased antioxidant status. *Diabetes* 53:1046-1051
3. **Hunt JV, Smith CC, Wolff SP** 1990 Autooxidative glycosylation and possible involvement of peroxides and free radicals in LDL modification by glucose. *Diabetes* 39:1420-1424
4. **Havel RJ, Eder HA, Bragdon JH** 1955 The distribution and chemical composition of ultracentrifugally separated lipoproteins in human serum. *J Clin Invest* 34:1345-1353
5. **Bradford MM** 1976 A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248-254
6. **Véricel E, Croset M, Sedivy P, Courpron Ph, Dechavanne M, Lagarde M** 1988 Platelets and aging. I- Aggregation, arachidonate metabolism and antioxidant status. *Thromb Res* 49:331-342
7. **Therasse J, Lemonnier F** 1987 Determination of plasma lipoperoxides by high-performance liquid chromatography. *J Chromatogr* 413:237-241
8. **Duell PB, Oram JF, Bierman EL** 1990 Nonenzymatic glycosylation of HDL resulting in inhibition of high-affinity binding to cultured human fibroblasts. *Diabetes* 39:1257-1263
9. **Lagarde M, Bryon PA, Guichardant M, Dechavanne M** 1980 A simple and efficient method for platelet isolation from their plasma. *Thromb Res* 17:581-588

10. **Coulon L, Calzada C, Moulin P, Véricel E, Lagarde M** 2003 Activation of p38 mitogen-activated protein kinase/cytosolic phospholipase A2 cascade in hydroperoxide-stressed platelets. *Free Radic Biol Med* 35:616-625
11. **Meraji S, Moore CE, Skinner O, Bruckdorfer KR** 1992 The importance of oxidation or glycosylation of low-density lipoproteins in relation to platelet activation. *Platelets* 3:155-162
12. **Weidtmann A, Scheithe R, Hrboticky N, Pietsch A, Lorenz R, Siess W** 1995 Mildly oxidized LDL induces platelet aggregation through activation of phospholipase A2. *Arterioscler Thromb Vasc Biol* 15:1131-1138
13. **Hackeng CM, Relou IA, Pladet MW, Gorter G, van Rijn HJ, Akkerman JW** 1999 Early platelet activation by low density lipoprotein via p38MAP kinase. *Thromb Haemost* 82:1749-1756
14. **Knott HM, Brown BE, Davies MJ, Dean RT** 2003 Glycation and glycooxidation of low-density lipoproteins by glucose and low-molecular mass aldehydes. Formation of modified and oxidized particles. *Eur J Biochem* 270:3572-3582
15. **Ferretti G, Rabini RA, Bacchetti T, Vignini A, Salvolini E, Ravaglia F, Curatola G, Mazzanti L** 2002 Glycated low density lipoproteins modify platelet properties: a compositional and functional study. *J Clin Endocrinol Metab* 87:2180-2184
16. **Relou IA, Hackeng CM, Akkerman JW, Malle E** 2003 Low-density lipoprotein and its effect on human blood platelets. *Cell Mol Life Sci* 60:961-971
17. **Mowri H, Frei B, Keaney JF** 2000 Glucose enhancement of LDL oxidation is strictly metal ion dependent. *Free Radic Biol Med* 29:814-824
18. **Watanabe J, Wohltmann HJ, Klein RL, Colwell JA, Lopes-Virella MF** 1988 Enhancement of platelet aggregation by low-density lipoproteins from IDDM patients. *Diabetes* 37:1652-1657

19. **Griesmacher A, Kindhauser M, Andert SE, Schreiner W, Toma C, Knoebl P, Pietschmann P, Prager R, Schnack C, Schernthaner G, Mueller MM** 1995 Enhanced serum levels of thiobarbituric-acid-reactive substances in diabetes mellitus. *Am J Med* 98:469-475
20. **Jenkins AJ, Thorpe SR, Alderson NL, Hermayer KL, Lyons TJ, King LP, Chassereau CN, Klein RL** 2004 In vivo glycated low-density lipoprotein is not more susceptible to oxidation than nonglycated low-density lipoprotein in type 1 diabetes. *Metabolism* 53:969-976

FIG. 1 Effect of *in vitro* modified LDL on platelet p38 MAPK phosphorylation (A) and TxB₂ concentration (B).

Platelets were incubated for 2 h at 37°C in the absence (0) or presence of 500µg/ml control LDL, glycated LDL (LDL + 50 mmol/liter glucose), oxidized LDL (LDL + 1µmol/liter CuSO₄) or glycoxidized LDL (LDL + 50 mmol/liter glucose + 1µmol/liter CuSO₄). Data are means ± SD of 5 to 8 experiments. ^a*P* < 0.05 vs platelets.

FIG. 2 Effect of LDL from control subjects and type 2 diabetic patients on platelet p38 MAPK phosphorylation (A) and TxB₂ concentration (B).

Platelets were incubated for 2 h at 37°C in the absence (0) or presence of LDL (500µg/ml). Results are means ± SD (n=10). ^a*P* < 0.05 vs platelets, ^b*P* < 0.05 vs platelets + LDL from control subjects.



A. p38 MAPK phosphorylation**B. TxB₂ concentration**

FIG.1



A. p38 MAPK phosphorylation**B. TxB₂ concentration**

FIG. 2