



## **Birth characteristics and childhood malignant central nervous system tumors: the ESCALE study (French Society for Childhood Cancer).**

Nathalie Mallol-Mesnard, Florence Menegaux, Brigitte Lacour, Olivier Hartmann, Didier Frappaz, François Doz, Anne-Isabelle Bertozzi, Pascal Chastagner, Denis Hémon, Jacqueline Clavel

### **► To cite this version:**

Nathalie Mallol-Mesnard, Florence Menegaux, Brigitte Lacour, Olivier Hartmann, Didier Frappaz, et al.. Birth characteristics and childhood malignant central nervous system tumors: the ESCALE study (French Society for Childhood Cancer).. Cancer Detection and Prevention, 2008, 32 (1), pp.79-86. 10.1016/j.cdp.2008.02.003 . inserm-00273538

**HAL Id: inserm-00273538**

**<https://inserm.hal.science/inserm-00273538>**

Submitted on 15 Apr 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Birth characteristics and childhood malignant central nervous system tumours: the ESCALE study (SFCE\*)**

**Short title:** Childhood CNS tumours and birth characteristics.

**Category:** original article

**Condensed abstract:** Pre- and perinatal characteristics (birth weight, gestational age, birth order, breastfeeding, congenital anomalies, use of assisted reproductive technology, and foetal losses) were obtained for 209 cases of childhood malignant central nervous system tumour and 1681 frequency matched controls. The results suggest an association with a maternal history of miscarriages.

## **Authors:**

Nathalie Mallol-Mesnard (MD)<sup>1,2</sup>, Florence Menegaux (MD, PhD)<sup>1,2</sup>, Brigitte Lacour (MD)<sup>1,2,3</sup>, Olivier Hartmann (MD)<sup>4</sup>, Didier Frappaz (MD)<sup>5</sup>, François Doz (MD)<sup>6</sup>, Anne-Isabelle Bertozzi (MD)<sup>7</sup>, Pascal Chastagner (MD)<sup>8</sup>, Denis Hémon (PhD)<sup>1,2</sup>, Jacqueline Clavel (MD, PhD)<sup>1,2,9</sup>

## **Affiliations:**

<sup>1</sup>INSERM, U754, Villejuif, France. <sup>2</sup>Univ Paris-Sud, UMR-S754, IFR69, Villejuif, France. <sup>3</sup>Registre National des Tumeurs solides de l'Enfant, Nancy, France. <sup>4</sup>Institut Gustave Roussy, Villejuif, France. <sup>5</sup>Centre Léon Bérard, Lyon, France. <sup>6</sup>Institut Curie, Paris, France. <sup>7</sup>Hôpital des Enfants, Toulouse, France. <sup>8</sup>Hôpital d'Enfants, Vandoeuvre les Nancy, France. <sup>9</sup>Registre National des Hémopathies malignes de l'Enfant, Villejuif, France.

\*SFCE: Société Française de lutte contre les Cancers de l'Enfant et de l'Adolescent

## **Corresponding author:**

Jacqueline Clavel

Inserm U754

16, av. Paul Vaillant-Couturier

F-94807 VILLEJUIF Cedex

Tel: 33 1 45 59 50 38; Fax: 33 1 45 59 60 20; E-mail: [clavel@vjf.inserm.fr](mailto:clavel@vjf.inserm.fr)

**Sources of support:** This work was supported by grants from INSERM, the Fondation de France, the Association pour la Recherche contre le Cancer (ARC), the Agence Française de Sécurité Sanitaire des Produits de Santé (AFSSAPS), the Agence Française de Sécurité Sanitaire de l'Environnement et du Travail (AFSSET) and the association, Cent pour Sang la Vie.

**Liste of abbreviations**

ESCALE: Etude Sur les Cancers et les Leucémies de l'Enfant (Study on childhood cancers and leukemia)

EUROCAT: European Surveillance of Congenital Anomalies

ICCC: International Classification on Childhhod Cancer

ICD-I-3: International Classification of Diseases for Oncolgy, Third Edition

RNHE: Registre National des Hémopathies malignes de l'Enfant (National Registry of Childhood Haematological Cancers)

RNTSE: Registre National des Tumeurs Solides de l'Enfant (National Registry of Childhood Solid Tumours)

SFCE: Société Française de lutte contre les Cancers de l'Enfant et de l'adolescent (French society for childhood cancer)

## Abstract

**Background:** Determining the role of pre- and perinatal factors in the aetiology of childhood malignant central nervous (CNS) tumours, using data from the French national case-control study, ESCALE. **Methods:** ESCALE included all children in France less than 15 years old with a diagnosis of acute leukaemia, lymphoma, malignant CNS tumour, or neuroblastoma (2003-2004). In all, 209 malignant CNS tumour cases (80% of the eligible cases) and 1681 population-based controls (71%) were included using quotas ensuring frequency matching with the cases by age and gender. Case and control mothers were interviewed using a standardised telephone interview, which elicited birth characteristics, congenital malformation, maternal reproductive history, and use of assisted reproductive technologies for the index child. **Results:** The cases and controls did not differ in terms of gestational age at birth, birth weight, birth order, breastfeeding, or parental age at birth. There was no association between assisted reproduction for the index child and malignant CNS tumour (OR = 1.1 [0.6-2.2]). A positive association between a maternal history of one miscarriage and malignant CNS tumour was observed (OR = 1.4 [1.0-2.0],  $p < 0.05$ ), especially for glial cell tumours (other glioma: OR = 2.0 [1.1-3.6]). **Conclusion:** The results suggest a possible association between a maternal history of one miscarriage and the risk of malignant CNS tumour.

**Keywords:** Epidemiology ; risk factors ; childhood CNS tumour ; birth characteristics

## Introduction

Central Nervous System [CNS] tumours are the most common childhood cancer, after leukaemia, in developed countries [1], accounting for 20 to 25% of all childhood neoplasms. In France, about 200 new cases of malignant CNS tumours are diagnosed each year [2]. Most of the cases emerge before the age of 5 years, suggesting that antenatal, perinatal, or early postnatal exposures may be considered potential risk factors [3]. Most of the central nervous system's development occurs during foetal life and over the first 2 or 3 years of extrauterine life, with a trend toward heterogeneity of the various types of central nervous system cells. It is therefore important to consider the period of exposure as well as the type of exposure suspected. Apart from ionizing radiation and a few rare genetic syndromes, the aetiology of childhood CNS tumours is largely unknown. Maternal reproductive history, such as foetal losses and assisted reproduction technologies for the index child, has often been studied, but associations with CNS tumors have rarely been observed. Birth characteristics, such as gestational age, birth weight, parent's age at birth, birth order and congenital malformations, have also been studied. Only congenital malformations seem to be frequently associated with CNS tumors [4, 5]. This paper reports the results for maternal reproductive history, birth characteristics, and the risk of childhood malignant CNS tumors, based on data from the ESCALE study.

## Materials and methods

ESCALE is a French national population-based case-control study conducted in 2003 and 2004, which addressed 4 childhood neoplastic diseases (leukaemia, lymphoma, neuroblastoma, and malignant central nervous system tumour) and infectious, environmental, and genetic risk factors. This paper concerns the cases of malignant central nervous system tumours.

### *Study population and data collection*

The design of the study has already been published elsewhere [6]. The cases were all children (0-14 years) whose malignant central nervous system tumour was diagnosed between 2003 and 2004. The cases were living in France at the time of diagnosis. All the cases were recruited directly by investigators assigned to each French paediatric oncology hospital

department with the support of the two National Registries of childhood cancer: the National Registry of Childhood Haematological Cancers (RNHE) [7] and the National Registry of Childhood Solid Tumours (RNTSE) [8]. The diagnoses were confirmed by crosschecking with the RNTSE, which was also able to give a histological confirmation for 81% of the cases and a clinical/radiological confirmation for 16%. All the cases have been classified into 4 groups using the third version of the International Classification of Childhood Cancer (ICCC) based on the ICD-O-3 [9]: embryonal tumours (ICD-O-3 codes 9470/3, 9471/3, 9473/3, 9474/3, 9480/3, 9508/3), astrocytomas (codes 9380/3, 9400/3, 9401/3, 941/3, 9420/3, 9424/3, 9440/3), other gliomas (codes 9380/3, 9381/3, 9382/3, 9430/3, 9450/3, 9451/3), and ependymomas (codes 9390/3-9393/3).

For ethical reasons, children who had died or who were receiving palliative care before the inclusion date were not eligible. The cases that were adopted, whose mothers were not French-speaking or who had serious psychiatric disorder were not eligible. Out of the 343 cases of malignant central nervous system tumours (with a /3 behaviour ICD-O-3 codes) diagnosed in France from January 1, 2003, to December 31, 2004, 82 were therefore not eligible (57 early deaths, 1 child receiving palliative care, 14 non-French speaking mothers, 5 non-biological mothers, and 5 psychiatric disorders). Out of the 261 eligible cases, 19 refused to participate and 33 could not be contacted by interviewers despite repeated attempts. Finally, 209 (80%) of the eligible cases of malignant CNS tumour were included in the study.

The control group was selected from the French population by sampling from 60,000 representative addresses taken from the French national telephone directory plus unlisted phone numbers generated randomly. Age and gender quotas were applied to ensure the same age and sex distribution as the group of cases. Additional quotas for the number of children living in the household, based on the last population census, were used to improve the comparability of the controls with the French population with regard to birth order. The latter is related to a number of variables of interest, such as early day-care or breast-feeding, and might otherwise have been biased by sampling. Out of the 50,217 phone numbers contacted, 46,994 were ineligible numbers (22,584 business or inactive numbers, 18,456 households without children, 5,277 outside of the quotas, and 677 with no living, French-speaking biological mother). The eligibility of 862 phone numbers whose owners hung up too early could not be determined. Out of the 2361 remaining

eligible controls, 679 parents refused and 1 child had a prior history of neuroblastoma. Finally, 1681 controls (71%) were included in the study.

The case and control mothers' interviews were conducted by telephone by trained interviewers using the same standardised questionnaire. The interview elicited data on demographic and socio-economic characteristics, parental occupational history, childhood environment, familial and personal medical history, and history of pregnancy. The mother's obstetric history was specifically described: foetal losses (spontaneous, voluntary, or therapeutic) and assisted reproduction for the index child. The mothers were also asked to describe the characteristics of the index child at birth: gestational age, birth weight, parents' age at birth, birth order, breastfeeding, and presence and site of congenital malformations. Malformations were classified as major or minor malformations using the European Surveillance of Congenital Anomalies (EUROCAT) classification [10].

### ***Statistical analysis***

The SAS<sup>®</sup> software package (version 9, Cary, North Carolina) was used for all the analyses. Analyses were carried out for all central nervous system tumours considered malignant, in combination and separately for each large group of malignant CNS tumours: ependymomas, embryonal tumours, astrocytomas, and other gliomas. All the variables were treated as categorical variables. Odds ratios (OR) and their 95% confidence intervals (95% CI) were estimated using unconditional logistic regression, closely adjusted for age and gender. Polytomous logistic regression adjusted for age and gender was also used to study the large groups of CNS tumours simultaneously. Additional adjustments for potential confounding factors and mutual adjustments were also made.

## **Results**

For the 209 cases, the diagnoses were as follows: ependymoma in 33 cases; embryonal tumours in 100 cases; high grade astrocytoma in 26 cases; other glioma in 45 cases; and other types of CNS tumour in 5 cases. In 7 cases, type 1 neurofibromatosis (NF1) had been diagnosed.

The cases did not differ from the controls with respect to gender (table I). Since age quotas for selection of the controls had been defined to enable comparison with the whole case

group, there was a small age discrepancy between the CNS cases and controls, particularly for very young children. There were, however, at least 5 controls for each case in each age group, and the mean ages of the cases and controls were very similar (6.3 vs. 6.0,  $p = 0.4$ ). The cases and controls were also similar with respect to the parents' marital status, educational level and socioeconomic category (table I). The number of children living in the household, made comparable to that in the French population by quotas, differed significantly between cases and controls: single children were more frequent for the controls than for the cases.

The cases' mothers reported slightly more frequent history of one miscarriage (OR = 1.4 [1.0-2.0],  $p < 0.05$ ), but this association was restricted to some glial cell tumours (other gliomas: OR = 2.0 [1.1-3.6]) (table II). When mothers who had also had voluntary or therapeutic abortions were excluded, the results were unchanged. Childhood CNS tumour was not related to voluntary or therapeutic abortion, except for the ependymomas, which were associated with voluntary abortion. There was no association between childhood CNS tumour and assisted reproduction, irrespective of the type of tumour.

Overall, gestational age was not associated with CNS tumour, except for embryonal tumours, which were significantly associated with a gestational age at birth of less than 37 weeks (table III). No association between malignant CNS tumours and birth weight or parental age at birth or breast-feeding was observed, irrespective of the type of tumour. An isolated association between birth order and embryonal tumours was observed, with a higher OR for second born children but not for children born subsequently.

A total of 8 cases presented with congenital malformations, 6 of which were major malformations (1 cleft lip, 1 cleft palate, 1 congenital malformation of the circulatory system, 1 congenital dislocation of the hip, 1 child with webbed toes, and 1 child with spade-like hands) and 2 of which were minor (1 non-neoplastic giant congenital nevus and 1 case of depressions in the skull). No association between major or minor malformations and CNS tumour was evidenced (table III). However, malformations, and especially major malformations, were more often positively, but non-significantly, associated with glial cell tumours (astrocytomas: OR = 1.9 [0.2-14], other gliomas: OR = 2.9 [0.9-10]) than with the other types of CNS tumour.

The results were unchanged after adjusting for the parents' educational level and socioeconomic category. The results were also unchanged when maternal reproductive history



and birth characteristics were included in the same model. Lastly, the exclusion of cases with type 1 neurofibromatosis did not modify the results.

## Discussion

In short, the present study did not provide evidence of any association between malignant CNS tumours and gestational age at birth, birth weight, birth order, breast-feeding, parental age at birth, or medically assisted reproduction. Glial cell tumours (astrocytoma or other glioma) were significantly associated with a maternal history of miscarriages, and, to a lesser extent and not significantly, with the existence of major congenital malformations.

With a statistical power of 80% and an  $\alpha$  error of 5%, the study was able to detect odds ratios from 1.5 (30% exposed controls) to 2.2 (5% exposed controls). The statistical power, however, was lower for subtype differences, particularly for uncommon exposures such as malformations.

All cases were identified directly by the network of investigators working with the 2 French National Registries of Childhood Cancer (RNHE and RNTSE), making selection bias due to the case identification process unlikely. The ineligibility of cases who had died or were under palliative care may have induced a survival bias. To the authors' knowledge, however, none of the variables of interest are known to be survival factors, after accounting for age and tumour type [11]. The study included 19 children who died early, after the interview. They were younger at diagnosis than the children who survived (5.4 vs. 6.1 years old), but comparable, although the numbers were small, with regard to the other characteristics. Therefore, the exclusion of early deaths is unlikely to have induced a major selection bias. The 52 cases (20%) who did not answer the questionnaire had similar age and gender distributions to those of the respondent cases.

All the diagnoses were ascertained by the National Childhood Registry of Solid Tumours (RNTSE) and confirmed by histology (81%) and radiology (16%). The results did not change after exclusion of cases whose tumour had not been histologically confirmed.

The controls were selected from the general population using the national telephone directory as a random basis together with randomly generated unlisted numbers. The sampling process also made the controls similar to the French population in terms of number of children

living in the household and, subsequently, in terms of birth order, when compared with the national perinatal surveys [12, 13]. Comparison with those surveys also showed that the controls were very similar to the French population in terms of paternal age at birth, mother's educational level, voluntary abortion, assisted reproduction, gestational age at birth and birth weight, and maternal reproductive history. The controls' mothers were slightly older than in the overall population. However, no association between the mother's age at birth and childhood CNS tumour was observed and adjusting for that variable did not modify the results. The proportion of congenital malformations in the controls (3.4%) was very close to what was expected on the basis of the Paris Registry of Congenital Malformations (2.2% to 3.2% for children born between 1988 and 2000) [14].

In order to avoid misclassification with respect to gestational age, birth weight, abortions, and assisted reproduction, only biological mothers were interviewed and identical questionnaires were used for the cases and controls. The frequent examinations and computed tomography and ultrasound scans of children with brain tumours are liable to promote malformation detection and reporting, and may have led to differential misclassification. However, this probably mainly concerns minor, rather than major, malformations and very young children. Major malformations are unlikely to remain undetected after the age of one year. In this study, all the cases with malformations were at least one year old. Because the information on malformations was only given by the mothers, ascertainment may have been incomplete, but similarly, for both the cases and controls.

As was the case in this study, the previous studies did not evidence any association between birth order and childhood CNS tumour [15-25]. The association between embryonal tumours and second born children observed in the present study was isolated, with no linear trend with increasing birth order, and would appear to have no particular biological plausibility.

A potential link between birth weight and childhood leukemia has been suggested in a meta-analysis of 18 studies [26] and in several studies which were not included in the meta-analysis [27-31]. Insulin-like growth factor has been proposed as a possible explanation [32]. For CNS tumours, no relationship with high birth weight was reported in most of the previous studies [17, 18, 21, 22, 33-37]. The only relationship with high birth weight was reported for astrocytomas [15, 19, 25]. When the results were reported separately for astrocytomas, they were discordant [15, 25].

257           Only 2 of the 10 previous studies [19, 20, 23, 25, 38-42] showed a positive relationship  
258 between CNS tumours and premature birth [40, 42]. As in the present study, one of those studies  
259 reported a relationship with medulloblastoma [42].

260           Several studies have suggested that breastfeeding may have a protective affect in  
261 childhood cancer [43], especially in childhood leukaemia. Nevertheless, none of the studies  
262 investigating the role of breastfeeding in childhood CNS tumours, including the present study,  
263 observed such inverse association [21, 23, 37, 40].

264           Most studies considering maternal history of foetal losses have not shown any association  
265 with CNS tumours [15, 16, 19-21, 23-25]. However, one study investigating malignant CNS  
266 tumours also reported a positive association with malignant astrocytoma [15]. A variety of terms  
267 were used in the previous studies and it is not always possible to distinguish between  
268 spontaneous, voluntary and therapeutic abortions. The terms used by the mothers themselves are  
269 probably not always reliable. In the present study, no association was found when all the types of  
270 foetal losses (spontaneous, therapeutic, and voluntary abortions) were considered together. The  
271 observed association between voluntary abortion and ependymoma remained stable after  
272 adjusting for well known confounding factors such as socioeconomic characteristics or smoking  
273 during pregnancy. It is therefore difficult to explain except by misclassification of the type of  
274 abortion.

275           Many studies have evidenced an association between congenital malformations and CNS  
276 tumours [4, 5, 44-46], sometimes with a stronger association when the malformation had a CNS  
277 site [44, 46]. In the present study, none of the controls and only one case presented with a brain  
278 malformation, and that malformation was minor. Congenital malformations have decreased in  
279 France since 1995, thanks to active prevention (folate supplementation) and early screening for  
280 major CNS malformations such as spina bifida, enabling therapeutic abortion [14].

281           To the authors' knowledge, only 3 studies on CNS tumour and medically assisted  
282 reproduction have been published. None of them reported any association [23, 47, 48]. More  
283 generally, no association between childhood malignancies and medically-assisted reproduction  
284 has ever been reported [49-52]. The present results are therefore consistent with those of the  
285 published studies.

## Conclusion

The present study does not support a role of prematurity, birth weight, breastfeeding or parental age in the occurrence of malignant CNS tumours. It suggests that a maternal history of miscarriages may be associated with the risk of malignant CNS tumours.

## Acknowledgements

We are grateful to: the coordinators of the recruitment of the cases; the staff of the French National Registry of Childhood Solid Tumours, who contributed to case detection and verification; the CSA Institute, who coordinated the selection of the controls and the interviews; the team of interviewers, who interviewed the cases and the controls; and our translator for his skilful revision of the manuscript.

## References

1. Baldwin, RT, and Preston-Martin S. Epidemiology of brain tumours in childhood - a review. *Toxicol Appl Pharmacol.* 2004; 199(2): 118-31.
2. Desandes E, Clavel J, Berger C, Bernard JL, Blouin P, de Lumley L, et al. Cancer incidence among children in France, 1990-1999. *Pediatr Blood Cancer.* 2004; 43(7): 749-57.
3. Autrup H. Transplacental transfer of genotoxins and transplacental carcinogenesis. *Environ Health Perspect.* 1993; 101 Suppl 2: 33-8.
4. Altmann AE, Halliday JL and Giles GG. Associations between congenital malformations and childhood cancer. A register-based case-control study. *Br J Cancer.* 1998; 78(9): 1244-9.
5. Agha MM, Williams JI, Marrett L, To T, Zipursky A and Dodds L. Congenital abnormalities and childhood cancer. *Cancer.* 2005; 103(9): 1939-48.
6. Rudant J, Menegaux F, Leverger G, Baruchel A, Nelken B, Bertrand Y, et al. Family history of cancer in children with acute leukemia, Hodgkin's lymphoma or non-Hodgkin's lymphoma: the ESCALE study (SFCE\*). *Int J Cancer.* 2007; 121(1): 2873-80.

7. Clavel J, Goubin A, Auclerc MF, Auvrignon A, Waterkeyn C, Patte C, et al. Incidence of childhood leukaemia and non-Hodgkin's lymphoma in France: National Registry of Childhood Leukaemia and Lymphoma, 1990-1999. *Eur J Cancer Prev.* 2004; 13(2): 97-103.
8. <http://www.chu-nancy.fr/rntse/>.
9. Steliarova-Foucher E, Stiller C, Lacour B and Kaatsch P. International Classification of Childhood Cancer, third edition. *Cancer.* 2005; 103(7): 1457-67.
10. De Wals P, Dolk H, Bertrand F, Gillerot Y, Weatherall JA and Lechat MF. [Epidemiologic surveillance of congenital abnormalities using the EUROCAT Register]. *Rev Epidemiol Sante Publique.* 1988; 36(4-5): 273-82.
11. Gurney JG and Kadan-Lottick N. Brain and other central nervous system tumours: rates, trends, and epidemiology. *Curr Opin Oncol.* 2001; 13(3): 160-6.
12. Blondel B, Breart G, du Mazaubrun C, Badeyan G, Wcislo M, Lordier A, et al. [The perinatal situation in France. Trends between 1981 and 1995]. *J Gynecol Obstet Biol Reprod (Paris).* 1997; 26(8): 770-80.
13. Blondel B, Supernant K, Du Mazaubrun C and Breart G. [Trends in perinatal health in metropolitan France between 1995 and 2003: results from the National Perinatal Surveys]. *J Gynecol Obstet Biol Reprod (Paris).* 2006; 35(4): 373-87.
14. De Vigan C, Khoshnood B, Lhomme A, Vodovar V, Goujard J and Goffinet F. [Prevalence and prenatal diagnosis of congenital malformations in the Parisian population: twenty years of surveillance by the Paris Registry of congenital malformations]. *J Gynecol Obstet Biol Reprod (Paris).* 2005; 34(1 Pt 1): 8-16.
15. Emerson JC, Malone KE, Daling JR and Starzyk P. Childhood brain tumour risk in relation to birth characteristics. *J Clin Epidemiol.* 1991; 44(11): 1159-66.
16. Cordier S, Iglesias MJ, Le Goaster C, Guyot MM, Mandereau L and Hemon D. Incidence and risk factors for childhood brain tumours in the Ile de France. *Int J Cancer.* 1994; 59(6): 776-82.
17. McCredie M, Maisonneuve P and Boyle P. Perinatal and early postnatal risk factors for malignant brain tumours in New South Wales children. *Int J Cancer.* 1994; 56(1): 11-5.
18. Savitz DA and Ananth CV. Birth characteristics of childhood cancer cases, controls, and their siblings. *Pediatr Hematol Oncol.* 1994; 11(6): 587-99.

19. Linet MS, Gridley G, Cnattingius S, Nicholson HS, Martinsson U, Glimelius B, et al. Maternal and perinatal risk factors for childhood brain tumours (Sweden). *Cancer Causes Control*. 1996; 7(4): 437-48.
20. McCredie M, Little J, Cotton S, Mueller B, Peris-Bonet R, Choi NW, et al. SEARCH international case-control study of childhood brain tumours: role of index pregnancy and birth, and mother's reproductive history. *Paediatr Perinat Epidemiol*. 1999; 13(3): 325-41.
21. McKinney PA, Juszczak E, Findlay E, Smith K and Thomson CS. Pre- and perinatal risk factors for childhood leukaemia and other malignancies: a Scottish case control study. *Br J Cancer*. 1999; 80(11): 1844-51.
22. Schuz J, Kaatsch P, Kaletsch U, Meinert R and Michaelis J. Association of childhood cancer with factors related to pregnancy and birth. *Int J Epidemiol* 1999; 28(4): 631-9.
23. Schuz J, Kaletsch U, Kaatsch P, Meinert R and Michaelis J. Risk factors for pediatric tumours of the central nervous system: results from a German population-based case-control study. *Med Pediatr Oncol*. 2001; 36(2): 274-82.
24. Mogren I, Malmer B, Tavelin B and Damber L. Reproductive factors have low impact on the risk of different primary brain tumours in offspring. *Neuroepidemiology*. 2003; 22(4): 249-54.
25. Von Behren J and Reynolds P. Birth characteristics and brain cancers in young children. *Int J Epidemiol* 2003; 32(2): 248-56.
26. Hjalgrim LL, Westergaard T, Rostgaard K, Schmiegelow K, Melbye M, Hjalgrim H, et al. Birth weight as a risk factor for childhood leukemia: a meta-analysis of 18 epidemiologic studies. *Am J Epidemiol*. 2003; 158(8): 724-35.
27. Hjalgrim LL, Rostgaard K, Hjalgrim H, Westergaard T, Thomassen H, Forestier E, et al. Birth weight and risk for childhood leukemia in Denmark, Sweden, Norway, and Iceland. *J Natl Cancer Inst*. 2004; 96(20): 1549-56.
28. Roman E, Simpson J, Ansell P, Lightfoot T, Mitchell C and Eden TO. Perinatal and reproductive factors: a report on haematological malignancies from the UKCCS. *Eur J Cancer*. 2005; 41(5): 749-59.
29. McLaughlin CC, Baptiste MS, Schymura MJ, Nasca PC and Zdeb MS. Birth weight, maternal weight and childhood leukaemia. *Br J Cancer*. 2006; 94(11): 1738-44.

- 373 30. Podvin D, Kuehn CM, Mueller BA and Williams M. Maternal and birth characteristics in  
374 relation to childhood leukaemia. *Paediatr Perinat Epidemiol.* 2006; 20(4): 312-22.
- 375 31. Dorak MT, Pearce MS, Hammal DM, McNally RJ and Parker L. Examination of gender  
376 effect in birth weight and miscarriage associations with childhood cancer (United Kingdom).  
377 *Cancer Causes Control.* 2007; 18(2):219-28.
- 378 32. Ross JA, Perentesis JP, Robison LL and Davies SM. Big babies and infant leukemia: a  
379 role for insulin-like growth factor-1? *Cancer Causes Control.* 1996; 7(5): 553-9.
- 380 33. Preston-Martin S, Yu MC, Benton B and Henderson BE. N-Nitroso compounds and  
381 childhood brain tumours: a case-control study. *Cancer Res.* 1982; 42(12): 5240-5.
- 382 34. Daling JR, Starzyk P, Olshan AF and Weiss NS. Birth weight and the incidence of  
383 childhood cancer. *J Natl Cancer Inst.* 1984; 72(5): 1039-41.
- 384 35. Howe GR, Burch JD, Chiarelli AM, Risch HA and Choi BC. An exploratory case-control  
385 study of brain tumours in children. *Cancer Res.* 1989; 49(15): 4349-52.
- 386 36. Yeazel MW, Ross JA, Buckley JD, Woods WG, Ruccione K and Robison LL. High birth  
387 weight and risk of specific childhood cancers: a report from the Children's Cancer Group. *J*  
388 *Pediatr.* 1997; 131(5): 671-7.
- 389 37. Hardell L and Dreifaldt AC. Breast-feeding duration and the risk of malignant diseases in  
390 childhood in Sweden. *Eur J Clin Nutr.* 2001; 55(3): 179-85.
- 391 38. Kuijten RR, Bunin GR, Nass CC and Meadows AT. Gestational and familial risk factors  
392 for childhood astrocytoma: results of a case-control study. *Cancer Res.* 1990; 50(9): 2608-12.
- 393 39. Heuch JM, Heuch I, Akslen LA and Kvale G. Risk of primary childhood brain tumours  
394 related to birth characteristics: a Norwegian prospective study. *Int J Cancer.* 1998; 77(4): 498-  
395 503.
- 396 40. Smulevich VB, Solionova LG, and Belyakova SV. Parental occupation and other factors  
397 and cancer risk in children: I. Study methodology and non-occupational factors. *Int J Cancer.*  
398 1999; 83(6): 712-7.
- 399 41. Fear NT, Roman E, Ansell P and Bull D. Malignant neoplasms of the brain during  
400 childhood: the role of prenatal and neonatal factors (United Kingdom). *Cancer Causes Control.*  
401 2001; 12(5): 443-9.

- 402 42. Mellemkjaer L, Hasle H, Gridley G, Johansen C, Kjaer SK, Frederiksen K, et al. H. Risk  
403 of cancer in children with the diagnosis immaturity at birth. *Paediatr Perinat Epidemiol*. 2006;  
404 20(3): 231-7.
- 405 43. Martin RM, Gunnell D, Owen CG and Smith GD. Breast-feeding and childhood cancer:  
406 A systematic review with metaanalysis. *Int J Cancer*. 2005; 117(6): 1020-1031.
- 407 44. Windham GC, Bjerkedal T, and Langmark F. A population-based study of cancer  
408 incidence in twins and in children with congenital malformations or low birth weight, Norway,  
409 1967-1980. *Am J Epidemiol*. 1985; 121(1): 49-56.
- 410 45. Mili F, Khoury MJ, Flanders WD and Greenberg RS. Risk of childhood cancer for infants  
411 with birth defects. I. A record-linkage study, Atlanta, Georgia, 1968-1988. *Am J Epidemiol*.  
412 1993; 137(6): 629-38.
- 413 46. Narod SA, Hawkins MM, Robertson CM and Stiller CA. Congenital anomalies and  
414 childhood cancer in Great Britain. *Am J Hum Genet*. 1997; 60(3): 474-85.
- 415 47. Klip H, Burger CW, de Kraker J and van Leeuwen FE. Risk of cancer in the offspring of  
416 women who underwent ovarian stimulation for IVF. *Hum Reprod*. 2001; 16(11): 2451-8.
- 417 48. Brinton LA, Kruger Kjaer S, Thomsen BL, Sharif HF, Graubard BI, Olsen JH, et al.  
418 Childhood tumour risk after treatment with ovulation-stimulating drugs. *Fertil Steril*. 2004;  
419 81(4): 1083-91.
- 420 49. Doyle P, Bunch KJ, Beral V and Draper GJ. Cancer incidence in children conceived with  
421 assisted reproduction technology. *Lancet*. 1998; 352(9126): 452-3.
- 422 50. Bergh T, Ericson A, Hillensjo T, Nygren KG and Wennerholm UB. Deliveries and  
423 children born after in-vitro fertilisation in Sweden 1982-95: a retrospective cohort study. *Lancet*.  
424 1999; 354(9190): 1579-85.
- 425 51. Bruinsma F, Venn A, Lancaster P, Speirs A and Healy D. Incidence of cancer in children  
426 born after in-vitro fertilization. *Hum Reprod*. 2000; 15(3): 604-7.
- 427 52. Lerner-Geva L, Toren A, Chetrit A, Modan B, Mandel M, Rechavi G, et al. The risk for  
428 cancer among children of women who underwent in vitro fertilization. *Cancer*. 2000; 88(12):  
429 2845-7.



**Table I:** Characteristics of the cases and controls

	<b>Cases n=209 (%)</b>	<b>Controls n=1681 (%)</b>	<b>Co/Ca<sup>(a)</sup> ratio</b>	<b>p</b>
<b>Histological subtypes</b>				
Embryonal tumours	100 (48)			
Ependymomas	33 (16)			
Astrocytomas	26 (12)			
Other gliomas	45 (21)			
Other specified tumors	5 (3)			
<b>Gender</b>				ns <sup>(b)</sup>
Male	125 (60)	932 (55)		
<b>Age at the reference date (years)</b>				0.03 <sup>(b)</sup>
< 2	34 (16)	369 (22)	10.5	
2	15 (7)	153 (9)	10.2	
3	17 (8)	166 (10)	9.8	
4	25 (12)	145 (9)	5.6	
5-6	34 (16)	228 (14)	6.5	
7-8	33 (16)	163 (10)	4.8	
9-11	27 (13)	225 (13)	8.3	
12-14	24 (11)	232 (14)	9.7	
<b>Parent's marital status at interview</b>				ns <sup>(c)</sup>
Married	191 (92)	1541 (92)		
Separated, divorced, widowed	4 (2)	65 (4)		
Single	14 (7)	75 (5)		
<b>Paternal educational level</b>				ns <sup>(c)</sup>
≤ High school	126 (60)	1063 (63)		
> High school	79 (38)	601 (36)		
missing data	4 (2)	17 (1)		
<b>Maternal educational level</b>				ns <sup>(c)</sup>
≤ High school	125 (60)	979 (58)		
> High school	84 (40)	701 (42)		
<b>Socioprofessional categories</b>				ns <sup>(c)</sup>
Intellectual and scientific jobs, intermediate profession	92 (44)	713 (42)		
Administrative employees and sales workers	51 (24)	477 (28)		
Service workers	29 (13)	215 (13)		
Farmers, agricultural, craftsmen and factory workers	37 (18)	274 (16)		

<sup>(a)</sup> Co/Ca ratio: Controls/Cases ratio<sup>(b)</sup> Chi square test, ns: p>0.05<sup>(c)</sup> Non-conditional logistic regression adjusted for age and gender

432 **Table II:** Mother's obstetrical history and childhood central nervous tumours

	Controls n=1681 (%)		All CNS tumours n=209 (%)				Ependymomas (n=33)			Embryonal tumours (n=100)			Astrocytomas (n=26)			Other gliomas (n=45)		
	n	(%)	n	(%)	OR	[IC <sub>95%</sub> ] <sup>(1)</sup>	n	OR	[IC <sub>95%</sub> ] <sup>(2)</sup>	n	OR	[IC <sub>95%</sub> ] <sup>(2)</sup>	n	OR	[IC <sub>95%</sub> ] <sup>(2)</sup>	n	OR	[IC <sub>95%</sub> ] <sup>(2)</sup>
All foetal losses																		
No	1031	(61)	115	(55)	1.0	ref	20	1.0	ref	57	1.0	ref	12	1.0	ref	23	1.0	ref
Yes	650	(39)	94	(45)	1.3	[1.0-1.7]	13	1.1	[0.5-2.2]	43	1.2	[0.8-1.8]	14	1.8	[0.8-4.0]	22	1.5	[0.8-2.6]
1	403	(24)	58	(28)	1.3	[0.9-1.8]	8	1.1	[0.5-2.5]	26	1.2	[0.7-1.9]	10	2.2	[0.9-5.0]	12	1.3	[0.6-2.7]
2	247	(15)	36	(17)	1.3	[0.9-1.9]	5	1.1	[0.4-3.0]	17	1.2	[0.7-2.1]	4	1.3	[0.4-4.2]	10	1.7	[0.8-3.6]
Miscarriages																		
No	1239	(74)	142	(68)	1.0	ref	29	1.0	ref	68	1.0	ref	16	1.0	ref	26	1.0	ref
Yes	442	(26)	67	(32)	1.3	[1.0-1.8]	4	0.4	[0.1-1.1]	32	1.3	[0.9-2.0]	10	1.7	[0.8-3.8]	19	2.0	[1.1-3.6]
1	303	(18)	50	(24)	1.4*	[1.0-2.0]	2	0.3	[0.1-1.2]	24	1.4	[0.9-2.3]	10	2.5	[1.1-5.6]	12	1.8	[0.9-3.6]
≥ 2	138	(8)	17	(8)	1.1	[0.6-1.8]	2	0.6	[0.1-2.6]	8	1.1	[0.5-2.3]	0	-	-	7	2.4	[1.0-5.5]
Voluntary abortion																		
No	1429	(85)	173	(83)	1.0	ref	23	1.0	ref	85	1.0	ref	21	1.0	ref	39	1.0	ref
Yes	252	(15)	36	(17)	1.2	[0.8-1.7]	10	2.6	[1.2-5.7]	15	1.0	[0.5-1.7]	5	1.4	[0.5-3.8]	6	0.8	[0.4-2.0]
1	209	(12)	27	(13)	1.1	[0.7-1.6]	6	1.9	[0.7-4.6]	12	0.9	[0.5-1.8]	4	1.4	[0.5-4.1]	5	0.9	[0.3-2.2]
≥ 2	43	(3)	9	(4)	1.7	[0.8-3.6]	4	7.2	[2.3- 22]	3	1.1	[0.3-3.7]	1	1.5	[0.2- 12]	1	0.8	[0.1-5.9]
Therapeutic abortion																		
No	1594	(95)	203	(97)	1.0	ref	33	1.0	ref	97	1.0	ref	24	1.0	ref	44	1.0	ref
Yes	87	(5)	6	(3)	0.5	[0.2-1.2]	0	-	-	3	0.5	[0.2-1.8]	2	1.4	[0.3-6.2]	1	0.4	[0.1-2.8]
Assisted reproduction for the index child																		
No	1599	(95)	198	(95)	1.0	ref	29	1.0	ref	97	1.0	ref	25	1.0	ref	43	1.0	ref
Yes	82	(5)	11	(5)	1.1	[0.6-2.2]	4	2.6	[0.9-7.8]	3	0.6	[0.2-2.1]	1	0.8	[0.1-6.3]	2	0.9	[0.2-3.9]

433 <sup>(1)</sup> Non-conditional logistic regression adjusted on age and gender, <sup>(2)</sup> Non-conditional polytomic regression adjusted on age and gender

434 \* p&lt;0.05

435 **Table III:** Characteristics of children at birth and risk of childhood malignant central nervous tumours

	<b>Controls</b> <b>n=1681</b> <b>(%)</b>		<b>All CNS tumours</b> <b>n=209 (%)</b>				<b>Ependymomas</b> <b>(n=33)</b>			<b>Embryonal</b> <b>tumours</b> <b>(n=100)</b>			<b>Astrocytomas</b> <b>(n=26)</b>			<b>Other gliomas</b> <b>(n=45)</b>		
	n	(%)	n	(%)	OR	[IC <sub>95%</sub> ] <sup>(1)</sup>	n	OR	[IC <sub>95%</sub> ] <sup>(2)</sup>	n	OR	[IC <sub>95%</sub> ] <sup>(2)</sup>	n	OR	[IC <sub>95%</sub> ] <sup>(2)</sup>	n	OR	[IC <sub>95%</sub> ] <sup>(2)</sup>
<b>Gestational age</b>																		
< 37 weeks	107	(6)	20	(10)	<b>1.6</b>	[0.8-2.9]	3	<b>1.5</b>	[0.4-6.2]	13	<b>2.5</b>	[1.1-5.4]	1	<b>0.4</b>	[0.1-3.6]	2	<b>0.6</b>	[0.1-3.1]
37-41 weeks	1515	(91)	177	(87)	<b>1.0</b>	ref	29	<b>1.0</b>	ref	83	<b>1.0</b>	ref	22	<b>1.0</b>	ref	39	<b>1.0</b>	ref
> 41 weeks	40	(2)	6	(3)	<b>1.4</b>	[0.6-3.3]	1	<b>1.4</b>	[0.2- 11]	1	<b>0.5</b>	[0.1-3.8]	1	<b>1.9</b>	[0.2- 15]	3	<b>2.7</b>	[0.8-9.3]
<b>Birth weight</b>																		
< 2500	94	(6)	16	(7)	<b>1.0</b>	[0.5-2.5]	2	<b>0.8</b>	[0.1-4.3]	8	<b>0.9</b>	[0.3-2.3]	3	<b>2.2</b>	[0.4- 11]	3	<b>1.7</b>	[0.4-7.1]
2500-4000	1451	(86)	178	(85)	<b>1.0</b>	ref	29	<b>1.0</b>	ref	83	<b>1.0</b>	ref	23	<b>1.0</b>	ref	38	<b>1.0</b>	ref
> 4000	135	(8)	15	(7)	<b>1.0</b>	[0.5-1.7]	2	<b>0.8</b>	[0.2-3.3]	9	<b>1.3</b>	[0.6-2.6]	0	-	-	4	<b>1.2</b>	[0.4-3.5]
<b>Paternal Age</b>																		
< 25	69	(4)	8	(4)	<b>1.0</b>	[0.4-2.0]	0	-	-	5	<b>1.3</b>	[0.5-3.3]	2	<b>1.9</b>	[0.4-8.4]	1	<b>0.5</b>	[0.1-3.9]
25-34	1109	(66)	135	(64)	<b>1.0</b>	ref	22	<b>1.0</b>	ref	63	<b>1.0</b>	ref	17	<b>1.0</b>	ref	30	<b>1.0</b>	ref
≥ 35	503	(30)	65	(32)	<b>1.1</b>	[0.8-1.5]	11	<b>1.0</b>	[0.5-2.1]	32	<b>1.2</b>	[0.7-1.8]	7	<b>1.0</b>	[0.4-2.4]	14	<b>1.1</b>	[0.6-2.1]
<b>Maternal Age</b>																		
< 25	163	(10)	17	(8)	<b>0.8</b>	[0.5-1.4]	2	<b>0.7</b>	[0.2-2.9]	7	<b>0.7</b>	[0.3-1.6]	3	<b>1.2</b>	[0.3-4.1]	5	<b>1.1</b>	[0.4-2.9]
25-34	1235	(73)	158	(76)	<b>1.0</b>	ref	25	<b>1.0</b>	ref	76	<b>1.0</b>	ref	19	<b>1.0</b>	ref	34	<b>1.0</b>	ref
≥ 35	283	(17)	34	(16)	<b>1.0</b>	[0.6-1.4]	6	<b>1.0</b>	[0.4-2.4]	17	<b>1.0</b>	[0.6-1.8]	4	<b>1.0</b>	[0.3-2.9]	6	<b>0.8</b>	[0.3-2.0]
<b>Birth order</b>																		
1	708	(42)	80	(39)	<b>1.0</b>	ref	13	<b>1.0</b>	ref	31	<b>1.0</b>	ref	11	<b>1.0</b>	ref	22	<b>1.0</b>	ref
2	608	(36)	94	(44)	<b>1.3</b>	[0.9-1.8]	15	<b>1.3</b>	[0.6-2.8]	52	<b>1.9</b>	[1.2-2.9]	13	<b>1.3</b>	[0.6-3.0]	14	<b>0.7</b>	[0.4-1.4]

3 +	365	(22)	35	(17)	<b>0.8</b>	[0.5-1.2]	5	<b>0.7</b>	[0.3-2.0]	17	<b>1.0</b>	[0.6-1.9]	2	<b>0.4</b>	[0.1-1.6]	9	<b>0.8</b>	[0.3-1.7]
<b>Breast feeding<sup>(3)</sup></b>																		
Never	775	(49)	93	(45)	<b>1.0</b>	<b>ref</b>	16	<b>1.0</b>	<b>ref</b>	43	<b>1.0</b>	<b>ref</b>	12	<b>1.0</b>	<b>ref</b>	19	<b>1.0</b>	<b>ref</b>
Ever	816	(51)	112	(55)	<b>1.2</b>	[0.9-1.6]	16	<b>0.8</b>	[0.4-1.7]	55	<b>1.3</b>	[0.8-1.9]	13	<b>1.1</b>	[0.5-2.5]	26	<b>1.4</b>	[0.8-2.6]
Breast & Bottle	92	(6)	10	(5)	<b>0.9</b>	[0.5-1.9]	1	<b>0.5</b>	[0.1-3.6]	6	<b>1.3</b>	[0.5-3.1]	2	<b>1.5</b>	[0.3-7.0]	1	<b>0.5</b>	[0.1-3.5]
Breast only	724	(45)	102	(50)	<b>1.2</b>	[0.9-1.6]	15	<b>0.9</b>	[0.4-1.8]	49	<b>1.3</b>	[0.8-2.0]	11	<b>1.1</b>	[0.5-2.5]	25	<b>1.5</b>	[0.8-2.8]
<b>Duration</b>																		
≤ 3 months	484	(30)	67	(33)	<b>1.2</b>	[0.5-1.9]	9	<b>0.8</b>	[0.4-1.9]	28	<b>1.1</b>	[0.7-1.8]	10	<b>1.4</b>	[0.6-3.4]	19	<b>1.7</b>	[0.9-3.2]
> 3 months	240	(15)	35	(17)	<b>1.3</b>	[0.8-1.7]	6	<b>1.0</b>	[0.4-2.6]	21	<b>1.7</b>	[1.0-2.9]	1	<b>0.3</b>	[0.1-2.4]	6	<b>1.2</b>	[0.5-3.0]
<b>Congenital malformations</b>																		
No	1624	(97)	201	(96)	<b>1.0</b>	<b>ref</b>	32	<b>1.0</b>	<b>ref</b>	97	<b>1.0</b>	<b>ref</b>	25	<b>1.0</b>	<b>ref</b>	42	<b>1.0</b>	<b>ref</b>
Yes	57	(3)	8	(4)	<b>1.1</b>	[0.5-2.3]	1	<b>1.0</b>	[0.1-7.2]	3	<b>0.8</b>	[0.2-2.6]	1	<b>1.2</b>	[0.2-9.2]	3	<b>2.0</b>	[0.6-6.6]
Major	38	(2)	6	(3)	<b>1.2</b>	[0.5-2.9]	0	-	-	2	<b>0.8</b>	[0.2-3.5]	1	<b>1.9</b>	[0.2- 14]	3	<b>2.9</b>	[0.9- 10]

436 <sup>(1)</sup> Non-conditional logistic regression adjusted on age and gender; <sup>(2)</sup> Non-conditional polytomic regression adjusted on age and  
437 gender; <sup>(3)</sup> Analysis were performed in children older than 6 months