



Effect of hypocaloric diet-induced weight loss in obese women on plasma apelin and adipose tissue expression of apelin and APJ.

Isabelle Castan-Laurell, Michaela Vítková, Danièle Daviaud, Cédric Dray,
Michaela Kovacikova, Zuzana Kováčová, Jindriska Hejnová, Vladimir Stich,
Philippe Valet

► To cite this version:

Isabelle Castan-Laurell, Michaela Vítková, Danièle Daviaud, Cédric Dray, Michaela Kovacikova, et al.. Effect of hypocaloric diet-induced weight loss in obese women on plasma apelin and adipose tissue expression of apelin and APJ.. European Journal of Endocrinology, 2008, 158 (6), pp.905-10. 10.1530/EJE-08-0039 . inserm-00277309

HAL Id: inserm-00277309

<https://inserm.hal.science/inserm-00277309>

Submitted on 7 Apr 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TABLE 1. Clinical characteristics and metabolic parameters of lean controls and obese subjects before and after LCD.

	Obese		P
	Before diet	After diet	
Weight (kg)	89.8 ± 3.7	83.1 ± 3.6	<0.0001
BMI (kg/m ²)	32.2 ± 1.4	29.8 ± 1.4	<0.0001
Fat mass (%)	41.3 ± 1.1	39.0 ± 1.4	<0.001
Fat (kg)	37.8 ± 2.6	33.3 ± 2.7	<0.0001
Waist circumference (cm)	96.4 ± 3.1	88.8 ± 2.8	<0.0001
Waist-to-hip ratio (WHR)	0.81 ± 0.02	0.79 ± 0.01	<0.005
Systolic blood pressure (mm Hg)	116 ± 2	115 ± 3	0.260
Diastolic blood pressure (mm Hg)	73 ± 2	72 ± 2	0.522
Cholesterol (mmol/L)	5.76 ± 0.23	5.27 ± 0.18	<0.01
Triglycerides (mmol/L)	1.28 ± 0.09	1.00 ± 0.07	<0.005
HDL-C (mmol/L)	1.70 ± 0.07	1.51 ± 0.05	<0.001
Glucose (mmol/L)	5.14 ± 0.10	4.97 ± 0.09	0.0862
HOMA-IR	1.89 ± 0.18	1.47 ± 0.16	<0.01

The data presented are means ± SEM in obese subjects before (n= 20) and after (n= 20) LCD.

P: probabilities obtained with nonparametric Wilcoxon's test for paired observations (before and after LCD)