

**Table 1.** Short review of studies assessing occupational exposure among nuclear workers by using job-exposure matrix approach

Reference	Industry	Exposure	Exposure estimator	Exposure assessment method	Purpose, study design and main results	Reviewer's comments
Country	Period					
Eheman et al., 1999	All involving radiation	External radiation dose	Discrete annual dose distribution with 6 dose categories, Distribution of cumulative dose,	Published dosimetry data for different occupational groups	To estimate radiation doses and uncertainty for individuals reported	Use of radioprotection dosimetry data. No assessment of internal radiation exposure
Germany	exposure		Monte Carlo method geometric mean and standard deviation			
1960-80						
Krishnamadasan et al., 2007	Rocket engine and nuclear power testing facility	TCE, benzene, hydrazine, PHA, mineral oils	Exposure score for each chemical= intensity of exposure (4-level scale) by 3 time periods x duration of employment	Coding of likelihood and intensity of exposure by job title by hygienist based on knowledge from facility records and workers survey and interviews	To assess a relationship between occupational exposure to chemicals in the Nested Case-Control study of prostate cancer incidence	Available data on life style habits and familial history of prostate cancer, occupational physical activity intensity.
USA	1950-90					
					Dose-response relationship with TCE	Study with etiological issue

					exposure
Krishnadasan et al., 2008	Rocket engine and nuclear power testing facility	Occupational physical activity	Score of occupational physical activity based on intensity (3-level scale) of occupational physical activity for a job held longest x duration of employment	Coding of intensity of physical activity by job title by hygienist based on knowledge from facility records and workers survey and interviews	To assess a relationship between occupational physical activity in the Nested Case-Control study of prostate cancer incidence
USA	1950-90				Job titles used as an entry to JEM despite discrepancies in physical activity level results
Boice et al., 2006	Rocket engine testing facility SSFL	TCE and hydrazine. Other chemicals as surrogate for exposure to all other chemicals	Potential of exposure (no, possible and likely exposure) and cumulative potential exposure for hydrazine and TCE.	Review of worker's exposure history, year, place, and type of work. Validation based on walkthrough visits, interviews with workers, review of workers' medical records	Only potential of exposure assessed by crude JEM. Comprehensive review of TCE and hydrazine use at the facility.
USA	1948-1999		Duration of employment as a test stand mechanic		relationship between TCE and kidney cancer. No relationship between test

				for other chemicals	stand mechanic job and mortality.
Ritz et al., 1999a USA	Rocket engine testing facility SSFL*	Hydrazine and other chemicals 1950-90	Relative intensity of presumptive exposure (4-level scale) by 3 time periods. Time-dependent cumulative exposure score 1950-90	3 experts consensus based on walkthrough visits, interviews with workers, review of historical facility reports	Retrospective cohort study of cancer incidence and mortality. Dose-response relationship for lung, colorectal, and pancreatic cancer.
Rooney et al., 1993 UK	UK Atomic Energy Authority	13 specific radionuclides 6 metals: Be, Br, Cd, Pb, Hg, Zn; 3 types of chemicals:	For each radionuclides - Level of exposure (none, possible, probable but relatively low, probable but relatively high).	Review of worker's exposure history, year, place, and type of work. 125 work areas classified by health physicists or experienced staff in 7 work environments: Reactor, reactor maintenance, fuel examination after irradiation, fuel fabrication, decontamination and was disposal and laundry, fuel reprocessing.	No details on JEM. No reporting of exposure results
		aromatic, halogenated, other organic compounds; asbestos, metal dusts, metal fumes	For other agents, level of exposure on 3 level scale (none, low, high) Duration of work	No significant findings according to other kinds of exposure	

Records of internal contamination						
Carpenter et al., 1988	Y-12 and ORNL USA	26 chemicals or chemical groups including Be, U, Th, Carbon dusts, welding fumes, cutting oils, hydrazine	Rank of potential exposure to each chemical (0-no, 1-low, 2-moderate, 3-high potential for exposure).	Subjective evaluation by industrial hygienist of job title/department combination with accounting for time period, exposure duration=duration review of industrial processes, of employment in job titles on-site visits, interview with people, urinalysis and air monitoring data.	Nested case-control of primary CNS cancer. Analyses according to exposure levels. No reference on exposure evaluation report.	No cumulative exposure assessment. No results according to exposure levels. No reference on exposure evaluation report.
Wing et al., 1993	ORNL USA	Be, Hg, Pb	15 Job categories based on potential for similar occupational environment and activities.	Employment records: Job title, department, payment status and monitoring status for Be, Hg, Pb exposure Time spent in job category	To consider the role of possible selection and confounding factors on dose-risk estimators in previous cohort study of solid cancer and leukaemia. No significant changes on dose-risk estimates except Hg	

							Good description of
Ritz, 1999b	Fernald	TCE, kerosene and	Exposure level (0 to 3) and	Plant experts: a plant foreman,	Cohort study of chemical		
USA	FFMPC	cutting oils	duration of exposure (with	hygienists, an engineer	exposure effects on	highly exposed jobs,	
			15 year lag)	determined likelihood of	cancer mortality with	jobs with concomitant	
			Internal and external	chemical exposure for each job	accounting for radiation	exposure, cutting oils	
			radiation dose	title and plant area for	exposure for specific	composition and	
					cancer sites.	definition discussed.	
				1952-77 period			
Ritz at al.,	Rocketyne	Internal lung dose	Exposure level (0 to 3)	? Job titles, employment	Cohort study of internal	Results adjusted for	
2000	1950-94	Asbestos	defined on job titles.	periods, and job locations used	radiation exposure effect	chemical exposure are	
USA	Hydrazine	Internal and external	as proxy measures for chemical	on mortality from specific	not shown.		
		radiation dose	exposures	cancer sites. Dose-			
				response relationship with			
				internal radiation			
Rutterber et	Rocky Flats	Be, CCl4, Cr, Pb, Ni,	Time-weighted average	Published estimates.	Improving estimates of	Validation of Be	
al., 2001a	Nuclear	TCE, PCE,	(max/min) annual air	For Be, data from a 100s	exposure for	exposure estimators.	
USA	weapon	formaldhyde,	exposure	personnel air samplers.			
	facility	methylene chloride,		Estimates for concentration in	epidemiological studies of	Results in % and Nb of	
		asbestos		breathing zone of workers,	plutonium workers	workers exposed to each	
						agent.	

					Published estimates for similar	To develop a JEM for	No reference of
Ruttenber et al., 2001b	Rocky Flats	Be, CCl4, Cr, Pb, Ni, Time-weighted average	(max/min) daily air concentration.	production process. Hours worked paid/year from employer's roster.	epidemiological studies and for determination of eligibility for a medical screening program for highly exposed former workers	epidemiological studies and for determination of exposure levels.	published estimates of exposure levels.
USA	Nuclear weapon facility	TCE, PCE, formaldehyde, methylene chloride, asbestos	Cumulative exposure for period of interest=AV. concentration x Nb hours worked			No detail on final nb of job groups, on mostly exposed jobs, on 10/20 selected chemicals. Solid estimators allowing accounting for uncertainties. Reporting of exposures according to LOEL	
Henn et al., 2007	Chemical laboratories at Y-12, X-10, K-25 and SRS plants	Organic and inorganic chemicals including radioisotopes	Individual cumulative potential exposure indices (PEIs)=Nb of days with task- and time-dependent combination x task- and time-dependent weighting factors	Task-dependent weighting factors determined by industrial hygienists (IH) through jobs review and interviews with workers. Time-dependent weighting factors	To assess chemical exposure for a mortality study of 6157 chemical laboratory workers	PEIs are not specific of any chemical. Waiting factor for time were established on the basis of exposure trends data available for only 4 chemicals from 54	

considered

**Table 2.** Summary of exposure agents evaluated in the job exposure matrix and exposure characteristics of the AREVA NC Pierrelatte plant workers (N=2709)

Exposure agents	Detail	Exposed workers n (%)	Individual cumulative exposure score in exposed workers	Mean ± SD (Median)
1-Natural U compounds f*	UF <sub>6</sub> , UF <sub>4</sub> , UO <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> , (UO <sub>4</sub> , nH <sub>2</sub> O)	2253 (83.23)	67.49 ± 65.32 (44.50)	
2-Natural U compounds m*	(U <sub>2</sub> O <sub>7</sub> )(NH <sub>4</sub> ) <sub>2</sub> , U <sub>3</sub> O <sub>8</sub> , UO <sub>2</sub> F <sub>2</sub> , UO <sub>3</sub>	1815 (67.05)	42.93 ± 51.01 (23.21)	
3-Natural U compounds s*	UO <sub>2</sub>	992 (36.65)	42.22 ± 40.52 (16.40)	
4-Reprocessed U compounds f*	UF <sub>6</sub> , UF <sub>4</sub> , UO <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> , (UO <sub>4</sub> , nH <sub>2</sub> O)	851 (31.44)	31.89 ± 40.52 (16.27)	
5-Reprocessed U compounds m*	(U <sub>2</sub> O <sub>7</sub> )(NH <sub>4</sub> ) <sub>2</sub> , U <sub>3</sub> O <sub>8</sub> , UO <sub>2</sub> F <sub>2</sub> , UO <sub>3</sub>	656 (24.23)	36.62 ± 43.32 (20.53)	
6-Reprocessed U compounds s*	UO <sub>2</sub>	475 (17.55)	35.40 ± 43.76 (17.95)	
7-Chlorinated agents	Perchloroethylene, tetrachloroethene, trichloroethylene, dichloromethane, polychlorinated biphenyls (PCBs), carbon tetrachloride	1784 (65.90)	18.49 ± 21.72 (13.37)	
8-Fluoride agents	Fluorhydric acid, tungsten hexafluoride, fluorine, potassium fluoride	1652 (61.03)	28.47 ± 42.39 (13.37)	
9-Nitrogenous agents	Ammonia, ammonia anhydride, nitrogen acid, nitrous	1415 (47.80)	31.55 ± 45.60 (12.04)	

	vapours		
10- Solvents containing aromatic hydrocarbons	Benzene, toluene, xylene, styrene	1255 (46.36)	30.41 ± 39.90 (15.26)
11-Welding fumes	Fumes and metal dusts	425 (15.70)	12.07 ± 14.57 (7.78)
12-Vitreous fibres and rock wool	Rock, slag, and glass wools	1280 (47.28)	21.05 ± 26.05 (10.21)
13-Asbestos	Asbestos fireproofing, insulation and braids	1894 (69.97)	19.28 ± 18.73 (15.94)
14-Refractive ceramic fibres		555 (20.50)	9.98 ± 7.93 (8.56)
15-Chromate	Potassium dichromate and chromium trioxide	102 (3.77)	7.56 ± 6.18 (6.42)
16- Chlorine trifluoride		1341 (49.54)	14.94 ± 15.35 (10.04)
17-Trichloroethylene (TCE)		1685 (62.25)	27.27 ± 33.57 (15.25)
18-Lead	Paints, plates and dust	331 (12.23)	8.49 ± 8.32 (6.46)
19-Mercury	Vapours	827 (30.55)	16.46 ± 23.19 (8.96)
20-Silica	Silica gel, silica grains	695 (25.67)	8.71 ± 8.06 (6.73)
21-Hydrazine and other fuels	Hydrazine, domestic fuel, gas-oils, four-star fuel, petroleum	1164 (42.99)	16.18 ± 19.89 (10.29)
22-Heat	Temperature at the workstation > 30°C	2361 (87.22)	54.63 ± 53.71 (34.47)

\* Uranium compounds were classified in terms of absorption types (f-fast, m-moderate or s-slow) according to the Human

Respiratory Tract Model described in ICRP Publication 66 (32)

**Table 3.** AREVA NC Pierrelatte job exposure matrix validity characteristics compared to  
AREVA NC Pierrelatte workers' medical records used as reference

Exposure category	Observed agreement coefficient	Kappa coefficient	Sensitivity	Specificity
Uranium compounds derived from NU*	0.85	0.66	0.81	0.96
Uranium compounds derived from RPU**	0.92	0.83	0.82	0.98
Asbestos	0.61	0.09	1.00	0.60
Vitreous fibres and wools	0.74	0.27	0.50	0.80
Chlorinated agents	0.71	0.42	0.77	0.67
Fluoride agents	0.58	0.15	0.57	0.60
Nitrogenous agents	0.68	0.36	0.59	0.79
Total	0.78	0.56	0.72	0.83

\* Natural uranium

\*\* Reprocessed uranium

**Table 4.** Generic jobs with the highest cumulated exposure levels for more than 3 categories of exposure agent.

Generic Jobs	Exposure agents with the highest scores
Dismantling-decontaminating technician	Compounds derived from RPUI*
	Compounds derived from RPUM*
	Compounds derived from RPUS*
TCE	
Fluorinated compounds	
Mercury	
Lead	
Refractive ceramic fibres	
Driver at enriched materials chemistry unit	Compounds derived from NUU*
	Compounds derived from NUm*
	Compounds derived from NUJ*
TCE	
Fluorinated compounds	
Heat	
Mechanic at enriched materials chemistry unit	Compounds derived from NUJ*
	Hydrazine and other fuels
	Welding fumes
	Wools
Technician and operator at container	Heat
	Compounds derived from RPUI*

maintenance shop	Compounds derived from RPUs*
	Compounds derived from RPUs*
	Fluorinated compounds
	Potassium dichromate
Physical-chemistry analysis technician at the plants' main laboratory	Mercury
	Nitrogenous compounds
	Silica gel, silica grains

\* Uranium compounds derived from natural (NU) and reprocessed uranium (RPU) were classified in terms of absorption types (f, m or s) according to (ICRP, 1994)

**Table 5.** Correlation between exposure to uranium compounds and other exposure agents. Values of Spearman's rho correlation coefficient (n=2709).

Exposure agents	Asbestos	Ceramic fibres	Chromate	Chloroethane	Chlorofluoride	Mercury	Lead	Silica	Hydrazin	Chlorinating agents	Fluoride agents	Nitrogenous solvents	Solvents	Welding fumes	Wools	Heat	
NUff**	<b>0.61*</b>	0.29*	0.10*	<b>0.72*</b>	0.40*	0.31*	0.25*	0.33*	0.24*	<b>0.56*</b>	<b>0.65*</b>	<b>0.52*</b>	0.43*	0.25*	0.26*	<b>0.66*</b>	
NUm	0.38*	<b>0.52*</b>	0.20*	0.38*	0.26*	<b>0.53*</b>	0.16*	0.40*	0.19*	0.42*	<b>0.55*</b>	<b>0.59*</b>	0.48*	0.36*	0.36*	0.47*	0.35*
NUs	0.21*	0.45*	0.31*	0.27*	0.10*	0.40*	0.23*	0.32*	0.17*	0.30*	<b>0.57*</b>	<b>0.65*</b>	<b>0.51*</b>	0.12*	0.24*	0.24*	0.08*
RPUf	0.03*	0.37*	0.34*	0.10*	-0.01	0.35*	0.16*	<b>0.57*</b>	0.16*	0.26*	0.37*	0.40*	0.27*	0.28*	0.19*	0.03*	
RPUm	-0.08*	0.44*	0.36*	-0.05	-0.04	0.08*	0.23*	0.35*	0.14*	0.15*	0.40*	0.46*	0.23*	0.12*	0.17*	0.01*	
RPUu	0.10*	0.37*	0.35*	0.16*	0.11*	0.26*	0.27*	0.31*	0.32*	0.22*	0.40*	0.47*	0.34*	0.03*	0.20*	0.07*	

\*Two-tailed significance test p value <0.01.

\*\* Uranium compounds derived from natural (NU) and reprocessed uranium (RPU) were classified in terms of absorption types (f, m or s)

Values of rho > 0.50 corresponding to a large correlation according to Cohen's criteria are bolded.