



Imaging E-selectin expression following traumatic brain injury in the rat using a targeted USPIO contrast agent.

Catherine Chapon, Florence Franconi, Franck Lacoeuille, François Hindré, Patrick Saulnier, Jean-Pierre Benoit, Jean-Jacques Le Jeune, Laurent Lemaire

► To cite this version:

Catherine Chapon, Florence Franconi, Franck Lacoeuille, François Hindré, Patrick Saulnier, et al.. Imaging E-selectin expression following traumatic brain injury in the rat using a targeted USPIO contrast agent.. Magnetic Resonance Materials in Physics, Biology and Medicine, 2009, 22 (3), pp.167-74. 10.1007/s10334-008-0161-z . inserm-00353311

HAL Id: inserm-00353311

<https://inserm.hal.science/inserm-00353311>

Submitted on 4 Jan 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Time (min)	Time course of signal reduction after BP-EG3		Time course of signal reduction after BP-IELLQAR	
	Contralateral cortex	Traumatized cortex	Contralateral cortex	Traumatized cortex
10	-30.3 ± 3.5 **	-25.8 ± 3.9 **	-34.2 ± 8.3 **	-30.1 ± 4.2 **
20	-23.3 ± 4.7 *	-16.7 ± 6.2 *	-30.1 ± 10.1 **	-32.4 ± 7.6 **
30	-17.4 ± 4.6 *	-18.3 ± 8.0 *	-13.1 ± 5.6 *	-29.7 ± 5.3 **
40	- 7.7 ± 4.8	-15.4 ± 8.8 *	- 3.8 ± 4.0	-23.4 ± 5.3 **
50	5.9 ± 3.0	- 3.1 ± 6.0	- 5.2 ± 3.5	-17.0 ± 6.4*
70	- 7.2 ± 2.8	- 5.9 ± 6.6	6.7 ± 2.5	-19.9 ± 3.6*

Table 1 : Variation of the T2* of traumatized and contralateral non-traumatized rat cortex after injection of targeted (BP-IELLQAR) or Non-targeted (BP-EG3) contrast agents expressed as the percentage of T2* value prior any injection

* P < 0.05 and ** P < 0.01 compared to initial T2* value of the considered tissue