



Primary biliary cirrhosis.

Xavier Roblin, Bruno Bonaz

► To cite this version:

Xavier Roblin, Bruno Bonaz. Primary biliary cirrhosis.. New England Journal of Medicine, 2005, 353 (25), 2719-20; author reply 2719-20. inserm-00410568

HAL Id: inserm-00410568

<https://inserm.hal.science/inserm-00410568>

Submitted on 10 Sep 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Primary Biliary Cirrhosis

TO THE EDITOR: The association between primary biliary cirrhosis and celiac disease should have been emphasized more clearly in the recent review of primary biliary cirrhosis. The prevalence of celiac disease among patients with primary biliary cirrhosis is 10 times as high as in the general population.¹ Moreover, Sorensen et al.² found an incidence ratio of primary biliary cirrhosis of 27.6 among patients with celiac disease. Thus, in the event of primary biliary cirrhosis, clinicians should systematically rule out celiac disease. Certain symptoms of primary biliary cirrhosis, such as osteoporosis, asthenia, and liver anomalies, regress with a gluten-free diet in people who also have celiac disease. This screening is made even simpler because there are sensitive and specific markers for each of the two diseases.

Xavier Roblin, M.D.
Bruno Bonaz, M.D., Ph.D.
Grenoble University
38000 Grenoble, France
xroblin@chu-grenoble.fr

1. Dickey W, McMillan SA, Callender ME. High prevalence of celiac sprue among patients with primary biliary cirrhosis. *J Clin Gastroenterol* 1997;25:328-32.
2. Sorensen HT, Thulstrup AM, Blomqvist P, Norgaard B, Fonager K, Ekbom A. Risk of primary biliary liver cirrhosis in patients with coeliac disease: Danish and Swedish cohort data. *Gut* 1999;44:736-8.