



**HAL**  
open science

## **AlphaIIbeta3 integrin: new allelic variants in Glanzmann thrombasthenia, effects on ITGA2B and ITGB3 mRNA splicing, expression, and structure-function.**

Vincent Jallu, Mathilde Dusseaux, Simon Panzer, Marie-Françoise Torchet, Nathalie Hezard, Jenny Goudemand, Alexandre de Brevern, Cécile Kaplan

### ► To cite this version:

Vincent Jallu, Mathilde Dusseaux, Simon Panzer, Marie-Françoise Torchet, Nathalie Hezard, et al.. AlphaIIbeta3 integrin: new allelic variants in Glanzmann thrombasthenia, effects on ITGA2B and ITGB3 mRNA splicing, expression, and structure-function.. *Human Mutation*, 2010, 31 (3), pp.237-46. 10.1002/humu.21179 . inserm-00448024

**HAL Id: inserm-00448024**

**<https://inserm.hal.science/inserm-00448024v1>**

Submitted on 3 Oct 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**$\alpha$ IIb $\beta$ 3 integrin : New allelic variants in Glanzmann Thrombasthenia,  
effects on mRNA splicing, expression and structure-function**

**V. JALLU<sup>1</sup>, M. DUSSEAUX<sup>1</sup>, S. PANZER<sup>2</sup>, M.F. TORCHET<sup>3</sup>, N. HEZARD<sup>4</sup>,  
J. GOUDEMANT<sup>5</sup>, A.G. de BREVERN<sup>6</sup>, C. KAPLAN<sup>1</sup>**

**<sup>1</sup> Platelet Immunology Laboratory, INTS, Paris, France**

**<sup>2</sup> Department of Blood Group Serology and Transfusion Medicine, Division of Blood  
Group Serology, Medical University, Vienne, Austria**

**<sup>3</sup> Centre des Hémophiles, Groupe Hospitalier Necker-Enfant malade, Paris, France**

**<sup>4</sup> Laboratoire d'Hématologie, CHU Robert Debré, Reims, France**

**<sup>5</sup> Hôpital Cardiologique, Centre Hospitalier Régional Universitaire, Lille, France**

**<sup>6</sup> Dynamique des Structures et Interactions des Macromolécules Biologiques, INSERM  
UMR-S 665, Université Paris Diderot - Paris 7, INTS, Paris, France**

**Short title :**

**New mutations in Glanzmann patients and carriers**

**Correspondance**

**Dr Cécile KAPLAN**

**Laboratoire d'Immunologie Plaquettaire**

**INTS**

**6, rue Alexandre Cabanel**

**75015 Paris, France**

**Email : [ckaplan@ints.fr](mailto:ckaplan@ints.fr)**

1  
2  
3 **Phone : 33 1 44 49 30 66 Fax : 33 1 45 67 19 30**  
4

5  
6 **Abstract**  
7

8  
9  
10 Glanzmann thrombasthenia (GT) is an autosomal recessive inherited bleeding disorder  
11 characterized by an impaired platelet aggregation due to defects in integrin  $\alpha$ IIB $\beta$ 3 (ITGA2B,  
12 ITGB3), a fibrinogen receptor. Mutations from 24 GT patients and 2 carriers of various  
13 origins, Caucasian, North-African and Asian were characterized. Promoter and exon  
14 sequences of  $\alpha$ IIB and  $\beta$ 3 genes were amplified and directly sequenced. Among 29 identified  
15 mutations, 17 new allelic variants resulting from nonsense, missense and deletion / insertion  
16 mutations were described. RNA alterations were evaluated by using web-servers. The  $\alpha$ IIB  
17 p.S926L, p.V903F, and  $\beta$ 3 p.C38Y, p.M118R, p.G221D substitutions prevented complex  
18 expression at the surface of Cos-7 cells by altering the  $\alpha$ IIB or the  $\beta$ 3 subunit structure. As  
19 shown by free energy analyses applied on the resolved structure of  $\alpha$ IIB $\beta$ 3 and structural  
20 modeling of the mutant, the p.K253M substitution of  $\beta$ 3 helped to define a key role of the  
21 K253 in the interaction of the  $\alpha$ IIB  $\beta$ -propeller and the  $\beta$ 3  $\beta$ -I domains. Finally, the  $\alpha$ IIB  
22 p.Q595H substitution allowed cell surface expression of the complex but its corresponding  
23 c.2800G>T mutation is predicted to alter normal RNA splicing. Characterization of GT  
24 mutations is of interest for prenatal diagnosis, genetic counseling and provides useful tools for  
25 structure-function relationship studies of  $\alpha$ IIB $\beta$ 3.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

54 **Keywords**

55  $\alpha$ IIB $\beta$ 3, ITGA2B, ITGB3, Glanzmann, protein structure modeling  
56  
57  
58  
59  
60

## INTRODUCTION

Glanzmann thrombasthenia (GT) is an autosomal recessive bleeding disorder characterized by a prolonged bleeding time and an impaired platelet aggregation in response to physiological agonists such as ADP, thrombin, collagen and epinephrine [Caen, 1972]. GT results from quantitative and / or qualitative defects of the fibrinogen (Fg) platelet receptor, the glycoprotein  $\alpha$ IIb $\beta$ 3 complex (OMIM accession number #273800). Classically GT has been divided into 3 subtypes defined by the  $\alpha$ IIb $\beta$ 3 content of patient's platelets : Type I, no or only trace amounts of  $\alpha$ IIb $\beta$ 3; Type II and variant, respectively with low and near normal platelet surface content of functionally impaired  $\alpha$ IIb $\beta$ 3 [Nurden, 2005]. To date, more than 100 different mutations involved in GT phenotype have been described in the  $\alpha$ IIb gene and 60 in the  $\beta$ 3 gene (<http://sinaicentral.mssm.edu/intranet/research/glanzmann/menu>). The nature of GT mutations is highly variable excepting some that are commonly identified in consanguineous populations such as the  $\alpha$ IIb c.409-3\_418del (IVS3-3del13) and the  $\beta$ 3 c.2031\_2041del (2031del11) respectively detected in Arab and Iraqi-Jewish GT populations [Newman et al., 1991], the  $\alpha$ IIb c.1544+1G>A (IVS15+1/G>A) mutation in French Gypsies [Schlegel et al., 1995], or the  $\beta$ 3 c.1053\_1058del (1053CCCAGG) deletion in North-Africans [Morel-Kopp et al., 1997].

Characterization of GT mutations in patients is critical for prenatal diagnosis and genetic counseling. We report here the characterization of 29 mutations identified in 24 thrombasthenic patients and 2 asymptomatic carriers. Using site directed mutagenesis, cell expression models, *in silico* tools and the resolved structure of the  $\alpha$ IIb $\beta$ 3 complex, we have analyzed the pathogenic role of 17 newly described allelic variants. Structural modeling studies of an allelic variant of GT has revealed the importance of the Lys253 in the interaction of the  $\alpha$ IIb  $\beta$ -propeller and the  $\beta$ 3  $\beta$ -I domains.

## MATERIAL AND METHODS

### Patients

Blood samples from patients were sent to our laboratory for confirmation of GT diagnosis and genetic analysis. Propositi and their detailed platelet phenotype are listed in Supp Table S1. Propositi 4 and 15 are GT carriers. Propositus 4 who had presented mild bleeding and healing difficulties during tympanoplasty was addressed to the laboratory for GT investigation after flow cytometry analysis revealed a 50 % reduction of  $\alpha$ IIB $\beta$ 3 platelet surface expression. Propositus 15 was unexpectedly found to be a GT type I carrier during neonatal alloimmune thrombocytopenia diagnosis following in utero fetal death. Her platelets were phenotyped HPA-3a negative but genetically HPA-3a/b as determined by PCR-Sequence Specific Primer testing (data not shown).

As shown in Supp Table SI, 19 propositi of 22 of confirmed platelet phenotype\_ (86 %) were found to be of type I (< 5 % of  $\alpha$ IIB $\beta$ 3 and < 25 % of Fg normal platelet contents). Three propositi were found to be of type II as their platelets presented low but significant  $\alpha$ IIB $\beta$ 3 content ( $\geq$  10 %) and a relatively high level of intra-platelet Fg ( $\geq$  50 %). None of these patients had been previously studied for molecular gene defects. Patients described herein gave their informed consent for genetic analysis.

### Monoclonal antibodies

Monoclonal antibodies (Moab) SZ1 (anti-GPIX), SZ22 (anti- $\alpha$ IIB), SZ21 (anti- $\beta$ 3), P2 (anti- $\alpha$ IIB $\beta$ 3) and E3 (anti-Fg) were purchased from Beckmann Coulter-France (Villepinte, France). Moabs AP2 (anti- $\alpha$ IIB $\beta$ 3) and AP3 (anti- $\beta$ 3) were obtained from Lucron Bioproducts

1  
2  
3 (Lille, France). Moab Y2-51 (anti- $\beta$ 3) was purchased from DAKO ( Trappes, France). Moab  
4  
5 XIIF9 to  $\beta$ 3 was a generous gift from G.Vezon (CRTS, Bordeaux, France). In some flow  
6  
7 cytometry experiments, Moabs SZ22, SZ21 and P2 were directly conjugated to FITC.  
8  
9

### 10 11 12 **Blood samples**

13  
14  
15 Blood samples from patients were collected by venipuncture using 5mM ethylene-diamine-  
16  
17 tetra-acetic acid as an anticoagulant. Platelet were separated by differential centrifugation and  
18  
19 used or stored in 70% ethanol at  $-80^{\circ}\text{C}$  until RNA was extracted. In some cases, 500  $\mu\text{l}$  of  
20  
21 whole blood were mixed with 1.3 ml of RNA Later (Applied Biosystem, Courtaboeuf,  
22  
23 France) and were kept frozen at  $-80^{\circ}\text{C}$  for subsequent RNA extraction.  
24  
25  
26  
27  
28  
29  
30  
31  
32

### 33 34 35 **Flow cytometry**

36  
37 For platelet glycoprotein and Fg content analysis, 50  $\mu\text{l}$  of EDTA anticoagulated whole  
38  
39 blood was diluted in 500  $\mu\text{l}$  of phosphate saline buffer (PBS) pH 7.2 containing 1 % (wt/vol)  
40  
41 BSA-fraction V (Euromedex, Souffelweyersheim, France) and 3 mM EDTA (PBE buffer).  
42  
43 Cell were fixed by adding 1.5 ml of PBS - 3 mM EDTA, pH 7.2, containing 1.33 % (vol/vol)  
44  
45 of a methanol-stabilized formaldehyde solution (Fisher Scientific, Illkirch, France) for 15 min  
46  
47 at room temperature (RT). The reaction was stopped by adding 2 ml of PBE before cells were  
48  
49 pelleted by centrifugation at 1500 g for 10 min. Pelleted cells were washed in 2 ml PBE  
50  
51 before to be resuspended in 1 ml PBE containing 0.1 % (wt/vol) sodium azide and store at  $4^{\circ}\text{C}$   
52  
53 until use. 40  $\mu\text{l}$  of the fixed cells suspension were incubated with 3  $\mu\text{g}/\text{ml}$  Moab in PBE (100  
54  
55  $\mu\text{l}$  final volume), for 1 h at RT. After two washes in 2 ml PBE, the bound Moabs were  
56  
57 detected by using a polyclonal goat anti-mouse IgG conjugated to FITC (Jackson  
58  
59 Immunoresearch Europe, Newmarket, UK) diluted in PBE (1  $\mu\text{l}$  for 10 ml) for 30 min at RT.  
60  
Following one wash in 2 ml PBE, mean fluorescence intensity of cells was analyzed on an

1  
2  
3 XL-MCL Flow cytometer (Beckman Coulter, Villepinte, France). To detect intraplatelet Fg,  
4  
5 each incubation steps were done in presence of 1% (wt/vol) saponine (Fischer Scientific).  
6  
7  
8 Amounts of proteins are expressed as a percentage of the mean amounts obtained with normal  
9  
10 platelets from 10 healthy donors.  
11  
12  
13  
14

15  
16 To analyze  $\alpha$ IIb $\beta$ 3 wild type or mutant complexes expressed at the surface of transfected  
17  
18 Cos-7 cells, cells were collected by Trypsin-EDTA treatment and washed 2 times in PBS-1%  
19  
20 (wt/vol) BSA.  $5 \times 10^5$  cells were incubated in 100  $\mu$ l PBS containing 3  $\mu$ g/ml Moabs (SZ22,  
21  
22 SZ21 or P2) conjugated to FITC for 1 h at RT. Cells were washed twice in 2 ml PBS-BSA  
23  
24 before to be analyzed by flow cytometry. Dead cells were discarded from analysis by using  
25  
26 the 7-AAD viability dye (Beckmann-Coulter). Results are expressed as a percentage of  
27  
28 positive cells for the concerned antigen.  
29  
30  
31  
32

### 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

**DNA/RNA extractions**

Genomic DNA was isolated from peripheral blood according to a published procedure [Jallu et al., 2002] or by using the MagNA Pure Compact Instrument and the Nucleic Acid Isolation Kit I (Roche Diagnostics, Meylan, France). In some cases, DNA samples were obtained by using QIAamp DNA Blood Mini kit (Qiagen, Courtaboeuf, France). RNA extractions were done on platelets by using the Rneasy Protect MiniKit (Qiagen). DNAs and RNAs were extracted according to the manufacturers' instructions.

**Sequencing analysis**

Sequencing was performed as previously described [Quintanar et al., 1998] or by GenoScreen (Lille, France). Primers defined by Kato and coworkers [Kato et al., 1992] and Jin and co-workers [Jin et al., 1993] were respectively used to amplify coding sequences of

1  
2  
3  $\alpha$ IIb and  $\beta$ 3 genes from genomic DNA. Primers to amplify promoter sequences of  $\alpha$ IIb and  
4  
5  $\beta$ 3 have been respectively defined according to the genes' reference sequences NG\_008331.1  
6  
7 and NG\_008332.  $\alpha$ IIb promoter sequences were studied by using the following primer pairs  
8  
9 ("a" forward, "b" reverse) :

10  
11  
12 IIbP1a CGCCAGAATTCTCAGTCACGAAGCTTGGCTCAAGACGGAG

13  
14 IIbP1b CCAGCCAGATATCAGCTATGTACTACCACCGTGCTAGTCC

15  
16 IIbP2a CGCCAGAATTCTCAGTCACGGGTAAAGATTCAAGAGACAT

17  
18 IIbP2b CCAGCCAGATATCAGCTATGTCAGATTCCTCCACAGGAAG

19  
20 IIbP3a CGCCAGAATTCTCAGTCACGGAAGGGAAGGAGGAGGAGCT

21  
22 IIbP3b CCAGCCAGATATCAGCTATGCTTCCCTTACGGCTCACCTC

23  
24  
25 Underlined sequences are non genomic sequence for sequencing PCR products. Promoter  
26  
27 and exon 1 sequences of  $\beta$ 3 gene were amplified with the primer pair

28  
29 IIIaE1a ATGTGGTCTTGCCCTCAACAGGTAG

30  
31 IIIaE1b CACGCTCTCACCCAGGAAGTTACAG by using the Advantage GC genomic  
32  
33 PCR kit (Clontech, Saint-Germain-en-Laye, France) and then sequencing was done with the  
34  
35 forward primer 3aseq-P1a CCGGAAAACCAAATAAGGC and the reverse primer 3aseq-  
36  
37 P1b GACCAGTCCTCCGCGTTTGC.

38  
39 All identified mutations were confirmed by PCR-Restriction Fragment Length  
40  
41 Polymorphism (PCR-RFLP) analysis on a second PCR product. Restriction enzymes are  
42  
43 listed in tables 1 and 2 with the corresponding mutations. The experimental details of these  
44  
45 RFLP assays can be obtained upon request. When the mutation did not introduce restriction  
46  
47 modification, sequencing was done on a second PCR-product. For nucleotide numbering the  
48  
49 A nucleotide of the ATG start was designated +1 (cDNA ITGA2B and ITGB3 GenBank  
50  
51 accession numbers NM\_000419.3 and NM\_000212.2 respectively). For amino acid  
52  
53 numbering used in the text, the first amino acids of the mature  $\alpha$ IIb and  $\beta$ 3 proteins were  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 designated +1. Residue numbering according to the Human Genome Variation Society  
4 (HGVS) nomenclature that designed as +1 the translation initiating methionine  
5 (http://www.hgvs.org/mutnomen) is also shown in tables 1 and 2. Both numberings differ by  
6  
7  
8  
9  
10 31 and 26 residues that correspond to the signaling peptides of  $\alpha$ IIB and  $\beta$ 3 respectively. All  
11  
12 identified allelic variations have been deposited on the NCBI dbSNP database  
13 (http://www.ncbi.nlm.nih.gov/SNP/) and their corresponding accession numbers are indicated  
14  
15  
16  
17  
18 in tables 1 and 2.

### 21 22 **Site directed mutagenesis and in-vitro expression**

23  
24 Site directed mutagenesis and *in vitro* expression in Cos-7 cells were done as previously  
25 described [Jallu et al., 2002] or by using the QuikChange® II XL Site-Directed Mutagenesis  
26 Kit (Agilent Technologies, Massy, France) according to the manufacturer's instructions. In  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
some mutagenesis experiments, plasmid pcDNA3.1-Zeo containing the  $\alpha$ IIB cDNA was used  
instead of pcDNA3.1-Neo (Invitrogen, Cergy Pontoise, France).

### 41 42 **Minigene study**

43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
To study a mutation predicted to affect mRNA splicing, an  $\alpha$ IIB gene fragment covering  
exons 17 to 20 and containing the mutation was synthesized and introduced in the plasmid  
Exontrap (Mobitech, Goettingen, Germany). The wild type form of the sequence was obtained  
by site directed mutagenesis. Synthesis, cloning and site directed mutagenesis were done by  
the Epoch Biolabs, Inc. Company (Sugar Land, TX, USA). Following transient transfection in  
Cos cells, total RNA was extracted and mRNA amplified for the exon 17 to 20 and directly  
sequenced.

### Western-blot analysis

Trypsin-EDTA collected Cos-7 cells were washed 2 times in PBS and 1 time in 50 mM Tris-HCL pH 7.6, 150 mM NaCl. Cells were resuspended ( $1.5 \times 10^7$  cells) in 100  $\mu$ l of the same Tris buffer containing 1% (wt/vol) Triton-X 100 and 2 mM para-Methyl Sulfonyl Fluoride (PMSF). They were lysed for 30 min at 4°C and the lysate centrifuged at 16.000 g for 30 min at 4°C. The supernatant was diluted with 4 volumes of a 150 mM NaCl solution containing 2.5 % (wt/vol) Sulfate Dodecyl Sodium (SDS), 31.25 mM n-ethylmaleimide and 2.5 mM PMSF. The SDS lysate was heated to 100°C for 5 min before 35  $\mu$ l were used for western-blot analysis as previously described [Jallu et al., 1994] with a minor modification : bound antibodies were detected by using anti-mouse IgG conjugated to peroxidase associated with a chemiluminescence detection technique.

### Computational analyses

The effect of mutations on intron splicing has been predicted *in silico* by using the following tools : Genscan (<http://genes.mit.edu/GENSCAN.html> [Burge and Karlin, 1998]) and NNSPLICE 0.9 ([http://www.fruitfly.org/seq\\_tools/splice.html](http://www.fruitfly.org/seq_tools/splice.html) [Reese et al., 1997]). Modifications of the splicing factors binding sites pattern resulting from the mutations were evaluated by using the ESEFinder tool [Cartegni et al., 2003; Smith et al., 2006].

To compare wild type and a mutant headpiece of  $\alpha$ IIB $\beta$ 3, modeling has been done using the structure (PDB code 2VDL) published by Springer et co-workers [Springer et al., 2008]. Residues 1-452 of  $\alpha$ IIB and 109-352 of  $\beta$ 3 were taken into account. The interface between the protein chains has a correct resolution (2.75 Å) and the residues implicated at the interface have all good normalized B-factor values. A structural model of the mutant has been built. This model was checked using ProCheck software to analyze the quality of the modification [Laskowski et al., 1993] and the side-chain position was analyzed with SCWRL [Canutescu et

1  
2  
3 al., 2003]. Finally, the FastContact web server (<http://structure.pitt.edu/servers/fastcontact/>)  
4  
5 was used to estimate the direct electrostatic and desolvation interaction free energy between  
6  
7 two proteins [Camacho and Zhang, 2005; Champ and Camacho, 2007]. The computational  
8  
9 estimation of the binding free energy is based on a statistically determined desolvation contact  
10  
11 potential and Coulomb electrostatics with a distance-dependent dielectric constant [Camacho  
12  
13 and Zhang, 2005]. This free energy provides a reasonably good estimate of experimental  
14  
15 binding affinities from complex crystal structures.  
16  
17  
18  
19  
20  
21

## 22 RESULTS

23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Clinical GT diagnosis was confirmed in our laboratory by immunological or biochemical studies (see Supp Table S1). Genomic DNA samples from patients were studied for molecular defects in promoter and exon sequences of genes coding for  $\alpha$ IIB and  $\beta$ 3 by direct sequencing of the corresponding PCR products. Allelic variations identified in these genes are reported in tables 1 and 2. To avoid erroneous identification due to amplification errors, all sequence alterations have been confirmed on a second PCR product either by a PCR-RFLP technique when a restriction site modification was observed or by new sequencing analysis in other cases. Restriction enzymes that have been used are listed in tables 1 and 2 with their corresponding allelic variant. When available, DNAs from patient's relatives were also analyzed and used to confirm the allelic variations (data not shown).

### Sequence alterations detected in the $\alpha$ IIB gene

For 18 propositi, a total of 20 mutations was identified in the gene coding for  $\alpha$ IIB (see table 1). Five lead to nonsense codons : patients 1, 6, 7, 8 and 17. It should be noted that patient 17 was phenotype GT type II because the second mutation that had not been identified allowed residual expression of  $\alpha$ IIB $\beta$ 3 (see Supp Table S1). Three mutations directly affect a splice site : patients 9 and 12 with a single point mutation and 3 unrelated propositi (GT

1  
2  
3 patients 3 and 5, and the carrier 4) who present the same 13 base pair deletion c.1442-  
4 13\_1442-1del. However for these latter propositi, an unknown common ancestry cannot be  
5  
6 excluded as all are European Caucasians who live in the north-east of France. Two  
7  
8 heterozygous single base insertions were identified in patient 2. These mutations are predicted  
9  
10 to result in frame-shift anomalies creating a downstream stop codon and premature  
11  
12 termination. A 10 base pair deletion identified in patient 14 also leads to a frame shift  
13  
14 anomaly. Other gene anomalies identified affecting  $\alpha$ IIb, were missense mutations. Three of  
15  
16 them (patients 6, 10, 15 and 16) have not been described so far and 6 (patients 3, 11, 12, 13  
17  
18 and 18) have already been reported (see references in Table 1). Among the 9 heterozygous  
19  
20 compounds, 4 second obligatory mutations (patients 5, 7, 17 and 18) remain to be identified  
21  
22 (NI abbreviation (not identified) in tables).  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

### 33 **Sequence alterations detected in the $\beta$ 3 gene**

34 Ten mutations were identified in the gene coding for the  $\beta$ 3 glycoprotein for 8 propositi  
35  
36 (see Table 2). One single point mutation (patient 20) is responsible for a nonsense codon and  
37  
38 a second (patient 19) disrupts an acceptor splice site. An homozygous 4 base pair deletion  
39  
40 located downstream of the donor splice site (c.1125+3\_1125+6del) of intron 8 has been  
41  
42 identified as the unique mutation in patient 25. In patient 26, two silent homozygous  
43  
44 mutations, c.1143C>A and c.1260G>A (20507C>A and 20624G>A with genomic  
45  
46 numbering : accession number J05427) located in exon 9 are observed. These mutations have  
47  
48 been reported to be responsible for type I GT [Jin et al., 1996]. Though silent, their  
49  
50 association induces exon 9 skipping and the introduction of five intronic bases in the final  
51  
52 transcript [Jin et al., 1996]. Analysis of the mRNA from patient 26 has confirmed this deletion  
53  
54 / insertion mechanism (data not shown). Finally five single point missense mutations were  
55  
56  
57  
58  
59  
60

1  
2  
3 identified (patients 21, 22, 23 and 24). In two heterozygous patients the second obligatory GT  
4  
5 mutation remains to be determined.  
6  
7  
8  
9

### 10 **Flow cytometry analysis of mutant complexes expressed in COS-7 cells**

11  
12 Seven missense mutations not previously reported in the literature have been further  
13 studied by expressing the mutant complexes in COS-7 cells to assess their role in the GT  
14 phenotype. Cell surface expression of the mutant complexes was assessed by flow cytometry  
15 using Moabs SZ22 to  $\alpha$ IIB, SZ 21 to  $\beta$ 3 or P2 to the  $\alpha$ IIB $\beta$ 3 complex (see Figure 1).  $\alpha$ IIB  
16 p.S926L (patient 16) and p.V903F (patients 6 and 15) and  $\beta$ 3 p.C38Y (patient 21), p.M118R  
17 (patient 22), p.G221D and p.K253M (patient 24) substitutions prevented normal complex  
18 expression, mimicking the type I GT phenotype observed for the patients. For the  $\alpha$ IIB  
19 p.S926L (patient 16) and  $\beta$ 3 p.G221D and p.K253M (patient 24) mutants the percentage of  
20 positive cells expressing  $\beta$ 3 was slightly higher than in other mutants tested (see Figure 1).  
21 This low level expression could be correlated to the 10 % of  $\beta$ 3 expression at the surface of  
22 the patients' platelets (see Supp Table S1).  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Surprisingly, the  $\alpha$ IIB p.Q595H substitution identified in a GT type I patient did not affect  
the expression of the mutant complex which is similar to that obtained with the wild type  
form.

As control, untransfected cells did not significantly bind any of the Moabs tested and  
irrelevant mouse IgG gave less than 1 % of positive cells whatever the transfectant tested.

### **In silico analyses of mRNA processing for missense allelic variations and for the intronic deletion c.1125+3\_1125+6del**

Some exonic single point mutations can alter normal processing of the mRNA and lead to  
the synthesis of an aberrant mRNA, associated or not with its degradation. Genscan,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

NNSPLICE 0.9 and ESEFinder web servers were used to predict the effect of the missense mutations responsible for the substitutions that have been tested in flow cytometry. From this panel, the c.1878G>C responsible for the p.Q595H substitution was the unique variation predicted to modify normal splicing and to affect ESE binding sites. Splice site prediction by NNSPLICE 0.9 cannot identify the normal donor splice site of intron 18 in the mutant. Genscan analysis did not predict splice anomalies, but scores were lowered for both exons 18 (0.987 to 0.791) and 19 (0.995 to 0.849) implying a probability of correct splicing of 75 % instead of 99 % for the normal sequence. Furthermore ESEFinder shows mutation 1878G>C to increase the score of an SF2/ASF binding site from 2.67 to 3.28 and to create a new one with a score of 2.50 (threshold 1.95). Finally an SRp55 binding site that presents a high score (4.33 - threshold 2.67) encompassing the normal splice site is also created.

In the  $\beta 3$  gene, the homozygous 4 bp deletion c.1125+3\_1125+6del (patient 25) was the single intronic mutation identified which does not directly affect a splice site. However Genscan analysis predicts the use of a cryptic donor splice site located 70 bp downstream the normal one. If this transcript is not degraded, this would lead to an in frame introduction of 23 amino-acids between  $\beta 3$   $\beta$ -I and hybrid domains. ESEFinder shows no modifications but NNSPLICE 0.9 suggests the contiguous donor splice site would be ignored in the mutant.

#### **Western-blot analysis of the p.C38Y $\beta 3$ mutation effect on the complex expression.**

All the  $\beta 3$  mutants tested strongly affect complex expression (Figure 1) but the p.C38Y substitution also strongly and specifically affects that of  $\beta 3$  when tested with anti- $\beta 3$  Moab SZ21. To understand this specific effect, synthesis of the Y38 $\beta 3$  form of the complex in the transfected cells was analyzed by using a Western-blot technique with Moabs SZ22 to  $\alpha$ Ib and SZ21, XIIF9 and AP3 to  $\beta 3$  (see Figure 2). SZ22 showed that  $\alpha$ Ib is clearly detectable in the lysate of the wild type  $\alpha$ Ib $\beta 3$  transfectant (lane c) as in the control platelet lysate (lane a).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For the  $\alpha$ IIb-Y38 $\beta$ 3 mutant transfectant, only a band of slightly greater relative mobility which might correspond to a degradation product of  $\alpha$ IIb was detected (lane d). Moab XIIF9 revealed the presence of the Y38 $\beta$ 3 mutant in the  $\alpha$ IIb-Y38 $\beta$ 3 transfectant (lane d). However, this mutant glycoprotein presents a lower relative mobility when compared to the wild type forms detected in lysates from platelets or wild type  $\alpha$ IIb $\beta$ 3 transfectant (lane c). Both SZ21 and AP3 failed to react with the Y38 $\beta$ 3 mutant (lanes d) but reacted with normal  $\beta$ 3 expressed in platelet and wild type  $\alpha$ IIb $\beta$ 3 transfectant (lanes a and c respectively). XIIF9 and AP3 but not SZ21 reacted with the endogenous  $\beta$ 3 subunit of the vitronectin receptor expressed in Cos-7 cells as shown by the faint band detected in non transfectant cells (lane b). These results suggested that the p.C38Y mutation should impair the complex formation, or at least its normal processing, as no  $\alpha$ IIb of normal mobility could be detected.

### **Role of the Lys253 of the $\beta$ -I domain of $\beta$ 3 in the interaction with the $\alpha$ IIb $\beta$ -propeller**

To understand precisely the impact of the p.K253M mutation, we have analyzed the interface between  $\alpha$ IIb (residues 1-452) and  $\beta$ 3 (residues 109-352) subunits using the protein structures obtained by the group of Springer (PDB code 2VDL [Springer et al., 2008]). From a simple visualization, the side chain of the  $\beta$ 3 K253 protrudes from the  $\beta$ -I domain toward the  $\alpha$ IIb beta-propeller surface and makes contact with different residues of the  $\alpha$ IIb subunit (see Figure 3A). Table 3 summarizes the different results concerning the estimations of the direct electrostatic and desolvation interaction free energy between the  $\alpha$ IIb and  $\beta$ 3 chains. Interestingly, K253 has a key role in terms of binding free energy with the most negative contribution (-5.5 kcal/mol) and also in terms of electrostatics (-9.7 kcal/mol). Inversely, desolvation free energy is high (+4.1 kcal/mol). The desolvation of charged and polar amino

1  
2  
3 acid is unfavorable from an energy point of view. This penalty of desolvation is greatly  
4 compensated by Coulomb interactions and by the hydrogen bonds which are formed in the  
5  
6 complex.  
7  
8

9  
10 Analysis of important interactions of by K253 underlines its weight in the interaction as it  
11 has the 1<sup>st</sup> (-6.7 kcal/mol with D232 of  $\alpha$ IIb), the 6<sup>th</sup> (-2.0 kcal/mol with F231) and the 13<sup>th</sup> (-  
12 0.8 kcal/mol with P238) most negative interactions of the interface. One interaction with  
13  
14 W262 is slightly unfavorable. Concerning free energy, the same observations are made, with  
15  
16 two strong interactions with D232 (-5.5 kcal/mol) and F231 (-1.2 kcal/mol). One unfavorable  
17  
18 interaction is observed again with W262. These results highlight the significance of this  
19  
20 residue, the most important in terms of electrostatic and free energy computation.  
21  
22

23  
24 A structural model of M253 mutant was constructed. Regarding the whole  $\alpha$ IIb and  $\beta$ 3  
25  
26 interface, the substitution provokes a striking loss of electrostatic energy (+9.1 kcal/mol) but a  
27  
28 gain in desolvation free energy (-5.72 kcal/mol) that cannot counter-balance the decrease of  
29  
30 electrostatic energy. As shown in Table 3, the diminution of the length of the side-chains and  
31  
32 the change from a charged polar amino acid to a hydrophobic amino acid has a deep impact.  
33  
34 Residue 253 has no more (negative) interaction with either D232 or F231, or P238. Only a  
35  
36 single interaction with W262 is conserved (see Figure 3B).  
37  
38

39  
40 From a visual point of view, the decrease of electrostatic energy between K253 (see  
41  
42 Figure 3C) and M253 (see Figure 3D) is striking. Overall, the most important electrostatic  
43  
44 energy implicated in the interaction becomes negligible. All these contacts are lost in presence  
45  
46 of the methionine, except for W262 that only preserves an electrostatic link. These results  
47  
48 indicate that it is rather the loss of K253 lost than the presence of the methionine which is  
49  
50 responsible for the defect of complex expression.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## DISCUSSION



1  
2  
3 The aim of the study was to identify mutations responsible for GT platelet phenotype.  
4  
5 SDS-PAGE, western-blot and flow cytometry analyses carried out on patients' platelets  
6  
7 confirmed GT clinical diagnosis. Twenty-nine mutations were identified in both  $\alpha$ IIb and  $\beta$ 3  
8  
9 genes. Most of these mutations (47 %) correspond to missense mutations but nonsense,  
10  
11 insertion, deletion, and single point mutations affecting splice sites were also found.  
12  
13

14  
15 Mutations previously reported in the literature (see tables 1 and 2) have not been further  
16  
17 studied in-vitro for their involvement in GT. It can be noted that their presence in the patients  
18  
19 studied herein and in one or more additional unrelated patients would confirm their role in GT  
20  
21 phenotype. New mutations directly affecting splice sites, creating nonsense codon or insertion  
22  
23 / deletion associated with frame shift anomaly were considered as known cause of GT. The  
24  
25 pathological effects of new allelic variations responsible for missense substitutions and one  
26  
27 new allelic variation leading to a splicing anomaly are discussed below.  
28  
29

30  
31 The substitutions p.V903F (patient 6 and propositus 15) and p.S926L (patient 16) affect  
32  
33 the extra-cellular N-terminal part of the  $\alpha$ IIb  $\beta$ -subunit that composes the calf-2 domain (see  
34  
35 Figure 4). In both cases they prevent  $\alpha$ IIb $\beta$ 3 complex expression at the Cos-7 cell surface.  
36  
37 They are predicted to impair the correct folding of the Calf-2 domain as the amino-acids V903  
38  
39 and S926 are located in its core (see Figure 4). To estimate their structural importance, the  
40  
41 human  $\alpha$ IIb primary sequence has been aligned with several corresponding integrin sequences  
42  
43 - marsupial, fish, amphibian, prochordates and insect (Supp Figure S1A). These sequences  
44  
45 share at least 45 % of identity with the human  $\alpha$ IIb (Psi-blast data from  
46  
47 <http://www.ncbi.nlm.nih.gov/BLAST>) and only 17 % of these residues are conserved through  
48  
49 all the seven organisms tested. It is noteworthy that V903 and S926 belong to these few  
50  
51 amino-acids, indicating that their conservation could be linked to a strong structural  
52  
53 evolutionary constraint. Furthermore, their poor conservation in other human integrin  $\alpha$ -  
54  
55 subunits (data not shown) reinforces their  $\alpha$ -IIb specificity.  
56  
57  
58  
59  
60

1  
2  
3 The missense mutation c.1878G>C (patient 10), contiguous with the donor splice site of  
4 intron 18, was predicted to be responsible for a p.Q595H substitution in  $\alpha$ IIB. However this  
5  
6 p.Q595H mutation identified in a type I GT patient does not prevent the expression of the  
7  
8 mutant complex at the surface of transfected Cos-7 cells. Study of the patient's platelet  
9  
10 mRNA revealed full normal  $\beta$ 3 mRNA synthesis but  $\alpha$ IIB mRNA remained undetectable. A  
11  
12 splicing anomaly can lead to mRNA decay and explain the lack of detection of the mRNA  
13  
14 coding for  $\alpha$ IIB in patient 10. *In silico* splicing testing by using the Genscan and\_NNSPLICE  
15  
16 0.9 web servers effectively predicted splicing anomalies. Similarly Jayo and coworkers [Jayo  
17  
18 et al., 2006] reported that the  $\alpha$ IIB c.2829C>T mutation (exon 27) predicted to induce a  
19  
20 p.L912P substitution, partially leads to exon 28 skipping. The authors pointed out that rather  
21  
22 than the disruption of an ESE site, the aberrant mRNA processing would result from the  
23  
24 strengthening of the SR protein SC35 binding score by 0.65 units (ESEFinder analysis).  
25  
26 Interestingly the ESEFinder web server predicted the mutation c.1878G>C to increase the  
27  
28 score by 0.61 of an existing SF2/ASF binding site and to create two new binding sites for  
29  
30 SF2/ASF and SRp55 factors with significant high scores (respectively 2.5 and 4.33). It can  
31  
32 therefore be hypothesized that the 1878G>C mutation alters the normal splice factor pattern  
33  
34 necessary to accurately splice intron 18. This was confirmed by a minigene study that showed  
35  
36 the c.1878G>C alteration to provoke exon 18 skipping during mRNA splicing (data not  
37  
38 shown).  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 In the  $\beta$ 3 gene, the homozygous 4 bp deletion c.1125+3\_1125+6del (patient 25) was the  
50  
51 single intronic mutation identified that does not directly affect a splice site but its contiguous  
52  
53 intronic sequence. Genscan analysis predict an altered splicing that correlates with the partial  
54  
55 loss of a sequence necessary to hybridize a small nuclear ribonucleoprotein involved in the  
56  
57 splicing process [Wang and Cooper, 2007].  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The mutation c.191G>A (patient 21) responsible for a p.C38Y substitution, is located in the PSI domain of the  $\beta 3$ . Transient co-transfection of the  $\beta 3$  mutated cDNA and with normal  $\alpha$ IIb cDNA showed the mutation to reproduce a GT type I phenotype as no complex was detected on the cell surface by any of the anti- $\alpha$ IIb $\beta 3$  Moabs tested herein and including testing with a panel to  $\beta 3$  (XIIF9, AP3 and Y2-51, data not shown). Computational analysis (Genscan, ESEFinder and NNSPLICE 0.9) indicated that the c.191G>A mutation would not induce splicing modification of the pre-mRNA. However, western-blot analyses showed that  $\alpha$ IIb and the mutated  $\beta 3$  were synthesized in transfected COS cells. The mutated Y38  $\beta 3$  presents a lower relative mobility than the wild type form which could result from structural modifications induced by the C16 – C38 disulfide bond disruption. A similar observation has been reported by Valentin and co-workers [Valentin et al., 1995], who have shown that alanine substitution of C38 lowered the relative mobility of  $\beta 3$  expressed in COS cells. The structural modification of the PSI domain induced by the p.C38Y mutation was evidenced by the lack of reactivity of Moabs AP3 and SZ21 whose epitopes are respectively located in amino-acids stretches 49-98 [Peterson et al., 2003] and 28-35 [Honda et al., 1995]. So the structural anomaly of the PSI domain impairs the complex association and / or its post-translational maturation leading to its retention inside the cell.

The missense mutations p.M118R, p.G221D and p.K253M are located in the  $\beta$ -I domain of  $\beta 3$  and are responsible for Type I GT. M118 and G221 are highly conserved in most integrin  $\beta$  subunits (data not Shown), and through invertebrates to vertebrates (see Supp Figure S1B). Most of the  $\beta$ -I domain mutations previously described, are responsible for Variant [Bajt et al., 1992; D'Andrea et al., 2002; D'Andrea et al., 2008; Lanza et al., 1992; Loftus et al., 1990] or type II GT [Jackson et al., 1998; Morel-Kopp et al., 2002; Nair et al., 2004; Tadokoro et al., 2002; Ward et al., 2000]. For comparison, Figure 5 locates the mutated amino-acids in the primary sequence (panel A) and in the  $\beta$ -I domain structure (panel B). Whatever the variant or

1  
2  
3 type II GT phenotype, the mutated complexes are located completely or partially at the  
4 surface of the  $\beta$ -I domain, and some of them (see Figure 5B, underlined amino-acids) present  
5  
6 significant electrostatic or / and free energies interactions with the  $\beta$ -propeller (data not  
7  
8 shown). Variant mutations concern the MIDAS or SyMBS sites [Zhu et al., 2008] directly  
9  
10 affecting amino-acids involved in metal coordination (p.D119Y, p.S123P) or their very  
11  
12 nearby environment (p.R217W or Q) (see Figure 5A). Both variant and GT Type II mutations  
13  
14 support at least a 10 % expression of  $\alpha$ I**II** $\beta$ 3 at the platelet surface. This contrasts with the  
15  
16 failure to detect the  $\beta$ 3-R118 and -D221 forms of the complex. Both the p.M118R and the  
17  
18 p.G221D mutations, located inside the  $\beta$ -I domain, should deeply alter its proper folding,  
19  
20 preventing complex expression at the platelet surface. Due to the nature of the amino-acid  
21  
22 involved, these mutations induce important changes of steric hindrance and charge. However,  
23  
24 in one case, the mutation p.L117W which affects a 100 % buried amino-acid still permits a 10  
25  
26 % expression of  $\alpha$ I**II** $\beta$ 3 at the platelet surface [Basani et al., 1997]. We could hypothesize that  
27  
28 the characteristics of these residues, both hydrophobic and aliphatic, could explain this  
29  
30 behavior. In contrast the p.G221D mutation, changes of volume and charge of the side-chains  
31  
32 has a too deep impact. Similarly, such a steric hindrance change could explain the type I  
33  
34 phenotype induced by the p.M124V mutation [Gonzalez-Manchon et al., 2004], as the M124  
35  
36 side chain pointed inward the  $\beta$ -I domain.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

47 Unlike for M118 and G221, K253 is not conserved in other integrin  $\beta$  subunits and if it is  
48  
49 present in  $\beta$ 3 from fish to primates it is replaced by an arginine in amphibian and absent from  
50  
51 invertebrate integrins (see Supp Figure S1B). However the p.K253M substitution results in  
52  
53 type I GT. The 3-dimensional structure of the  $\alpha$ I**II** $\beta$ 3 headpiece revealed that the side chain of  
54  
55 K253 protrudes from the  $\beta$ -I domain toward a pocket at the surface of the  $\alpha$ I**II** $\beta$  propeller  
56  
57 (see Figure 3A and 5B). Analyses of electrostatic, desolvation and free energies revealed that  
58  
59 K253 is one of the most important amino-acids involved in the  $\beta$ -I domain interaction with  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

the  $\beta$ -propeller of  $\alpha$ IIb. Furthermore, a mutant model of the  $\alpha$ IIb $\beta$ 3 headpiece shows that the p.K253M substitution does not introduce steric and energy clashes but only suppresses important contacts with the  $\alpha$ IIb  $\beta$ -propeller. So, the p.K253M mutation revealed K253 as a key amino-acid for the association of the  $\beta$ -I domain with the  $\beta$ -propeller. The last mutation reported so far of the  $\beta$ -I domain that leads to type I GT was a substitution of the I325-P326-G327 sequence by a methionine. This mutation impairs the  $\alpha$ IIb $\beta$ 3 association [Morel-Kopp et al., 1997]. However, the mutated amino-acids locate between the  $\alpha$ 6 helix and the  $\beta$ 7 sheet and do not directly face the  $\alpha$ IIb  $\beta$ -propeller surface (see Figure 5B). Furthermore the I325-P326-G327 sequence does not seem to have significant if any, interaction with the  $\beta$ -propeller as indicated by free and electrostatic energy analysis (data not shown). It might indirectly impair the interaction(s) of the  $\alpha$ 5 and / or  $\alpha$ 6 helix with the  $\beta$ -propeller. The last four mutations described, p.K137Q [Peterson et al., 2008], p.T140I (HPA-16bw) [Jallu et al., 2002], p.R143Q (HPA-4) [Wang et al., 1992] and p.T195N (HPA-17bw) [Stafford et al., 2008] only lead to allelic variants involved in alloimmune thrombocytopenia that are not related to GT phenotype. All these amino-acids locate at the surface of the  $\beta$ -I domain with their side chains directed outside.

In conclusion, we have identified 29 mutations in a panel of 24 GT patients and 2 GT carriers. Seventeen of these mutations have not been described so far. We have confirmed by *in vitro* studies that the missense mutations identified impaired normal expression of the mutant complex (except for one case). Nonsense and splice site mutations were considered to be related to the GT phenotype. This study emphasizes the high allelic heterogeneity of both the  $\alpha$ IIb and  $\beta$ 3 genes involved in GT phenotype. Finally the GT variant p.K253M of  $\beta$ 3 unveiled the key role of K253 in the  $\beta$ -propeller /  $\beta$ -I association of the  $\alpha$ IIb $\beta$ 3 headpiece.

## Acknowledgments

We thank the following colleagues for sending us blood samples from the patients we have studied here-in : B. Bastenaire, C. Caron, M. Dreyfus, F. Dutrillaux, I Elalami, Nizard, G. Le Roux, J. Peynet, S. C. Saladin-Thiron / MT Bricquet, P. Tron, J.P. Vernant. We also thank the following technician students for their help in mutation identification : M. Brément, S. Cantais, E. Divron, F. Jacquinot, A. Ledigarcher, P. Moffat, V. Noel and F. Ping. The work done by M. Dusseaux was supported by a grant from Novo Nordisk. Conflict Interests: no conflict financial interests exist.

## Reference List

- 1  
2  
3  
4  
5  
6 Bajt ML, Ginsberg MH, Frelinger III AL, Berndt MC, Loftus JC. 1992. A spontaneous  
7 mutation of integrin  $\alpha$ IIB $\beta$ 3 (platelet glycoprotein IIB-IIIa) helps define a ligand  
8 binding site. *J Biol Chem* 267:3789-3794.  
9
- 10 Basani RB, Brown DL, Vilaire G, Bennett JS, Poncz M. 1997. A Leu<sup>117</sup>--> Trp mutation  
11 within the RGD-peptide cross-linking region of  $\beta$ 3 results in Glanzmann  
12 thrombasthenia by preventing  $\alpha$ IIB $\beta$ 3 export to the platelet surface. *Blood* 90:3082-  
13 3088.  
14
- 15 Basani RB, French DL, Vilaire G, Brown DL, Chen F, Collier BS, Derrick JM, Gartner TK,  
16 Bennett JS, Poncz M. 2000. A naturally occurring mutation near the amino terminus  
17 of  $\alpha$ IIB defines a new region involved in ligand binding to  $\alpha$ IIB $\beta$ 3. *Blood* 95:180-188.  
18
- 19 Burge CB, Karlin S. 1998. Finding the genes in genomic DNA. *Curr Opin Struct Biol* 8:346-  
20 354.  
21
- 22 Caen JP. 1972. Glanzmann thrombasthenia. *J Clin Invest* 1:388-391.  
23
- 24 Caen JP. 1989. Glanzmann's thrombasthenia. *Baillieres Clin Haematol* 2:609-625.  
25
- 26 Camacho CJ, Zhang C. 2005. FastContact: rapid estimate of contact and binding free  
27 energies. *Bioinformatics* 21:2534-2536.  
28
- 29 Canutescu AA, Shelenkov AA, Dunbrack RL, Jr. 2003. A graph-theory algorithm for rapid  
30 protein side-chain prediction. *Protein Sci* 12:2001-2014.  
31
- 32 Cartegni L, Wang J, Zhu Z, Zhang MQ, Krainer AR. 2003. ESEfinder: A web resource to  
33 identify exonic splicing enhancers. *Nucleic Acids Res* 31:3568-3571.  
34
- 35 Champ PC, Camacho CJ. 2007. FastContact: a free energy scoring tool for protein-protein  
36 complex structures. *Nucleic Acids Res* 35:W556-W560.  
37
- 38 D'Andrea G, Bafunno V, Del VL, Amoriello A, Morabito P, Vecchione G, Grandone E,  
39 Margaglione M. 2008. A beta3 Asp217-->Val substitution in a patient with variant  
40 Glanzmann Thrombasthenia severely affects integrin  $\alpha$ IIB $\beta$ 3 functions. *Blood*  
41 *Coagul Fibrinolysis* 19:657-662.  
42
- 43 D'Andrea G, Colaizzo D, Vecchione G, Grandone E, Di Minno G, Margaglione M, on behalf  
44 of the Glanzmann's thrombasthenia Italian team (GLATIT). 2002. Glanzmann's  
45 thrombasthenia : Identification of 19 new mutations in 30 patients. *Thromb Haemost*  
46 87:1034-1042.  
47
- 48 French DL, Collier B. 1997. Hematologically important mutations: Glanzmann  
49 thrombasthenia. *Blood Cells, Molecules, and Diseases* 23:39-51.  
50
- 51 Gonzalez-Manchon C, Butta N, Larrucea S, Arias-Salgado EG, Alonso S, Lopez A, Parrilla  
52 R. 2004. A variant thrombasthenic phenotype associated with compound  
53 heterozygosity of integrin  $\beta$ 3-subunit : (Met124Val) $\beta$ 3 alters the subunit dimerization  
54 rendering a decreased number of constitutive active  $\alpha$ IIB $\beta$ 3 receptors. *Thromb*  
55 *Haemost* 92:1377-1386.  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Gonzalez-Manchon C, Fernandez-Pinel M, Arias-Salgado EG, Ferrer M, Alvarez MV, Garcia-Munoz S, Ayuso MS, Parrilla R. 1999. Molecular genetic analysis of a compound heterozygote for the glycoprotein (GP) IIb gene associated with Glanzmann's thrombasthenia : disruption of the 674-687 disulfide bridge in GPIIb prevents surface exposure of GPIIb-IIIa complexes. *Blood* 93:866-875.
- Honda S, Honda Y, Bauer B, Ruan C, Kunicki TJ. 1995. The impact of three-dimensional structure on the expression of PI<sup>A</sup> alloantigens on human integrin  $\beta_3$ . *Blood* 86:234-242.
- Jackson DE, White MM, Jennings LK, Newman PJ. 1998. A ser<sub>162</sub>--> Leu mutation within glycoprotein (GP) IIIa (integrin  $\beta_3$ ) results in an unstable  $\alpha_{IIb}\beta_3$  complex that retains partial function in a novel form of type II Glanzmann thrombasthenia. *Thromb Haemost* 80:42-48.
- Jallu V, Diaz-Ricart M, Ordinas A, Pico M, Vezon G, Nurden AT. 1994. Two human antibodies reacting with different epitopes on integrin  $\beta_3$  of platelets and endothelial cells. *Eur J Biochem* 222:743-751.
- Jallu V, Meunier M, Brement M, Kaplan C. 2002. A new platelet polymorphism Duv(a+), localized within the RGD binding domain of glycoprotein IIIa, is associated with neonatal thrombocytopenia. *Blood* 99:4449-4456.
- Jayo A, Pabon D, Lastres P, Jimenez-Yuste V, Gonzalez-Manchon C. 2006. Type II Glanzmann thrombasthenia in a compound heterozygote for the alphaIIb gene. A novel missense mutation in exon 27. *Haematologica* 91:1352-1359.
- Jin Y, Dietz HC, Montgomery RR, Bell WR, McIntosh I, Collier B, Bray PF. 1996. Glanzmann thrombasthenia. Cooperative between sequence variants in *Cis* during splice site selection. *J Clin Invest* 98:1745-1754.
- Jin Y, Dietz HC, Nurden A, Bray PF. 1993. Single-strand conformation polymorphism analysis is a rapid and effective method for the identification of mutations and polymorphisms in the gene for GP IIIa. *Blood* 82:2281-2288.
- Kato A, Yamamoto K, Miyazaki S, Jung SM, Maoi M, Aoki N. 1992. Molecular basis for Glanzmann's thrombasthenia (GT) in a compound heterozygote with glycoprotein IIb gene : a proposal for the classification of GT based on the biosynthetic pathway of glycoprotein IIb-IIIa complex. *Blood* 79:3212-3218.
- Lanza F, Stierlé A, Fournier D, Morales M, André G, Nurden AT, Cazenave JP. 1992. A new variant of Glanzmann's thrombasthenia (Strasbourg I). Platelets with functionally defective glycoprotein IIb-IIIa. Complexes and a glycoprotein IIIa<sup>214</sup> Arg--><sup>214</sup> Trp mutation. *J Clin Invest* 89:1995-2004.
- Laskowski RA, Mac Arthur MW, Moss DS, Thornton JM. 1993. PROCHECK : A program to check the stereochemical quality of protein structures. *J Appl Cryst* 26:286-291.
- Loftus JC, O'Toole TE, Plow EF, Glass A, Frelinger III AL, Ginsberg MH. 1990. A  $\beta_3$  integrin mutation abolishes ligand binding and alters divalent cation-dependent conformation. *Science* 249:915-918.



- 1  
2  
3 Mitchell WB, Li JH, Singh F, Michelson AD, Bussel J, Collier BS, French DL. 2003. Two  
4 novel mutations in the  $\alpha$ IIB calcium-binding domains identify hydrophobic regions  
5 essential for  $\alpha$ IIB $\beta$ 3 biogenesis. *Blood* 101:2268-2276.  
6  
7  
8 Morel-Kopp MC, Kaplan C, Proulle V, Jallu V, Melchior C, Peyruchaud O, Aurousseau MH,  
9 Kieffer N. 1997. A three amino acid deletion in glycoprotein IIIa is responsible for  
10 type I Glanzmann's thrombasthenia : Importance of residues Ile325 Pro326 Gly327 for  
11 B3 integrin subunit association. *Blood* 90:669-677.  
12  
13  
14 Morel-Kopp MC, Melchior C, Chen P, Ammerlaan W, Lecompte T, Kaplan C, Kieffer N.  
15 2002. A naturally occurring point mutation in the  $\beta$ 3 integrin MIDAS-like domain  
16 affects differently  $\alpha$ v $\beta$ 3 and  $\alpha$ IIB $\beta$ 3 receptor function. *Thromb Haemost* 86:1425-  
17 1434.  
18  
19  
20 Nair S, Ghosh K, Shetty S, Mohanty D. 2004. A novel Ser123Pro substitution in the MIDAS  
21 domain of integrin  $\beta$ 3 associated with variant Glanzmann's thrombasthenia in an Indian  
22 patient. *Haematologica* 89:1529-1530.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Negrier C, Vinciguerra C, Attali O, Grenier C, Larcher ME, Dechavanne M. 1998.  
Illegitimate transcription : its use for studying genetic abnormalities in lymphoblastoid  
cells from patients with Glanzmann thrombasthenia. *Br J Haematol* 100:33-39.
- Newman PJ, Seligsohn U, Lyman S, Collier BS. 1991. The molecular genetic basis of  
Glanzmann thrombasthenia in the iraqi-jewish and arab populations in Israel. *Proc  
Natl Acad Sci USA* 88:3160-3164.
- Nurden AT. 2005. Qualitative disorders of platelets and megakaryocytes. *J Thromb Haemost*  
3:1773-1782.
- Nurden AT, Breillat C, Jacquelin B, Combrie R, Freedman J, Blanchette VS, Schmutz M.  
2004. Triple heterozygosity in the integrin  $\alpha$ IIB subunit in a patient with Glanzmann's  
thrombasthenia. *J Thromb Haemost* 2:813-819.
- Nurden AT, Ruan J, Pasquet JM, Gauthier B, Combrié R, Kunicki T, Nurden P. 2002. A  
novel <sup>196</sup>Leu to Pro substitution in the  $\beta$ 3 subunit of the  $\alpha$ IIB $\beta$ 3 integrin in a patient  
with a variant form of Glanzmann thrombasthenia. *Platelets* 13:101-111.
- Peretz H, Rosenberg N, Landau M, Usher S, Nelson EJ, Mor-Cohen R, French DL, Mitchell  
BW, Nair SC, Chandy M, Collier BS, Srivastava A, Seligsohn U. 2006. Molecular  
diversity of Glanzmann thrombasthenia in southern India: new insights into mRNA  
splicing and structure-function correlations of alphaIIBbeta3 integrin (ITGA2B,  
ITGB3). *Hum Mutat* 27:359-369.
- Peterson JA, Gitter M, Kanack A, Curtis B, McFarland J, Aster RH. 2009. New low-  
frequency platelet glycoprotein polymorphisms associated with neonatal alloimmune  
thrombocytopenia. *Transfusion* published online : 10 Oct 2009
- Peterson JA, Nyree CE, Newman PJ, Aster RH. 2003. A site involving the "hybrid" and PSI  
homology domains of GPIIIa (beta 3-integrin subunit) is a common target for  
antibodies associated with quinine-induced immune thrombocytopenia. *Blood*  
101:937-942.

- 1  
2  
3 Quintanar A, Jallu V, Legros Y, Kaplan C. 1998. Human platelet antigen genotyping using a  
4 fluorescent SSCP technique with an automatic sequencer. *Br J Haematol* 103:437-444.  
5  
6  
7 Reese MG, Eeckman FH, Kulp D, Haussler D. 1997. Improved splice site detection in Genie.  
8 *J Comput Biol* 4:311-323.  
9  
10 Ruan J, Peyruchaud O, Alberio L, Valles G, Clemetson K, Bourre F, Nurden AT. 1998.  
11 Double heterozygosity of the GPIIb gene in a Swiss patient with Glanzmann's  
12 thrombasthenia. *Br J Haematol* 102:918-925.  
13  
14  
15 Schlegel N, Gayet O, Morel-Kopp MC, Wyler B, Hurtaud-Roux MF, Kaplan C, Mc Gregor J.  
16 1995. The molecular genetic basis of Glanzmann thrombasthenia in a Gipsy  
17 population in France. Identification of a new mutation on the  $\alpha$ -IIb gene. *Blood*  
18 86:977-982.  
19  
20  
21 Scott JP, Chao YL, Newman PJ, Ward CM. 1998. A frameshift mutation at Gly975 in the  
22 transmembrane domain of GPIIb prevents GPIIb-IIIa expression. Analysis of two  
23 novel mutations in a kindred with type I Glanzmann thrombasthenia. *Thromb*  
24 *Haemost* 80:546-550.  
25  
26  
27 Smith PJ, Zhang C, Wang J, Chew SL, Zhang MQ, Krainer AR. 2006. An increased  
28 specificity score matrix for the prediction of SF2/ASF-specific exonic splicing  
29 enhancers. *Hum Mol Genet* 15:2490-2508.  
30  
31  
32 Springer TA, Zhu J, Xiao T. 2008. Structural basis for distinctive recognition of fibrinogen  
33 gammaC peptide by the platelet integrin alphaIIbbeta3. *J Cell Biol* 182:791-800.  
34  
35  
36 Stafford P, Garner SF, Rankin A, Kekomaki R, Watkins NA, Ouwehand WH. 2008. A single-  
37 nucleotide polymorphism in the human ITGB3 gene is associated with the platelet-  
38 specific alloantigen Va (HPA-17bw) involved in fetal maternal alloimmune  
39 thrombocytopenia. *Transfusion* 48:1432-1438.  
40  
41  
42 Tadokoro S, Tomiyama Y, Honda S, Arai M, Yamamoto N, Shiraga M, Kosugi S, Kanakura  
43 Y, Kurata Y, Matsuzawa Y. 1998. A Gln747 -->Pro substitution in the  $\alpha$ IIb subunit is  
44 responsible for a moderate  $\alpha$ IIb $\beta$ <sub>3</sub> deficiency in Glanzmann thrombasthenia. *Blood*  
45 92:2750-2758.  
46  
47  
48 Tadokoro S, Tomiyama Y, Honda S, Kashiwagi H, Kosugi S, Shiraga M, Kiyoi T, Kurata Y,  
49 Matsuzawa Y. 2002. Missense mutations in the  $\beta$ <sub>3</sub> subunit have a different impact on  
50 the expression and function between  $\alpha$ IIb $\beta$ <sub>3</sub> and  $\alpha$ <sub>v</sub> $\beta$ <sub>3</sub>. *Blood* 99:931-938.  
51  
52  
53 Valentin N, Visentin GP, Newman PJ. 1995. Involvement of the cystein-rich domain of  
54 glycoprotein IIIa in the expression of the human platelet alloantigen,  $PI^{A1}$ : Evidence  
55 for heterogeneity in the humoral response. *Blood* 85:3028-3033.  
56  
57  
58 Wang GS, Cooper TA. 2007. Splicing in disease: disruption of the splicing code and the  
59 decoding machinery. *Nat Rev Genet* 8:749-761.  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

glycoprotein IIIa is responsible for the formation of the Pena/Penb alloantigen system.  
J Clin Invest 90:2038-2043.

Ward CM, Kestin AS, Newman PJ. 2000. A Leu262Pro mutation in the integrin  $\beta 3$  subunit in an  $\alpha$ IIB- $\beta 3$  complex that binds fibrin but not fibrinogen. Blood 96:161-169.

Xiao T, Takagi J, Coller BS, Wang JH, Springer TA. 2004. Structural basis for allostery in integrins and binding to fibrinogen-mimetic therapeutics. Nature 432:59-67.

Zhu J, Luo BH, Xiao T, Zhang C, Nishida N, Springer TA. 2008. Structure of a complete integrin ectodomain in a physiologic resting state and activation and deactivation by applied forces. Mol Cell 32:849-861.

For Peer Review

**Figure legends :****Figure 1 : Flow cytometry analysis of the expression of the mutant forms of  $\alpha$ IIB $\beta$ 3.**

Transiently transfected COS-7 cells expressing the wild type (WT) or the following mutants of  $\alpha$ IIB $\beta$ 3, p.S926L, p.Q595H, p.V903F ( $\alpha$ IIB) and p.C38Y, p.M118R, p.G221D, p.K253 ( $\beta$ 3) were tested in flow cytometry by using Moabs SZ22 (hatched bar), SZ21 (dot bar) and P2 (white bar) or irrelevant mouse IgG (black bar). As control, untransfected cells (NT) were also tested. Following Moab incubation, bound IgGs were detected by using a polyclonal goat anti-mouse IgG conjugated to FITC. Washed cells were analyzed on an Epic-XML flow cytometer. Except for the  $\alpha$ IIB p.Q595H one, all mutants showed a markedly decreased expression of the complex when compared to the wild type. Results represent the mean  $\pm$  SD of 3 experiments.

**Figure 2 : Western blot analysis of the Y38  $\beta$ 3 form of the  $\alpha$ IIB $\beta$ 3 complex expressed in COS-7 cells.** SDS lysates from normal platelets (lane a), non-transfected COS-7 cells (lane b), or COS-7 cells expressing the wild-type (lane c) or the Y38  $\beta$ 3 (lane d) forms of the complex were subjected to SDS-polyacrylamide gel electrophoresis under non reduced conditions and transferred to nitrocellulose membrane. Following incubation with Moabs SZ22, XIIF9, SZ21 and AP3, membrane bound IgGs were revealed by using anti-mouse IgG conjugated to peroxidase and the enhanced chemiluminescence detection technique.

**Figure 3 : Location and modeling of the electrostatic effect of the p.K253M substitution.** (A) and (B) Ribbon diagram of the  $\alpha$ IIB  $\beta$ -propeller (in blue) and  $\beta$ 3  $\beta$ I (in red) complexed domains. For the  $\beta$ I domain, The residue 253 is shown in green with (A) a lysine and (B) a methionine, while for  $\beta$ -propeller partners the P228 is shown in yellow, the F231 in

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

grey, the W262 in pink and the D232 in orange. (C) and (D) Electrostatics visualization of the single point mutation p.K253M with (C) wild K253  $\beta 3$  and (D) mutated M253  $\beta 3$  subunits. The visualizations are done using PyMol software (<http://www.pymol.org>).

**Figure 4 : Localization of the V903 and S926 on a ribbon diagram of  $\alpha$ I**IIb**.** This  $\alpha$ I**IIb** cartoon shows the internal location of the V903 and S926 side chains (orange, ball and stick representation) in the calf-2 composed by the N-terminal part of the  $\beta$ -subunit of  $\alpha$ I**IIb** (blue) and the C-terminal part of the  $\alpha$ -subunit (yellow and red respectively for  $\beta$ -sheet and  $\alpha$ -helix). This ribbon diagram was obtained from the resolved structure of  $\alpha$ I**IIb** (PDB code 3FCS) using the Swiss-PdbViewer software (version 4.0.1).

**Figure 5 : Localization of the allelic variants of the  $\beta$ -I domain of  $\beta 3$ .** Residues identified in this study and from the literature as involved in GT or neonatal alloimmune thrombocytopenia and that locate in the  $\beta$ -I domain of  $\beta 3$  are reported on its primary sequence (panel A) and a ribbon diagrams showing the  $\beta$ -I domain and a partial view of the  $\beta$ -propeller of  $\alpha$ I**IIb** (panel B). Allelic variants of residues highlighted in magenta, blue, green are respectively responsible for variant, II and I GT phenotypes or yellow for those concerning alloimmunisation. The GT type induced by the allelic variant of the H192 (grey) is not reported [Peretz et al., 2006]. Amino-acids identified by the energy analyze as in contact with the  $\alpha$ I**IIb**  $\beta$ -propeller are underlined. The primary sequence (panel A) identifies the secondary structures [Xiao et al., 2004] and amino-acids responsible for MIDAS ( $\star$ ), ADMIDAS ( $\bullet$ ) and SyMBS ( $\blacklozenge$ ) sites [Springer et al., 2008]. As known, variant GT mutations GT specifically affect metal ion binding sites but other mutants concern residues evenly distributed along the primary sequence. Boxed residues have been identified in this study. The ribbon diagram (panel B) was obtained from the resolved structure of  $\alpha$ I**IIb** (PDB file 2VDL) using the Swiss-

1  
2  
3 PdbViewer software (version 4.0.1).  $\alpha$ -helix (red) and  $\beta$ -sheet (yellow) are shown. All  
4  
5 residues whose mutants are involved in type II GT lay on the  $\beta$ -I surface except for the L117,  
6  
7 some of them (underlined amino-acids) directly contacting the  $\beta$ -propeller. The Met 118 and  
8  
9 Gly 221 locate inside the  $\beta$ -I domain and the Lys 253 points out to the  $\beta$ -propeller surface  
10  
11 however their identified allelic variants completely prevent the  $\alpha$ IIb $\beta$ 3 complex expression at  
12  
13 the platelet surface.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

Table 1 :  $\alpha$ IIB mutations identified by genomic analysis<sup>a</sup>

N <sup>o</sup>	Proband	GT <sup>b</sup>	Ex/IVS <sup>c</sup>	Sequence alteration <sup>d</sup>	Status <sup>e</sup>	Restriction enzyme	Predicted HGVS <sup>f</sup>	Predicted Classical <sup>g</sup>	Accession number <sup>h</sup>
1	CabGT-1	I	Ex 4	c.531T>A	Hom.	Hga I	p.C177*	p.C146*	ss131010618
2	CabGT-2	I	Ex 10	c.917dupA	Het.	Alw I	p.R307Efs*22	p.R276E + FS <sup>k</sup>	ss131007603
3			Ex 19	c.1912dup T	Het.	No enzyme	p.C639Mfs*22	p.C608M + FS <sup>k,l</sup>	ss 131007606
4	CabGT-3	I	Ex 13	c.1214T>C	Het.	Sfc I	p.I405T	p.I374T <sup>l</sup>	ss 131007609
5			IVS14	c.1442-13_1442-1del	Het.	Hph I	Splice <sup>i</sup>	Splice	ss 131007612
6	CabGT-4	I (car <sup>w</sup> )	IVS14	c.1442-13_1442-1del	Het.	Hph I	Splice	Splice	ss 131007612
7	CabGT-5	I	IVS14	c.1442-13_1442-1del NI <sup>j</sup>	Het.	Hph I	Splice	Splice	ss 131007612
8	CabGT-6	I	Ex 14	c.1413C>G	Het.	Dde I	p.Y471*	p.Y440* <sup>m</sup>	ss 131007615
9			Ex 27	c.2800G>T	Het.	Mae III / Tsp45 I	p.V934F	p.V903F	ss 131007617
10	CabGT-7	I	Ex 16	c.1563T>A NI	Het.	No enzyme	p.C521*	p.C490*	ss 131007620
11	CabGT-8	NC <sup>x</sup>	Ex 17	c.1672C>T	Hom.	Mse I	p.Q558*	p.Q527*	ss 131007623
12	CabGT-9	I	IVS18	c.1879-2A>G	Hom.	Xba I	Splice	Splice	ss 131007625
13	CabGT-10	I	Ex 18	c.1878G>C	Hom.	Hpy CH IV	p.Q626H / Splice	p.Q595H / Splice	ss 131007628
14	CabGT-11	I	Ex 18	c.1787T>C	Hom.	No enzyme	p.I596T	p.I565T <sup>n</sup>	ss 131008441
15	CabGT-12	II	Ex 21	c.2113T>C	Het.	Fok I	p.C705R	p.C674R <sup>l,o,p</sup>	ss 131007632
16			IVS29	c.3060+2T>C	Het.	Hph I	p.V982_K1020del	p.V951_K989del <sup>q</sup>	ss 131007634
17	CabGT-13	I	Ex 23	c.2333A>C	Hom.	Bsr I / TspR I	p.Q778P	p.Q747P <sup>r</sup>	ss 131007637
18	CabGT-14	I	Ex 27	c.2748_2757del	Hom.	Rsa I	p.T917Sfs181	p.T886S + FS	ss 131007639
19	CabGT-15	I (car)	Ex 27	c.2800G>T	Het.	Mae III / Tsp45 I	p.V934F	p.V903F	ss 131007617
20	CabGT-16	I	Ex 28	c.2870C>T	Hom.	Hpy CH4 V	p.S957L	p.S926L	ss 131007642
21	CabGT-17	II	Ex 28	c.2929C>T	Het.	Ava I	p.R977*	p.R946* <sup>l,s,t</sup>	ss 131007644
22			NI						
23	CabGT-18	NC	Ex 29	c.2944G>A	Het.	Bst NI	p.V982M	p.V951M <sup>u,v</sup>	ss 131007647
24			Ex 29	c.2965G>A	Het.	Nci I	p.A989T	p.A958T <sup>u,v</sup>	ss 131007649
25				NI					

<sup>a</sup> cDNA ITGA2B GenBank accession number NM\_000419.3<sup>b</sup> GT : Glanzmann type

<sup>c</sup> Ex : Exon ; IVS : intronic sequence

<sup>d</sup> Nomenclature according to HGVS : +1 corresponds to the A of ATG translation initiation codon

<sup>e</sup> Status : Het. : heterozygous; Hom. : Homozygous

<sup>f</sup> Predicted effect, nomenclature according to HGVS : +1 corresponds to the initiating Met (signal peptide included)

<sup>g</sup> Predicted effect, classical nomenclature : +1 corresponds to the first amino acid of the mature  $\alpha$ IIb

<sup>h</sup> NCBI dSNP accession number

<sup>i</sup> Splice defect based upon modeling with GENSCAN / NNSPLICE and ESEFinder

<sup>j</sup> NI : 2<sup>nd</sup> mutation not Identified

<sup>k</sup> FS : Frame shift

References : <sup>l</sup> D'Andrea et al., 2002 [D'Andrea et al., 2002]; <sup>m</sup> Scott et al., 1998 [Scott et al., 1998]; <sup>n</sup> Ruan et al., 1998 [Ruan et al., 1998]; <sup>o</sup> Gonzalez-Manchon et al., 1999 [Gonzalez-Manchon et al., 1999]; <sup>p</sup> Mitchell et al., 2003 [Mitchell et al., 2003]; <sup>q</sup> French et al., 1997 [French and Coller, 1997]; <sup>r</sup> Tadokoro et al., 1998 [Tadokoro et al., 1998]; <sup>s</sup> Basani et al., 2000 [Basani et al., 2000]; <sup>t</sup> Peretz et al., 2006 [Peretz et al., 2006]; <sup>u</sup> Nurden et al., 2004 [Nurden et al., 2004]

<sup>v</sup> V951M is responsible for GT but not A958T [Nurden et al., 2004]

<sup>w</sup> Car : carrier

<sup>x</sup> NC : Not communicated



Table 2 :  $\beta 3$  mutations identified by genomic analysis<sup>a</sup>

N°	Proband	GT <sup>b</sup>	Ex/IVS <sup>c</sup>	Sequence alteration <sup>d</sup>	Status <sup>e</sup>	Restriction enzyme	Predicted HGVS <sup>f</sup>	Predicted Classical <sup>g</sup>	Accession number <sup>h</sup>
19	CabGT-19	NC <sup>j</sup>	IVS2	c.166-2A>G	Hom.	Apa I	Splice <sup>i</sup>	Splice	ss 131007652
20	CabGT-20	I	Ex 2	c.100C>T	Hom.	Hpy CH4 IV	p.R34*	p.R8* <sup>l</sup>	ss 131007655
21	CabGT-21	I	Ex 3	c.191G>A NI <sup>k</sup>	Het.	Tsp 45I	p.C64Y	p.C38Y	ss 131007657
22	CabGT-22	I	Ex 4	c.431T>G	Hom.	Bsm F1	p.M144R	p.M118R	ss 131007660
23	CabGT-23	II	Ex 5	c.665 T>C	Hom.	Acy I	p.L222P	p.L196P <sup>m</sup>	ss 131007662
24	CabGT-24	I	Ex 5 Ex 6	c.740 G>A c.836 A>T	Het. Het.	Fok I Fat I	p.G247D p.K279M	p.G221D p.K253M	ss 131007665 ss 131007667
25	CabGT-25	NC	IVS8	c.1125+3_1125+6del NI	Het.	No enzyme	Splice	Splice	ss 131007670
26	CabGT-26	I	Ex 9 Ex 9	c.1143C>A c.1260G>A	Het. Het.	Taq I Hph I	p.K376_T420del	p.K350_T394del <sup>n</sup>	ss 131007672 ss 131007674

<sup>a</sup>cDNA ITGB3 GenBank accession number NM\_000212.2

<sup>b</sup>GT : Glanzmann type I or II (< 5% or 10 – 25 % of normal platelet  $\alpha$ IIB $\beta$ 3[Caen, 1989]). See Supp\_mat tableS1 for detailed data.

<sup>c</sup>Ex : Exon ; IVS : intronic sequence

<sup>d</sup>Nomenclature according to HGVS : +1 corresponds to the A of ATG translation initiation codon

<sup>e</sup>Status : Het. : heterozygous; Hom. : Homozygous

<sup>f</sup>Nomenclature according to HGVS : +1 corresponds to the initiating Met (signal peptide included)

<sup>g</sup>Classical nomenclature : +1 corresponds to the first amino acid of the mature  $\beta 3$

<sup>h</sup>NCBI dSNP accession number

<sup>i</sup>Splice defect based upon modeling with GENSCAN, NNSPLICE and ESEFinder

<sup>j</sup>NC : Not communicated

<sup>k</sup>NI : 2<sup>nd</sup> mutation not Identified

<sup>l</sup>References : <sup>l</sup>Negrier et al., 1998 [Negrier et al., 1998]; <sup>m</sup>Nurden et al., 2002 [Nurden et al., 2002]; <sup>n</sup>Jin et al., 1996 [Jin et al., 1996]

Table 3 : Energetic effects of the Lys 253 Met substitution in the  $\alpha$ I**II**b/  $\beta$ 3 interaction

Energy of :		Lys 253			Met 253		
		Rank	Value (kcal/mol)	$\alpha$ I <b>II</b> b contact	rank	Value (kcal/mol)	$\alpha$ I <b>II</b> b contact
<b><math>\alpha</math>I<b>II</b>b<math>\beta</math>3 Interface</b>	<b>Electrostatic (4r) Energy</b>	/	-25.14	/	/	-16.08	/
	<b>Desolvation Free Energy</b>	/	-1.09	/	/	-6.81	/
	<b>Van der Waals (CHARMm19)</b>	/	-4894.63	/	/	-4903.95	/
<b>Residue</b>	<b>Binding free energy</b>	1 <sup>st</sup> negative	-5.541	/	12 <sup>th</sup> negative	-1.309	/
	<b>Desolvation free energy</b>	1 <sup>st</sup> positive	4.124	/	6 <sup>th</sup> negative	-1.347	/
	<b>Electrostatic energy</b>	1 <sup>st</sup> negative	-9.666	/	/	<i>low</i>	/
<b>Contact</b>	<b>Electrostatic</b>	1 <sup>st</sup> negative	-6.715	D232	/	/	/
		6 <sup>th</sup> negative	-2.076	F231	/	/	/
		13 <sup>th</sup> negative	-0.788	P228	/	/	/
		10 <sup>th</sup> positive	0.670	W262	6 <sup>th</sup> negative	-0.935	W262
	<b>Free energy</b>	1 <sup>st</sup> negative	-5.517	D232	/	/	/
		8 <sup>th</sup> negative	-1.166	F231	/	/	/
		2 <sup>th</sup> positive	2.057	W262	/	/	/




Figure 1 : Flow cytometry analysis of the expression of the mutant forms of  $\alpha$ IIb $\beta$ 3.  
235x165mm (96 x 96 DPI)





Figure 2 : Western blot analysis of the Y38  $\beta_3$  form of the  $\alpha_{IIb}\beta_3$  complex expressed in COS-7 cells.  
155x177mm (96 x 96 DPI)



CMYK color scale

Figure 3 : Location and modeling of the electrostatic effect of the p.K253M substitution.  
193x158mm (96 x 96 DPI)





Figure 3 : Location and modeling of the electrostatic effect of the p.K253M substitution.  
193x158mm (96 x 96 DPI)




CYMK color scale

Figure 4 : Localization of the V903 and S926 on a ribbon diagram of aIIb.  
135x115mm (96 x 96 DPI)

W

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





RGB color scale

Figure 4 : Localization of the V903 and S926 on a ribbon diagram of aIIb.  
135x115mm (96 x 96 DPI)









CMYK color scale

Figure 5 : Localization of the allelic variants of the  $\beta$ -I domain of  $\beta$ 3.  
212x200mm (96 x 96 DPI)



RGB color scale

Figure 5 : Localization of the allelic variants of the  $\beta$ -I domain of  $\beta$ 3.  
212x200mm (96 x 96 DPI)