

## **Clinical Image**

### **What is the patient's job?**

Alexis Descatha (1), Emmanuelle Deschamps(1)

(1) UVSQ AP-HP. Occupational health unit/ INSERM U687/ EMS of 92' district.  
Poincare Teaching Hospital, F-92380 Garches, France

**Address for reprints:** Dr. Alexis Descatha, Unité de pathologie professionnelle/ U687  
INSERM/SAMU92, CHU Poincaré, 104 bd Poincaré, 92380 Garches, France

Tel: +33 (1) 47 10 77 54; Fax: +33 (1) 47 10 77 68; email: alexis.descatha@rpc.aphp.fr

**One figure = 150 words (legend is the main manuscript file)**

### **Patient consent obtained**

### **Acknowledgments**

We would like to thank: Dr Potier, general practitioner, in charge of the patient, Dr Carlier, radiologist, who helps us to read the CT scan and Ms Laborde, English Supervisor in our University, who helps us to improve our English.

The CT scan shows characteristic but uncommon lesions, which are the results of almost 50 years of exposure.

A 55 year-old woman was suffering from pains in the feet. X-rays and the CT scan (sagittal CT scan reconstruction of the left foot) showed a bilateral flatfoot and a localized degenerative lesion of the tarsometatarsal joint. The joint between the medial cuneiform and the first metatarsal bone was particularly affected, so was the joint between the intermediate cuneiform and the second metatarsal bone.

Forty-nine years of dancing including 34 years as a professional dancer then as a teacher potentially caused the unusual localized osteoarthritis of the tarsometatarsal joint (1), or at least it accelerated the degenerative process.(2;3) One may suppose that repetitive or neglected microtraumas, especially when dancing on points,(4) led directly to the disorder or to an acquired flatfoot which worsened these lesions.(5) Because of the pain, she stopped dancing.

## References

- (1) Andersson S, Nilsson B, Hessel T, Saraste M, Noren A, Stevens-Andersson A et al. Degenerative joint disease in ballet dancers. *Clin Orthop Relat Res* 1989;(238):233-236.
- (2) Huwyler J. [Changes in the skeleton of the middle foot caused by professional classical dancing]. *Sportverletz Sportschaden* 1989; 3(1):14-20.
- (3) Teitz CC, Kilcoyne RF. Premature osteoarthrosis in professional dancers. *Clin J Sport Med* 1998; 8(4):255-259.
- (4) Nilsson C, Leanderson J, Wykman A, Strender LE. The injury panorama in a Swedish professional ballet company. *Knee Surg Sports Traumatol Arthrosc* 2001; 9(4):242-246.
- (5) Chu IT, Myerson MS, Nyska M, Parks BG. Experimental flatfoot model: the contribution of dynamic loading. *Foot Ankle Int* 2001; 22(3):220-225.

