

Table 2. Description of the disease and the technical aspects of HIPEC in the 7 patients

childbearing after the procedure. Y: years; m: months; PMP: pseudomyxoma peritonei; PCI: peritoneal carcinomatosis index; MMC: mitomycin-C.

Patient	Indication	Age at HIPEC (y/m)	PCI	Cytoreduction achieved (CC)	Drug used for HIPEC	Duration of HIPEC (minutes)	Mean temperature during HIPEC °C	Maximal temperature during HIPEC
1	PMP III	31y	26	CC-1	MMC – 17 mg	120	42° C	42° C
2	PMP II	21y 10m	10	CC-0	MMC – 20 mg/m ²	90	41° C	42.5° C
3	PMP I	34y 4m	1	CC-0	MMC– 40 mg	120	40° C	42.5° C
4	PMP I	26y 10m	11	CC-0	MMC– 50 mg	60	42° C	43.3° C
5	Epithelial mesothelioma	21y 2m	5	CC-0	Cisplatin – 250 mg/m ²	90	40° C	40° C
6	PMP I	29y 1m	-	CC-0	MMC + Cisplatin - 30 mg each	90	38.3° C	42° C
7	Papillary mesothelioma	17y 1m	10	CC-0	Cisplatin - 100 mg/m ²	90	40° C	40.5° C