



Role of CD4+CD25hiCD127lo/-FoxP3+ regulatory T lymphocytes in the pathogenesis of Behçet's disease in children

Tu-Anh Tran, Sylvia Monteil, Alexia Letierce, Benjamin Terrier, Guillaume Geri, David Saadoun, Isabelle Kone-Paut, Benoît Salomon, Michelle Rosenzwajg

► To cite this version:

Tu-Anh Tran, Sylvia Monteil, Alexia Letierce, Benjamin Terrier, Guillaume Geri, et al.. Role of CD4+CD25hiCD127lo/-FoxP3+ regulatory T lymphocytes in the pathogenesis of Behçet's disease in children. *Pediatric Rheumatology*, 2011, 9 (Suppl 1), pp.P84. inserm-00624793

HAL Id: inserm-00624793

<https://inserm.hal.science/inserm-00624793>

Submitted on 19 Sep 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

Role of CD4⁺CD25^{hi}CD127^{lo/-}FoxP3⁺ regulatory T lymphocytes in the pathogenesis of Behçet's disease in children

TA Tran^{1*}, S Monteil², A Letierce³, B Terrier², G Geri², D Saadoun⁴, I Kone-Paut¹, B Salomon⁵, M Rosenzweig²

From 18th Pediatric Rheumatology European Society (PReS) Congress
Bruges, Belgium. 14-18 September 2011

Introduction

Behçet's disease (BD) is an idiopathic multisystem recurrent inflammatory disorder. Physiopathology of BD shows a role of neutrophils and cytotoxic T lymphocytes.

Our aim

Were to assess the role of regulatory T lymphocytes (Tregs) in the pathogenesis of BD in children.

Patients and methods

19 patients with active BD (group A) and 8 patients with inactive BD (group B) were compared with 25 healthy controls (group C). Percentages of blood CD4⁺CD127^{lo/-}CD25^{hi}FoxP3⁺ Tregs and other T/B and NK cells subpopulations were analyzed by flow cytometry. The frequency of IL-17A and IFN- γ producing T cells was analyzed by flow cytometer from PBMC after 4 hours stimulation with PMA-ionomycin. We measured serum cytokines by Luminex and ELISA. We compared the 3 groups by using the Wilcoxon-Rank-signed test. Values were expressed as mean and median.

Results

Patients in the 3 groups (A, B, C respectively) were comparable in term of age and sex distribution (median age: 12.8, 9.9 and 9.7; F/M = 1/1). No differences were observed between the 3 groups concerning the absolute number of lymphocytes, CD4⁺ T cells and the percentage of total Tregs (median: A: 1.9, B:1.1, C:2.8). Percentages of naïve Treg/memory Treg and markers of Treg function (GITR, LAP, CD152, DR) were also

similar in the 3 groups. However, there was increased CD8⁺ T cells count in the BD patients groups compared to healthy controls (A: 552 $\pm$ 361, p=0.18; B: 627 $\pm$ 159, p=0.04, C: 479 $\pm$ 209). The NK cell (CD3-CD16+CD56+) were highest in group C compared to group A (p=0.4) or B (p=0.001). IL-17A secreting CD4⁺ T cells were significantly higher in active BD patients (n=6) compared to controls (n=6) (5.3 $\pm$ 2 vs 2.5 $\pm$ 1.47, p=0.043). Serum IL-6 level was significantly higher in BD populations compared to controls subjects (A: 4.3 $\pm$ 1.22 vs C:3 $\pm$ 0.7 pg/ml, p=0.016).

Conclusion

There is no deficit of Tregs number in BD patients. The high rate of peripheral IL-17 secreting CD4⁺ T cells suggests a possible role of Th17 cells in the occurrence of BD attacks. The Tregs functional ability to regulate CD4 and CD8 T cells needs to be studied further.

Author details

¹Department of Paediatrics, Pediatric Rheumatology. CEREMAI Bicêtre Hospital, University of Paris Sud., France. ²Service de Biothérapies/ UPMC CNRS 7211 INSERM 959. La Pitié Salpêtrière University Hospital. Paris, France. ³Unité de Recherche Clinique Paris Sud. Bicêtre University Hospital. Le Kremlin Bicêtre. France. ⁴Department of Internal Medicine. La Pitié Salpêtrière University Hospital, Paris, France. ⁵Unité 2. UPMC-CNRS U7087. La Pitié Salpêtrière University Hospital, Paris, France.

Published: 14 September 2011

doi:10.1186/1546-0096-9-S1-P84

Cite this article as: Tran et al.: Role of CD4⁺CD25^{hi}CD127^{lo/-}FoxP3⁺ regulatory T lymphocytes in the pathogenesis of Behçet's disease in children. *Pediatric Rheumatology* 2011 **9**(Suppl 1):P84.

¹Department of Paediatrics, Pediatric Rheumatology. CEREMAI Bicêtre Hospital, University of Paris Sud., France
Full list of author information is available at the end of the article