



HAL
open science

GnRH neurons directly listen to the periphery.

Vincent Prevot

► **To cite this version:**

Vincent Prevot. GnRH neurons directly listen to the periphery.. *Endocrinology*, 2011, 152 (10), pp.3589-91. 10.1210/en.2011-1544 . inserm-00626448

HAL Id: inserm-00626448

<https://inserm.hal.science/inserm-00626448>

Submitted on 26 Sep 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Submission: September 26, 2011
News and Views

GnRH neurons directly listen to the periphery

Vincent Prevot

Inserm, Jean-Pierre Aubert Research Centre, U837, Development and Plasticity of the postnatal Brain, F-59000 Lille, France; Univ Lille Nord de France, F-59000 Lille, France; UDSL, School of Medicine, F-59000 Lille, France; CHRU Lille, Department of Neurosurgery, Hôpital Roger Salengro, F-59000 Lille, France

The survival of a mammalian species relies on the ability of its individuals to promptly, efficiently and reproducibly transmit homeostatic signals to the hypothalamic neuronal population that releases gonadotropin-releasing hormone (GnRH) and controls fertility. This requires the intervention of neuronal networks that make continual adjustments to brain function in response to moment-to-moment changes in physiological inputs throughout life (1-6). Intriguingly, a report in this issue of *Endocrinology* (7) reveals that in complement to these transsynaptic regulatory mechanisms, GnRH neurons themselves may have direct access to peripheral information via the extension of dendrites outside the blood-brain barrier (BBB).

In rodents, the cell bodies of GnRH neurons are diffusely distributed throughout the basal forebrain and are particularly abundant in the preoptic hypothalamic region at the level of the organum vasculosum of the lamina terminalis (OVLT). They exhibit a simple bipolar morphology with one or two very long dendritic processes that can extend up to 1 mm (8, 9). GnRH is released from neuroendocrine terminals in the median eminence at the base of the hypothalamus into the pituitary portal circulation, and stimulates the synthesis and release of the gonadotropins luteinizing hormone (LH) and follicle stimulating hormone (FSH), which in turn act on the ovaries and testes to regulate the secretion of sex steroids and the production of eggs and sperm. In addition to sending abundant axonal projections to the median eminence, GnRH neurons also extend fibers towards the OVLT (10, 11), a discrete

periventricular area with highly permeable, fenestrated endothelial cells that, as in the median eminence (12, 13), form a brain-periphery interface outside the BBB (12) - features that are characteristic of the circumventricular organs. However, in contrast to the median eminence, in which the primary direction of communication is from neural tissue (neuroendocrine terminals) to the circulation, it is thought that the primary functional consequence of the absence of a BBB in the OVLT is that this allows circulating factors that do not normally cross the barrier to gain direct access to neurons within the central nervous system (14).

The presence of GnRH fibers in the OVLT has been known since the first description of the neuroanatomical distribution of GnRH neurons in the mammalian brain by Barry and colleagues in 1973 (10). The finding by Broadwell and Brightman in 1976 (15) using intravenously injected horseradish peroxidase (HRP) as a tracer, that the OVLT is a neurohemal region with permeable blood vessels, together with studies by the Kordon laboratory in 1977 (16) demonstrating that microdissected OVLT explants release GnRH *in vitro* as efficiently as median eminence explants, has led to the longstanding concept that the OVLT contains GnRH neurosecretory axons.

The current study by Herde *et al.* (7), however, elegantly shows that the GnRH processes in the OVLT are mainly dendritic rather than axonal in nature, and that they may play an important role in the regulation of the GnRH system. Using a combination of neuroanatomical and electrophysiological approaches, the authors (7) convincingly demonstrate that about 70% of the GnRH neurons located in the vicinity of the OVLT (> 20% of the total population of GnRH neurons in the rostral preoptic region) extend dendrites into this structure. Using the same HRP method as that employed by Broadwell and Brightman (15), they show that most of these dendrites lie outside the BBB and that they respond electrophysiologically to locally applied transmitters, including peptides that are unlikely to cross the BBB, such as kisspeptin. In addition, using c-Fos as a marker of neuronal activation, the authors demonstrate that the majority of GnRH neurons with dendrites in the OVLT are activated at the time of the GnRH/LH surge, and thus conclude that these GnRH neurons, identified more

than two decades ago as probably contributing to the GnRH/LH surge (17, 18), are exposed to blood-borne molecules through their dendrites. This work thus identifies a new mechanism for the modulation of GnRH neurons that considerably extends the range of factors that they can directly sense and integrate for the control of fertility. These findings also raise the exciting possibility that these GnRH neurons, which have direct access to plasma vs. brain levels of hormones, glucose or other metabolic substrates, act as metabolic sensing neurons (19) that could be used by the neuroendocrine brain to match fertility levels to the homeostatic status of the individual (20). In line with this idea, recent studies have demonstrated that GnRH neurons in brain slices are capable of directly sensing glucose in a concentration-dependent manner and that this glucosensing is modulated by gonadal steroids (21, 22).

The study by Herde *et al.* (7) showing that subsets of GnRH neurons have direct access to the periphery greatly improves our understanding of how abrupt changes in the plasma level of molecules can acutely modulate pulsatile GnRH/LH secretion (23, 24), thus establishing a direct and permanent link between metabolism and fertility (20). From a practical point of view, the realization that the GnRH system directly senses molecules in the periphery opens up avenues for future research and the development of new therapeutic tools for infertility.

Acknowledgments

This work was supported by the Agence National pour la Recherche (ANR, France) grants ANR-07-NEURO-026-03 and ANR-09-BLAN-0267, and the Fondation pour la Recherche Médicale (Equipe FRM 2005, France).

Address all correspondence and request of reprints to: Vincent Prevot, Inserm U837, Bâtiment Biserte, 59045 Lille cedex, France. E-mail: vincent.prevot@inserm.fr

Disclosure summary: The author has nothing to declare.

References

1. **Donato J, Jr., Cravo RM, Frazao R, Gautron L, Scott MM, Lachey J, Castro IA, Margatho LO, Lee S, Lee C, Richardson JA, Friedman J, Chua S, Jr., Coppari R, Zigman JM, Elmquist JK, Elias CF** 2011 Leptin's effect on puberty in mice is relayed by the ventral premammillary nucleus and does not require signaling in Kiss1 neurons. *J Clin Invest* 121:355-368
2. **Hill JW, Elias CF, Fukuda M, Williams KW, Berglund ED, Holland WL, Cho YR, Chuang JC, Xu Y, Choi M, Lauzon D, Lee CE, Coppari R, Richardson JA, Zigman JM, Chua S, Scherer PE, Lowell BB, Bruning JC, Elmquist JK** 2010 Direct insulin and leptin action on pro-opiomelanocortin neurons is required for normal glucose homeostasis and fertility. *Cell Metab* 11:286-297
3. **Donato J, Jr., Silva RJ, Sita LV, Lee S, Lee C, Lacchini S, Bittencourt JC, Franci CR, Canteras NS, Elias CF** 2009 The ventral premammillary nucleus links fasting-induced changes in leptin levels and coordinated luteinizing hormone secretion. *J Neurosci* 29:5240-5250
4. **Leshan RL, Louis GW, Jo YH, Rhodes CJ, Munzberg H, Myers MG, Jr.** 2009 Direct innervation of GnRH neurons by metabolic- and sexual odorant-sensing leptin receptor neurons in the hypothalamic ventral premammillary nucleus. *J Neurosci* 29:3138-3147
5. **I'Anson H, Manning JM, Herbosa CG, Pelt J, Friedman CR, Wood RI, Bucholtz DC, Foster DL** 2000 Central inhibition of gonadotropin-releasing hormone secretion in the growth-restricted hypogonadotropic female sheep. *Endocrinology* 141:520-527
6. **Berriman SJ, Wade GN, Blaustein JD** 1992 Expression of Fos-like proteins in gonadotropin-releasing hormone neurons of Syrian hamsters: effects of estrous cycles and metabolic fuels. *Endocrinology* 131:2222-2228
7. **Herde MK, Geist K, Campbell RE, Herbison AE** 2011 Gonadotropin-releasing hormone neurons extend complex dendritic trees outside the blood-brain barrier. *Endocrinology*:in press
8. **Campbell RE, Gaidamaka G, Han SK, Herbison AE** 2009 Dendro-dendritic bundling and shared synapses between gonadotropin-releasing hormone neurons. *Proc Natl Acad Sci U S A* 106:10835-10840
9. **Campbell RE, Han SK, Herbison AE** 2005 Biocytin filling of adult gonadotropin-releasing hormone neurons in situ reveals extensive, spiny, dendritic processes. *Endocrinology* 146:1163-1169
10. **Barry J, Dubois MP, Poulain P** 1973 LRF producing cells of the mammalian hypothalamus. A fluorescent antibody study. *Z Zellforsch Mikrosk Anat* 146:351-366
11. **Gross DS** 1976 Distribution of gonadotropin-releasing hormone in the mouse brain as revealed by immunohistochemistry. *Endocrinology* 98:1408-1417
12. **Ciofi P, Garret M, Lapirot O, Lafon P, Loyens A, Prevot V, Levine JE** 2009 Brain-endocrine interactions: a microvascular route in the mediobasal hypothalamus. *Endocrinology* 150:5509-5519
13. **Mullier A, Bouret SG, Prevot V, Dehouck B** 2010 Differential distribution of tight junction proteins suggests a role for tanycytes in blood-hypothalamus barrier regulation in the adult mouse brain. *J Comp Neurol* 518:943-962
14. **Ferguson AV, Bains JS** 1996 Electrophysiology of the circumventricular organs. *Front Neuroendocrinol* 17:440-475
15. **Broadwell RD, Brightman MW** 1976 Entry of peroxidase into neurons of the central and peripheral nervous systems from extracerebral and cerebral blood. *J Comp Neurol* 166:257-283
16. **Rotsztein WH, Charli JL, Pattou E, Kordon C** 1977 Stimulation by dopamine of luteinizing hormone-releasing hormone (LHRH) release from the mediobasal hypothalamus in male rats. *Endocrinology* 101:1475-1483
17. **Hoffman GE, Lee WS, Attardi B, Yann V, Fitzsimmons MD** 1990 Luteinizing hormone-releasing hormone neurons express c-fos antigen after steroid activation. *Endocrinology* 126:1736-1741

18. **Lee WS, Smith MS, Hoffman GE** 1990 Luteinizing hormone-releasing hormone neurons express Fos protein during the proestrous surge of luteinizing hormone. *Proc Natl Acad Sci U S A* 87:5163-5167
19. **Levin BE, Magnan C, Dunn-Meynell A, Le Foll C** 2011 Metabolic sensing and the brain: who, what, where, and how? *Endocrinology* 152:2552-2557
20. **Furman M, Wade GN** 2007 Animal models in the study of nutritional infertility. *Curr Opin Endocrinol Diabetes Obes* 14:475-481
21. **Zhang C, Bosch MA, Levine JE, Ronnekleiv OK, Kelly MJ** 2007 Gonadotropin-releasing hormone neurons express K(ATP) channels that are regulated by estrogen and responsive to glucose and metabolic inhibition. *J Neurosci* 27:10153-10164
22. **Roland AV, Moenter SM** 2011 Glucosensing by GnRH neurons: inhibition by androgens and involvement of AMP-activated protein kinase. *Mol Endocrinol* 25:847-858
23. **Bucholtz DC, Vidwans NM, Herbosa CG, Schillo KK, Foster DL** 1996 Metabolic interfaces between growth and reproduction. V. Pulsatile luteinizing hormone secretion is dependent on glucose availability. *Endocrinology* 137:601-607
24. **Nagatani S, Bucholtz DC, Murahashi K, Estacio MA, Tsukamura H, Foster DL, Maeda KI** 1996 Reduction of glucose availability suppresses pulsatile luteinizing hormone release in female and male rats. *Endocrinology* 137:1166-1170