

# GnRH neurons directly listen to the periphery

Vincent Prevot \* *Centre de recherche Jean-Pierre Aubert INSERM : U837, Université du Droit et de la Santé - Lille II, FR*

\* Correspondence should be addressed to: Vincent Prevot <vincent.prevot@inserm.fr>

**MESH Keywords** Animals ; Blood-Brain Barrier ; Dendrites ; physiology ; Female ; Gonadotropin-Releasing Hormone ; physiology ; Hypothalamus ; physiology

The survival of a mammalian species relies on the ability of its individuals to promptly, efficiently and reproducibly transmit homeostatic signals to the hypothalamic neuronal population that releases gonadotropin-releasing hormone (GnRH) and controls fertility. This requires the intervention of neuronal networks that make continual adjustments to brain function in response to moment-to-moment changes in physiological inputs throughout life (1 –6). Intriguingly, a report in this issue of *Endocrinology* (7) reveals that in complement to these transsynaptic regulatory mechanisms, GnRH neurons themselves may have direct access to peripheral information via the extension of dendrites outside the blood-brain barrier (BBB).

In rodents, the cell bodies of GnRH neurons are diffusely distributed throughout the basal forebrain and are particularly abundant in the preoptic hypothalamic region at the level of the organum vasculosum of the lamina terminalis (OVLT). They exhibit a simple bipolar morphology with one or two very long dendritic processes that can extend up to 1 mm (8, 9). GnRH is released from neuroendocrine terminals in the median eminence at the base of the hypothalamus into the pituitary portal circulation, and stimulates the synthesis and release of the gonadotropins luteinizing hormone (LH) and follicle stimulating hormone (FSH), which in turn act on the ovaries and testes to regulate the secretion of sex steroids and the production of eggs and sperm. In addition to sending abundant axonal projections to the median eminence, GnRH neurons also extend fibers towards the OVLT (10, 11), a discrete periventricular area with highly permeable, fenestrated endothelial cells that, as in the median eminence (12, 13), form a brain-periphery interface outside the BBB (12) - features that are characteristic of the circumventricular organs. However, in contrast to the median eminence, in which the primary direction of communication is from neural tissue (neuroendocrine terminals) to the circulation, it is thought that the primary functional consequence of the absence of a BBB in the OVLT is that this allows circulating factors that do not normally cross the barrier to gain direct access to neurons within the central nervous system (14).

The presence of GnRH fibers in the OVLT has been known since the first description of the neuroanatomical distribution of GnRH neurons in the mammalian brain by Barry and colleagues in 1973 (10). The finding by Broadwell and Brightman in 1976 (15) using intravenously injected horseradish peroxidase (HRP) as a tracer, that the OVLT is a neurohemal region with permeable blood vessels, together with studies by the Kordon laboratory in 1977 (16) demonstrating that microdissected OVLT explants release GnRH *in vitro* as efficiently as median eminence explants, has led to the longstanding concept that the OVLT contains GnRH neurosecretory axons.

The current study by Herde *et al.* (7), however, elegantly shows that the GnRH processes in the OVLT are mainly dendritic rather than axonal in nature, and that they may play an important role in the regulation of the GnRH system. Using a combination of neuroanatomical and electrophysiological approaches, the authors (7) convincingly demonstrate that about 70% of the GnRH neurons located in the vicinity of the OVLT (> 20% of the total population of GnRH neurons in the rostral preoptic region) extend dendrites into this structure. Using the same HRP method as that employed by Broadwell and Brightman (15), they show that most of these dendrites lie outside the BBB and that they respond electrophysiologically to locally applied transmitters, including peptides that are unlikely to cross the BBB, such as kisspeptin. In addition, using c-Fos as a marker of neuronal activation, the authors demonstrate that the majority of GnRH neurons with dendrites in the OVLT are activated at the time of the GnRH/LH surge, and thus conclude that these GnRH neurons, identified more than two decades ago as probably contributing to the GnRH/LH surge (17, 18), are exposed to blood-borne molecules through their dendrites. This work thus identifies a new mechanism for the modulation of GnRH neurons that considerably extends the range of factors that they can directly sense and integrate for the control of fertility. These findings also raise the exciting possibility that these GnRH neurons, which have direct access to plasma vs. brain levels of hormones, glucose or other metabolic substrates, act as metabolic sensing neurons (19) that could be used by the neuroendocrine brain to match fertility levels to the homeostatic status of the individual (20). In line with this idea, recent studies have demonstrated that GnRH neurons in brain slices are capable of directly sensing glucose in a concentration-dependent manner and that this glucosensing is modulated by gonadal steroids (21, 22).

The study by Herde *et al.* (7) showing that subsets of GnRH neurons have direct access to the periphery greatly improves our understanding of how abrupt changes in the plasma level of molecules can acutely modulate pulsatile GnRH/LH secretion (23, 24), thus establishing a direct and permanent link between metabolism and fertility (20). From a practical point of view, the realization that the GnRH system directly senses molecules in the periphery opens up avenues for future research and the development of new therapeutic tools for infertility.

## Acknowledgements:

This work was supported by the Agence National pour la Recherche (ANR, France) grants ANR-07-NEURO-026-03 and ANR-09-BLAN-0267, and the Fondation pour la Recherche Médicale (Equipe FRM 2005, France).

## Footnotes:

Disclosure summary: The author has nothing to declare.

## References:

- 1 . Donato J Jr , Cravo RM , Frazao R , Gautron L , Scott MM , Lachey J , Castro IA , Margatho LO , Lee S , Lee C , Richardson JA , Friedman J , Chua S Jr , Coppari R , Zigman JM , Elmquist JK , Elias CF . 2011 ; Leptin's effect on puberty in mice is relayed by the ventral preamillary nucleus and does not require signaling in Kiss1 neurons . *J Clin Invest* . 121 : 355 - 368
- 2 . Hill JW , Elias CF , Fukuda M , Williams KW , Berglund ED , Holland WL , Cho YR , Chuang JC , Xu Y , Choi M , Lauzon D , Lee CE , Coppari R , Richardson JA , Zigman JM , Chua S , Scherer PE , Lowell BB , Bruning JC , Elmquist JK . 2010 ; Direct insulin and leptin action on pro-opiomelanocortin neurons is required for normal glucose homeostasis and fertility . *Cell Metab* . 11 : 286 - 297
- 3 . Donato J Jr , Silva RJ , Sita LV , Lee S , Lee C , Lacchini S , Bittencourt JC , Franci CR , Canteras NS , Elias CF . 2009 ; The ventral preamillary nucleus links fasting-induced changes in leptin levels and coordinated luteinizing hormone secretion . *J Neurosci* . 29 : 5240 - 5250
- 4 . Leshan RL , Louis GW , Jo YH , Rhodes CJ , Munzberg H , Myers MG Jr . 2009 ; Direct innervation of GnRH neurons by metabolic- and sexual odorant-sensing leptin receptor neurons in the hypothalamic ventral preamillary nucleus . *J Neurosci* . 29 : 3138 - 3147
- 5 . l'Anson H , Manning JM , Herbosa CG , Pelt J , Friedman CR , Wood RI , Bucholtz DC , Foster DL . 2000 ; Central inhibition of gonadotropin-releasing hormone secretion in the growth-restricted hypogonadotropic female sheep . *Endocrinology* . 141 : 520 - 527
- 6 . Berriman SJ , Wade GN , Blaustein JD . 1992 ; Expression of Fos-like proteins in gonadotropin-releasing hormone neurons of Syrian hamsters: effects of estrous cycles and metabolic fuels . *Endocrinology* . 131 : 2222 - 2228
- 7 . Herde MK , Geist K , Campbell RE , Herbison AE . 2011 ; Gonadotropin-releasing hormone neurons extend complex dendritic trees outside the blood-brain barrier . *Endocrinology* . in press
- 8 . Campbell RE , Gaidamaka G , Han SK , Herbison AE . 2009 ; Dendro-dendritic bundling and shared synapses between gonadotropin-releasing hormone neurons . *Proc Natl Acad Sci U S A* . 106 : 10835 - 10840
- 9 . Campbell RE , Han SK , Herbison AE . 2005 ; Biocytin filling of adult gonadotropin-releasing hormone neurons in situ reveals extensive, spiny, dendritic processes . *Endocrinology* . 146 : 1163 - 1169
- 10 . Barry J , Dubois MP , Poulain P . 1973 ; LRF producing cells of the mammalian hypothalamus. A fluorescent antibody study . *Z Zellforsch Mikrosk Anat* . 146 : 351 - 366
- 11 . Gross DS . 1976 ; Distribution of gonadotropin-releasing hormone in the mouse brain as revealed by immunohistochemistry . *Endocrinology* . 98 : 1408 - 1417
- 12 . Ciofi P , Garret M , Lapirot O , Lafon P , Loyens A , Prevot V , Levine JE . 2009 ; Brain-endocrine interactions: a microvascular route in the mediobasal hypothalamus . *Endocrinology* . 150 : 5509 - 5519
- 13 . Mullier A , Bouret SG , Prevot V , Dehouck B . 2010 ; Differential distribution of tight junction proteins suggests a role for tanycytes in blood-hypothalamus barrier regulation in the adult mouse brain . *J Comp Neurol* . 518 : 943 - 962
- 14 . Ferguson AV , Bains JS . 1996 ; Electrophysiology of the circumventricular organs . *Front Neuroendocrinol* . 17 : 440 - 475
- 15 . Broadwell RD , Brightman MW . 1976 ; Entry of peroxidase into neurons of the central and peripheral nervous systems from extracerebral and cerebral blood . *J Comp Neurol* . 166 : 257 - 283
- 16 . Rotsztein WH , Charli JL , Pattou E , Kordon C . 1977 ; Stimulation by dopamine of luteinizing hormone-releasing hormone (LHRH) release from the mediobasal hypothalamus in male rats . *Endocrinology* . 101 : 1475 - 1483
- 17 . Hoffman GE , Lee WS , Attardi B , Yann V , Fitzsimmons MD . 1990 ; Luteinizing hormone-releasing hormone neurons express c-fos antigen after steroid activation . *Endocrinology* . 126 : 1736 - 1741
- 18 . Lee WS , Smith MS , Hoffman GE . 1990 ; Luteinizing hormone-releasing hormone neurons express Fos protein during the proestrous surge of luteinizing hormone . *Proc Natl Acad Sci U S A* . 87 : 5163 - 5167
- 19 . Levin BE , Magnan C , Dunn-Meynell A , Le Foll C . 2011 ; Metabolic sensing and the brain: who, what, where, and how? . *Endocrinology* . 152 : 2552 - 2557
- 20 . Furman M , Wade GN . 2007 ; Animal models in the study of nutritional infertility . *Curr Opin Endocrinol Diabetes Obes* . 14 : 475 - 481
- 21 . Zhang C , Bosch MA , Levine JE , Ronnekleiv OK , Kelly MJ . 2007 ; Gonadotropin-releasing hormone neurons express K(ATP) channels that are regulated by estrogen and responsive to glucose and metabolic inhibition . *J Neurosci* . 27 : 10153 - 10164
- 22 . Roland AV , Moenter SM . 2011 ; Glucosensing by GnRH neurons: inhibition by androgens and involvement of AMP-activated protein kinase . *Mol Endocrinol* . 25 : 847 - 858
- 23 . Bucholtz DC , Vidwans NM , Herbosa CG , Schillo KK , Foster DL . 1996 ; Metabolic interfaces between growth and reproduction. V. Pulsatile luteinizing hormone secretion is dependent on glucose availability . *Endocrinology* . 137 : 601 - 607
- 24 . Nagatani S , Bucholtz DC , Murahashi K , Estacio MA , Tsukamura H , Foster DL , Maeda KI . 1996 ; Reduction of glucose availability suppresses pulsatile luteinizing hormone release in female and male rats . *Endocrinology* . 137 : 1166 - 1170