



HAL
open science

HIV-1 integrase SUMOylation and viral replication

Alessia Zamborlini, Audrey Coiffic, Guillaume Beauclair, Olivier Delelis, Joris Paris, Fabien Magne, Yashuiro Koh, Marie-Lou Giron, Joelle Tobaly-Tapiero, Stephane Emiliani, et al.

► **To cite this version:**

Alessia Zamborlini, Audrey Coiffic, Guillaume Beauclair, Olivier Delelis, Joris Paris, et al.. HIV-1 integrase SUMOylation and viral replication. *Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts*, Sep 2009, Montpellier, France. pp.I14, 10.1186/1742-4690-8-S2-O4 . inserm-00637229


HAL Id: inserm-00637229

<https://inserm.hal.science/inserm-00637229>

Submitted on 31 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ORAL PRESENTATION

Open Access

HIV-1 integrase SUMOylation and viral replication

Alessia Zamborlini^{1,2*}, Audrey Coiffic¹, Guillaume Beauclair¹, Olivier Delelis³, Joris Paris¹, Fabien Magne^{1,2}, Yashuiro Koh⁴, Marie-Lou Giron¹, Joelle Tobaly-Tapiero¹, Stephane Emiliani⁵, Alan Engelman⁴, Hugues de The¹, Ali Saib^{1,2}

From *Frontiers of Retrovirology* 2011
Amsterdam, The Netherlands. 3-5 October 2011

Background

HIV-1 integrase catalyzes the integration of the reverse transcribed viral cDNA into the cellular genome, a key event of retroviral replication that is targeted by novel anti-HIV therapeutic agents. Numerous studies have contributed to understand of the molecular basis of integrase catalytic functions. Besides integration, HIV-1 integrase participates in other steps of viral replication such as reverse transcription and/or uncoating, PIC nuclear import and virion morphology. However, the underlying mechanisms are still not fully elucidated. Cellular and viral factors assist integrase in performing its multiple activities. Moreover, post-translational modifications (i.e. acetylation, ubiquitination and phosphorylation), which represent a common, rapid and generally reversible mechanism for fine tuning of protein activities, have also been shown to regulate integrase functions.

Materials and methods and results

By *in vitro* SUMOylation assay and purification on Ni-NTA beads in denaturing conditions, we show that HIV-1 integrase is covalently modified by the three SUMO paralogues. By mutating SUMO-acceptor lysine residues within phylogenetically conserved SUMOylation consensus motifs identified *in silico*, we demonstrate that they represent major sites of SUMO conjugation. We introduced the same changes in the integrase sequence of a plasmid encoding the viral genome, which was used to generate mutant viral particles, and we compare their infectivity to that of WT HIV-1. We find that viruses harboring SUMOylation-defective integrase mutants are less infectious than the WT counterpart. Real-time PCR analysis of viral cDNA synthesis reveals that cells infected with mutant viruses display similar content of traverse

transcripts, but a lower number of integrated proviruses, than WT HIV-1-infected cells. This integration defect at the integration step is not due to impairment of integrase binding to LEDGF/p75, a key chromatin-tethering factor of HIV-1 pre-integration complex. Indeed, both WT and SUMOylation-defective integrase proteins coprecipitated with similar efficiency with LEDGF/p75. Finally, we establish that SUMOylation-defective integrase mutants retained WT catalytic activity as determined by Vpr-fusion protein complementation assay.

Conclusions

Here, we show that HIV-1 IN is modified by SUMO proteins and that phylogenetically conserved SUMOylation consensus motifs represent major SUMO-acceptor sites. Viruses harboring SUMOylation-site IN mutants displayed a replication defect that was mapped during the early stages of infection, before integration but after reverse transcription. Since SUMOylation-defective IN mutants retained WT catalytic activity as well as LEDGF-binding, we hypothesize that SUMOylation might regulate the affinity of IN for co-factors, contributing to efficient HIV-1 replication.

Author details

¹CNRS UMR7212, INSERM U944, Université Paris7 Diderot, Institut Universitaire d'Hématologie, Paris, 75010, France. ²Conservatoire National des Arts et Métiers, Department CASER, Paris, 75003, France. ³LBPA, CNRS UMR8113, Ecole Normale Supérieure, Cachin, 94235, France. ⁴Dana-Farber Cancer Institute, Department of Cancer Immunology & AIDS, Boston, MA, 02115, USA. ⁵INSERM U1016, CNRS JMR8104, Université Paris Descartes, Cochin Institut, Paris, 75014, France.

Published: 3 October 2011

doi:10.1186/1742-4690-8-S2-O4

Cite this article as: Zamborlini *et al.*: HIV-1 integrase SUMOylation and viral replication. *Retrovirology* 2011 **8**(Suppl 2):O4.

¹CNRS UMR7212, INSERM U944, Université Paris7 Diderot, Institut Universitaire d'Hématologie, Paris, 75010, France
Full list of author information is available at the end of the article