



**HAL**  
open science

## **Epicardial fat volume is associated with coronary endothelium-dependent vasomotor response in healthy subjects**

Benedicte Gaborit, Pierre Julien Moro, Antonin Flavian, Frank Kober, Alexis Jacquier, Jacques Quilici, Thomas Cuisset, Umberto Simeoni, Patrick J Cozzone, Monique Bernard, et al.

► **To cite this version:**

Benedicte Gaborit, Pierre Julien Moro, Antonin Flavian, Frank Kober, Alexis Jacquier, et al.. Epicardial fat volume is associated with coronary endothelium-dependent vasomotor response in healthy subjects. *Journal of Cardiovascular Magnetic Resonance*, 2011, 13 (Suppl 1), pp.P71. inserm-00654376

**HAL Id: inserm-00654376**

**<https://inserm.hal.science/inserm-00654376v1>**

Submitted on 21 Dec 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

# Epicardial fat volume is associated with coronary endothelium-dependent vasomotor response in healthy subjects

Bénédicte Gaborit<sup>1\*</sup>, Pierre Julien Moro<sup>2</sup>, Antonin Flavian<sup>3</sup>, Frank Kober<sup>3</sup>, Alexis Jacquier<sup>3</sup>, Jacques Quilici<sup>4</sup>, Thomas Cuisset<sup>5</sup>, Umberto Simeoni<sup>6</sup>, Patrick Cozzone<sup>3</sup>, Monique Bernard<sup>3</sup>, Anne Dutour<sup>1</sup>

From 2011 SCMR/Euro CMR Joint Scientific Sessions  
Nice, France. 3-6 February 2011

## Introduction

Epicardial fat ( $E_{fat}$ ) is an active ectopic fat depot, directly surrounding coronary arteries, and secreting high level of inflammatory adipokines; its development has been associated with coronary atherosclerosis. We investigated the relationship between  $E_{fat}$  and endothelium dependent vasoreactivity of the coronary microcirculation.

## Methods

Myocardial blood flow (MBF) was determined by measuring coronary sinus flow with velocity-encoded cine magnetic resonance imaging at 3 teslas. We measured MBF at baseline and in response to sympathetic stimulation by cold pressor testing (CPT) in 17 healthy volunteers with normal left ventricular function (age  $24 \pm 6$  years, BMI =  $21.1 \pm 2.6$  kg/m<sup>2</sup>).  $E_{fat}$  volume was volumetrically assessed by manual delineation on short-axis views. CPT was applied by immersing one foot in ice water for 4 minutes.

## Results

A significant increase in MBF was observed:  $1.18 \pm 0.58$  vs  $0.84 \pm 0.47$  mL.min<sup>-1</sup>.g<sup>-1</sup>, CPT vs rest,  $p=0.002$ . Mean relative MBF increase ( $\Delta$ MBF) was  $50 \pm 47\%$ . Mean  $E_{fat}$  volume was  $82 \pm 31$  mL and varied from 43 to 131 mL; mean LV mass and Left ventricular ejection fraction were  $104 \pm 31$  g and  $64 \pm 5\%$ , respectively. CPT significantly increased heart rate (HR) by  $28 \pm 13\%$ , systolic blood pressure (BP) by  $17 \pm 13\%$ , diastolic BP by  $23 \pm 19\%$  and rate-pressure

product by  $52 \pm 25\%$ ,  $p < 0.01$ , indicating an increase in myocardial work load. The increase in HR, reflecting sympathetic stimulation, was not influenced by sex, age or  $E_{fat}$  volume. CPT induced a decrease in coronary vascular resistance ( $150 \pm 93$  vs  $114 \pm 44$  mmHg.mL<sup>-1</sup>.min.g) by trend ( $p=0.08$ ). Interestingly, we found a significant negative correlation between  $E_{fat}$  volume and  $\Delta$ MBF ( $r=-0.51$ ,  $p=0.03$ ), which remained significant after adjusting for age and sex.  $\Delta$ MBF was not associated with waist circumference, BMI, CRP, lipid or glycemic parameters.

## Conclusion

The increase in  $E_{fat}$  is associated with a decrease in endothelium dependent vasoreactivity response, suggesting that  $E_{fat}$  could early influence endothelial function.

## Author details

<sup>1</sup>INSERM U626, Marseille, F-13385 France, Centre de Résonance Magnétique Biologique et Médicale (CRMBM), CNRS UMR 6612, Department of Endocrinology, Metabolic Diseases and Nutrition, CHU Nord, Marseille, France, Marseille, France. <sup>2</sup>Centre de Résonance Magnétique Biologique et Médicale, CNRS UMR N°6612, Marseille, France. <sup>3</sup>Centre de Résonance Magnétique Biologique et Médicale (CRMBM), CNRS UMR 6612, Marseille, France. <sup>4</sup>Department of Cardiology, CHU Timone, Marseille, France, Marseille, France. <sup>5</sup>INSERM U626, Marseille, F-13385 France, Department of Cardiology, CHU Timone, Marseille, France, Marseille, France. <sup>6</sup>Department of Neonatology, Children and Parents Pole, CHU Timone, Marseille, France, Marseille, France.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P71

**Cite this article as:** Gaborit et al.: Epicardial fat volume is associated with coronary endothelium-dependent vasomotor response in healthy subjects. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1): P71.

<sup>1</sup>INSERM U626, Marseille, F-13385 France, Centre de Résonance Magnétique Biologique et Médicale (CRMBM), CNRS UMR 6612, Department of Endocrinology, Metabolic Diseases and Nutrition, CHU Nord, Marseille, France, Marseille, France

Full list of author information is available at the end of the article