



Upstream ORF affects MYCN translation depending on exon 1b alternative splicing.

Roger Besançon, Sandrine Valsesia-Wittmann, Clara Locher, Céline Delloye-Bourgeois, Lydie Furhman, Giovanni Tutrone, Christophe Bertrand, Anne-Catherine Jallas, Elisabeth Garin, Alain Puisieux

► To cite this version:

Roger Besançon, Sandrine Valsesia-Wittmann, Clara Locher, Céline Delloye-Bourgeois, Lydie Furhman, et al.. Upstream ORF affects MYCN translation depending on exon 1b alternative splicing.. BMC Cancer, 2009, 9 (1), pp.445. 10.1186/1471-2407-9-445 . inserm-00663547

HAL Id: inserm-00663547

<https://inserm.hal.science/inserm-00663547>

Submitted on 27 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research article

Open Access

Upstream ORF affects MYCN translation depending on exon 1b alternative splicing

Roger Besançon^{*1,2}, Sandrine Valsesia-Wittmann³, Clara Locher¹, Céline Delloye-Bourgeois¹, Lydie Furhman¹, Giovanni Tutrone¹, Christophe Bertrand¹, Anne-Catherine Jallas³, Elisabeth Garin³ and Alain Puisieux^{1,2,3}

Address: ¹INSERM UMR590, 28 rue Laënnec, Lyon, France, ²Université de Lyon, université Lyon 1, Institut des Sciences Pharmaceutiques et Biologiques, 8 avenue Rockefeller, Lyon, 69008, France and ³Centre Léon Bérard, Laboratoire de Recherche Translationnelle, 28 rue Laënnec, Lyon, France

Email: Roger Besançon^{*} - roger.besancon@recherche.univ-lyon1.fr; Sandrine Valsesia-Wittmann - WITTMANN@lyon.fnclcc.fr; Clara Locher - Clara.LOCHER@igr.fr; Céline Delloye-Bourgeois - delloye@lyon.fnclcc.fr; Lydie Furhman - lydiefurhmann@hotmail.com; Giovanni Tutrone - giovannitutrone@hotmail.com; Christophe Bertrand - bertrandchristophe@neuf.fr; Anne-Catherine Jallas - jallas@lyon.fnclcc.fr; Elisabeth Garin - garin@lyon.fnclcc.fr; Alain Puisieux - puisieux@lyon.fnclcc.fr

^{*} Corresponding author

Published: 17 December 2009

Received: 7 July 2009

BMC Cancer 2009, **9**:445 doi:10.1186/1471-2407-9-445

Accepted: 17 December 2009

This article is available from: <http://www.biomedcentral.com/1471-2407/9/445>

© 2009 Besançon et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The MYCN gene is transcribed into two major mRNAs: one full-length (MYCN) and one exon 1b-spliced (MYCN^{Δ1b}) mRNA. But nothing is known about their respective ability to translate the MYCN protein.

Methods: Plasmids were prepared to enable translation from the upstream (uORF) and major ORF of the two MYCN transcripts. Translation was studied after transfection in neuroblastoma SH-EP cell line. Impact of the upstream AUG on translation was evaluated after directed mutagenesis. Functional study with the two MYCN mRNAs was conducted by a cell viability assay. Existence of a new protein encoded by the MYCN^{Δ1b} uORF was explored by designing a rabbit polyclonal antibody against a specific epitope of this protein.

Results: Both are translated, but higher levels of protein were seen with MYCN^{Δ1b} mRNA. An upstream ORF was shown to have positive cis-regulatory activity on translation from MYCN but not from MYCN^{Δ1b} mRNA. In transfected SH-EP neuroblastoma cells, high MYCN dosage obtained with MYCN^{Δ1b} mRNA translation induces an antiapoptotic effect after serum deprivation that was not observed with low MYCN expression obtained with MYCN mRNA. Here, we showed that MYCNOT: MYCN Overlap Transcript, a new protein of unknown function is translated from the upstream AUG of MYCN^{Δ1b} mRNA.

Conclusions: Existence of upstream ORF in MYCN transcripts leads to a new level of MYCN regulation. The resulting MYCN dosage has a weak but significant anti-apoptotic activity after intrinsic apoptosis induction.

Background

The *MYCN* gene, located in 2p23-24 [1] has been demonstrated to be composed of three exons (Fig. 1): exon 1a/b, exon 2 and exon 3, and the coding sequence for MYCN has been mapped to exons 2 and 3 [2]. Several transcripts have been described; one full length: *MYCN* mRNA and two alternatively spliced mRNAs; one containing exons 1a, 2 and 3: *MYCN^{Δ1b}* mRNA [2,3] and the other containing exons 1a and 3: *MYCN^{Δ1b,2}* mRNA [4]. Only *MYCN* and *MYCN^{Δ1b}* transcripts might translate the MYCN protein. It was previously hypothesized but neither proven nor explored that an upstream AUG (uAUG) located in +1894 (according to GenBank: [Y00664](#)) of *MYCN^{Δ1b}* mRNA might translate a protein (GenBank: [ΔAG40001.1](#)) [5]. Moreover, as proposed by van Bokhoven *et al*, *MYCN^{Δ1b,2}* mRNA could be able to initiate translation of a new protein (ΔMYCN) from the same uAUG [4].

The existence of upstream open reading frames (uORFs) has been described in detail for many genes, and these uORFs could represent as much as half of the whole transcriptome [6,7]. Most studies have shown a negative cis-regulatory function of the major ORF [6-8].

MYCN, a basic helix-loop-helix transcription factor that belongs to the MYC family, was initially identified as a gene amplified in neuroblastoma, the most frequent pae-

diatric extra-cranial solid tumour [9]. Numerous experiments have demonstrated its importance during ontogenesis. In the developing mouse embryo, MYCN is required for normal organogenesis of the heart, neural tube, spinal ganglia, visceral arches, liver, stomach, limb buds and eyes [10]. More precisely, MYCN is essential to maintain a population of undifferentiated and proliferating progenitor cells in the brain [11] and the distal lung epithelium [12] of mouse embryos. MYCN is also able, in chick embryos, to initiate the ventral migration of cells from the neural crest to the sympathetic ganglia and to induce their differentiation into neurons [13].

The overexpression of MYCN in the presence of deregulated H-ras has been shown to contribute to the neoplastic transformation of rat embryo cells, suggesting an oncogenic role of MYCN [14]. This oncogenic role was later confirmed using MYCN-transgenic mice: targeted expression of MYCN to neural crest-derived cells under the control of the *Tyrosine Hydroxylase* promoter leads to the development of neuroblastoma-like tumours in all homozygous mice [15].

The amplification and overexpression of MYCN primarily found in 25% of neuroblastomas are associated with advanced tumour stage, tumour progression and poor outcome [16]. MYCN can also be overexpressed in other




Figure 1

The MYCN oncogene and encoded proteins. a) Annotated sequence of MYCN according to Genbank [Y00664](#). **b)** Alternative splicing and putative proteins encoded by MYCN. **c)** MYCNOT sequence. Bold aminoacids: shared homology with the putative ΔMYCN [4]. Underlined aminoacids: selected epitope for polyclonal antibody production in rabbit.

tumours including medulloblastoma, retinoblastoma, small cell lung cancer, glioblastoma and other embryonal tumours [17] suggesting the implication of MYCN dosage in cancer development and aggressive expressivity. In single copy MYCN (SCN) neuroblastoma cell lines, MYCN has also been demonstrated to enhance proliferation after bFGF administration [18]. Contrariwise, the protein exhibits pro-apoptotic properties in particular conditions such as drug-triggered apoptosis [19,20] or induction of the death receptor machinery by TRAIL [21].

Thus, MYCN protein exhibits dual properties in proliferation and apoptosis, both in physiological and in pathological conditions.

In the present work, we evidenced a link between MYCN transcription and the level of MYCN translation. *In vitro* translation of the two coding mRNAs led to different levels of MYCN expression, with MYCN $\Delta 1b$ mRNA being the most efficient. Their translation was differently regulated by an uORF located in the long 5' untranslated region of exon 1a/b. Moreover, we observed that the uORF of MYCN mRNA was able to up-regulate MYCN translation, whereas the uORF of MYCN $\Delta 1b$ mRNA was not. Our results demonstrated a MYCN dosage effect in SH-EP cells in which high amounts of MYCN were anti-apoptotic after serum deprivation, compared to low levels or absence of MYCN. Finally, the uORF of MYCN $\Delta 1b$ mRNA directed the translation of a new protein of unknown function, MYCNOT.

Methods

Cell culture

SH-EP cell line (an epithelial substrate-adherent Schwann-like clone issued from SK-N-SH neuroblastoma cell line[22] that do not express MYCN protein was cultured in RMPI 1640 (Sigma) supplemented with 10% foetal calf serum (GIBCO), penicillin G (200 IU/mL; GIBCO), streptomycin sulphate (200 µg/mL; GIBCO) and L-glutamine (2 mM; GIBCO) at 37°C in an atmosphere containing 5% CO₂.

Plasmids preparation

Plasmid preparations were based on the Invitrogen Gateway® technology. Inserts were prepared from human foetal brain mRNAs (Clontech) with a forward primer specific to the very beginning of exon 1a/b at +1865 (according to Genbank Y00664) flanked with an attB1 sequence. Depending on the desired construct, three reverse primers extended with an attB2 sequence were designed. The first one, for *p-uORF/MYCN* was complementary to the sequence of the penultimate nucleotides of the first stop codon of the MYCN mRNA in frame with the uAUG codon located at +1894. The second one, for *p-uORF/MYCN $\Delta 1b$* , was complementary to the sequence of the penultimate nucleotides of the first stop codon of

MYCN $\Delta 1b$ mRNA in frame with the same uAUG codon. The third reverse primer, for *p-MYCN* and *p-MYCN $\Delta 1b$* , was complementary to the sequence of the penultimate nucleotides of the MYCN stop codon in exon 3. DNAs amplified from MYCN and MYCN $\Delta 1b$ mRNAs were inserted into *pDEST40* according to Invitrogen Gateway® protocols with a C-terminal V5 epitope tag for visualizing protein translation.

The plasmids *p-MYCN^{mut}* and *p-MYCN $\Delta 1b$, ^{mut}* were obtained respectively from *p-MYCN* and *p-MYCN $\Delta 1b$* by mutation of the upstream AUG located in +1894 into an AUC using the Quickchange® Site-Directed Mutagenesis kit (Stratagene).

Cell transfection

For translational studies, three independent experiments were conducted in duplicate: SH-EP cells were sowed at a density of 200000 cells in 6-wells plates and transfected the next day with 3.6 µg of purified MYCN-derived plasmids or control *pcDNA/GW-40/lacZ* (Invitrogen), 0.4 µg *p-EGFP* (Clontech) and 8 µL Lipofectamine 2000 (Invitrogen). Cells were harvested 24 hours after transfection.

For apoptosis experiments, three independent experiments were conducted in quadruplicate: SH-EP cells were sowed at a density of 200000 cells in 6-wells plates and cultured in RMPI 1640 supplemented with 0.5% foetal calf serum, penicillin G (200 IU/mL), streptomycin sulphate (200 µg/mL) and L-glutamine (2 mM) for 48 hours at 37°C in an atmosphere containing 5% CO₂, then transfected with 3.6 µg of *p-MYCN*, *p-MYCN $\Delta 1b$* or *pcDNA/GW-40/lacZ*, 0.4 µg *p-EGFP* and 8 µL Lipofectamine 2000. After transfection, cells were cultured for an additional day in the same medium supplemented with 10% foetal calf serum.

Western blotting

Proteins (40 µg) from cell lysates or from a human foetal and adult brain (protein Medley™; BD Biosciences) were separated by SDS-PAGE and transferred to 0.2 µm-nitrocellulose membranes (Biorad). Protein levels were determined using:

- a mouse anti-V5 epitope monoclonal antibody (1:5000, Invitrogen),
- a mouse anti-human β-ACTIN monoclonal antibody (1:200000, Sigma),
- a rabbit anti-NGAERSPQSPAGRRA [anti-MYCNOT] peptide polyclonal antibody (1:100, Eurogentec), whose specificity was checked by manufacturer
- a mouse anti-human MYCN monoclonal antibody, clone B8.4.B (1:1000, BD Pharmingen).

Primary antibodies were detected using goat anti-mouse (1:10000, Sigma) or anti-rabbit (1:3000, Sigma) horse-radish peroxidase-conjugated secondary antibodies, and visualized using an enhanced chemiluminescence HRP substrate (Millipore).

Cell viability assay

Cell viability was measured with Uptibblue viable cell counting reagent (Uptima) before serum deprivation and one day after transfection: cells were incubated for two hours after addition of 10% Uptibblue, fluorescence was measured with 530 nm excitation wavelength and 590 nm emission wavelength.

A one-sided student-t test was performed on the serum deprivation-induced apoptosis calculated as the fluorescence difference between the two measures.

Results

MYCN and MYCN^{1b} mRNAs are differently translated

Since alternative splicing of exon 1b affects a potential upstream open reading frame (uORF), we speculated that it could influence the translation of MYCN protein. Vectors *p-MYCN* and *p-MYCN^{Δ1b}* were prepared to allow the translation of MYCN respectively from these two mRNAs. In SH-EP neuroblastoma cells, transfection assays followed by Western blot showed no MYCN expression in presence of the control plasmid (*pcDNA/GW40/lacZ*), weak but detectable translation from *p-MYCN*, whereas a very high amount of protein was obtained with *p-MYCN^{Δ1b}* (Fig. 2a). The observed differences are not a simple consequence of experimental variations in either transfection efficacy, because cotransfection of *p-EGFP* induced no difference in the number of GFP-expressing cells between experiments (data not shown), or RNA transcription or stability, because semi quantitative RT-PCR analysis showed no difference in the levels of MYCN and MYCN^{Δ1b} transcripts brought by plasmids (data not shown). Upstream ORF is usually known for *cis*-acting inhibitory effect on translation of the major protein. To test this hypothesis, the upstream AUG was mutated to AUC to produce *p-MYCN^{mut}* and *p-MYCN^{Δ1b, mut}* vectors. MYCN levels in SH-EP cells clearly diminished when it was translated from MYCN transcripts, but not from MYCN^{Δ1b} transcripts (Fig. 2b).

High MYCN dosage enhances cell viability

To explore the effect of MYCN dosage, SH-EP cells were subjected to apoptosis by 48-hour serum deprivation, then transfected with either control *pcDNA/GW40/lacZ*, *p-MYCN*, or *p-MYCN^{Δ1b}* plasmids. As shown in Table 1, when SH-EP cells did not express (*pcDNA/GW-40/lacZ* transfected cells) or expressed low amount (*p-MYCN* transfected cells) of MYCN protein, serum deprivation induced similar cell loss (respectively 35.56% +/- 6.68% and 34.94 +/- 7.19%; *p* = 0.60). When SH-EP cells




Figure 2

Exon 1b-splicing, uAUG and MYCN translation. **a)** Western blot analysis of 40 μg SH-EP neuroblastoma cells transfected with either control *pcDNA/GW40/lacZ* (lane 1), *p-MYCN* (lane 2) or *p-MYCN^{Δ1b}* (lane 3). Results obtained with -upper panel: a monoclonal antibody against MYCN (1:1000) and -lower panel: a monoclonal antibody against β-Actin (1:10000). **b)** Western blot analysis of 40 μg SH-EP cells transfected with either *p-MYCN* (lane 1), *p-MYCN^{mut}* (lane 2), *p-MYCN^{Δ1b}* (lane 3) or *p-MYCN^{Δ1b, mut}* (lane 4). Results obtained with -upper panel: the anti-MYCN antibody (1:1000) and -lower panel: the anti β-Actin antibody (1:10000).

expressed high levels (*p-MYCN^{Δ1b}* transfected cells) of MYCN protein, apoptosis (26.72% +/- 8.09%) was significantly (*p* = 0.01 and *p* = 0.02 respectively compared to previous conditions) reduced by one third.

MYCNOT is translated from a upstream AUG of MYCN^{1b} mRNA

We speculated that uAUG, located in +1894 (according to GenBank: [Y00664](#)), may initiate the translation and we assessed the importance of alternative splicing of exon 1b. In SH-EP cells, the transfection of *p-uORF/MYCN^{Δ1b}*,

Table 1: MYCN dose-dependant anti-apoptotic effect after serum deprivation.

Transfected plasmids	Cell viability (%)
<i>pcDNA/GW-40/lacZ</i>	64.44 +/- 6.68
<i>pMYCN</i>	65.06 +/- 7.19 ^a
<i>pMYCN^{Δ1b}</i>	73.27 +/- 8.09 ^{b,c}

MYCN dose-dependant enhanced SH-EP viability after serum deprivation measured with Uptibblue viable cell counting reagent (Uptima). ^a *p* = 0.60 compared with *pcDNA/GW-40/lacZ*; ^b *p* = 0.01 compared with *pcDNA/GW-40/lacZ*; ^c *p* = 0.02 compared with *pMYCN*

which contains the uORF of *MYCN*^{Δ1b} mRNA, allowed the production of a V5 epitope-tagged protein of approximately 15 kDa, whereas the transfection of *p-uORF/Myc*, which contains the uORF of the full length *MYCN* transcript, did not (Fig. 3a). This 11.8 kDa new protein was named MYCNOT: *MYCN* Overlap Transcript.

To validate the existence of endogenous MYCNOT, we performed a Western blot analysis using a specifically designed polyclonal antibody raised in rabbit against a N-terminal epitope (NGAERSPQSPAGRRA) [anti-MYCNOT] of the hypothetical protein. The designed MYCNOT antibody was able to detect in transfected SH-EP cells with *p-uORF/MYCN*^{Δ1b} the same protein detected with the V5 epitope antibody (Fig. 3b). The anti-MYCNOT antibody was able to detect a ~12 kDa protein in foetal but not in adult brain (Fig. 3c), thus confirming the existence of an endogenous MYCNOT protein. An analysis of MYCNOT

sequence with BLAST and PROSITE software did not show any sequence or motif homology.

Discussion

Because upstream ORF are known to influence protein translation [23], we investigated the *MYCN* translation from the full length *MYCN* mRNA and the exon 1b-spliced *MYCN*^{Δ1b} mRNA. Only these two mRNAs are able to translate the MYCN protein. More recently, in 2005, van Bokhoven *et al* described a new exon 1a/2-spliced mRNA able to translate a new protein named ΔMYCN [4]. In SH-EP cells, we demonstrated that both *MYCN* and *MYCN*^{Δ1b} were able to translate the MYCN protein, but *MYCN* translation from *MYCN*^{Δ1b} mRNA was much more efficient. This difference may not be attributed to a differential stability of the two transcripts because they have the same half-life of approximately 15 minutes [2]. But, that could be due to differences in the regulating activities of uORFs in the two mRNAs. Because, the coding AUG is several hundreds nucleotides after the stop codon of the uORF in *MYCN* mRNA compared to the overlapping uORF in *MYCN*^{Δ1b} transcript, we should have expected a higher translation of MYCN protein from full length mRNA due to reinitiation [6,23], but as already demonstrated, this mRNA contains an Internal Ribosome Entry Site (IRES) that could explain the higher initiation of translation at the major AUG in *MYCN*^{Δ1b} mRNA [5,24,25].

Upstream ORFs are usually known for their *cis*-acting inhibitory activity on the translation of the major protein [7,26-30]. To test this hypothesis, uAUG was mutated into AUC. This mutation had no effect on MYCN translation from *MYCN*^{Δ1b} mRNA but significantly impacted the translation from *MYCN* mRNA. Thus, *MYCN* uORFs have different *cis*-regulatory activities in MYCN translation depending on the alternative splicing of exon 1b. In our conditions, the uORF of the *MYCN*^{Δ1b} mRNA did not affect MYCN translation. Alternatively, the uORF transcribed from full length *MYCN* mRNA has *cis*-enhancing activity on MYCN translation. To our knowledge, only one team has shown that an uORF is able to augment the translation of the HIV-1 Env protein from the HIV-1 mRNA [30]. Taken together, our results suggest the existence of a new level of MYCN regulation. One hypothesis should be further explored: in cap-dependent conditions, low-level MYCN would be produced from the full length mRNA and this translation would be sustained by the activity of the uORF whereas in IRES-dependent conditions, high-level MYCN would be produced from exon 1b-spliced mRNA, independently of the uORF.

Numerous experiments have demonstrated important ontogenic [31] and oncogenic [14] roles. The amplification and overexpression of *MYCN* primarily found in 25% of neuroblastomas [16] and in other tumours




Figure 3

Translation of uORF and existence of MYCNOT. a) SH-EP cells were transfected with either *p-uORF/MYCN* (lane 1) or *p-uORF/MYCN*^{Δ1b} (lane 2). Western blot was conducted on 40 μg proteins, 48 hours after transfection with -upper panel: a monoclonal antibody against the V5 epitope (1:300) and -lower panel: the anti β-Actin antibody (1:10000). L: protein ladder; **b)** SH-EP cells were transfected with *p-uORF/MYCN*^{Δ1b}. Western blot was conducted on 40 μg protein, 48 hours after transfection with -left panel: a monoclonal antibody against the V5 epitope (1:300) and -right panel: a polyclonal rabbit anti-NGAERSPQSPAGRRA peptide [anti-MYCNOT] polyclonal antibody (1:100) L: protein ladder; **c)** Western blot analysis of 40 μg protein extracts from adult (lane 1) and foetal (lane 2) brain. Results obtained with -upper panel: the anti-MYCNOT antibody (1:100) and -lower panel: the anti β-Actin antibody (1:10000).

including medulloblastoma, retinoblastoma, small cell lung cancer, glioblastoma and other embryonal tumours [17] suggest the implication of MYCN dosage in cancer development and aggressive expressivity.

Our results showed that, in transfected SH-EP neuroblastoma cells, high MYCN dosage obtained with MYCN^{Δ1b} mRNA translation induces a weak but significant antiapoptotic effect after serum deprivation that was not observed with low MYCN expression obtained with MYCN mRNA. Our results do not fit the accepted proapoptotic properties of MYCN: using a MYCN-inducible SH-EP Tet21/N cell line, it has been demonstrated that MYCN overexpression enhances either doxorubicin- or platinum-induced apoptosis [19]. Using a similar approach, Cui et al have shown that the transfection of SH-EP cells with a *pBabe-hygro/MYCN* vector sensitises the cells to TRAIL-triggered apoptosis [21]. But, as we did, Jasty et al have reported a significant increase of cell viability with high MYCN expression after 6 days of serum deprivation [32]. These different results and ours suggest that MYCN overexpression in SH-EP cells might have different effects depending on the existence of an intrinsic (serum deprivation) or extrinsic (TRAIL or drug-induced apoptosis) pathway-mediated apoptosis. Thus, MYCN dosage, depending on translation of the alternatively exon 1b splicing mRNA, enhances cell viability after intrinsic pathway-mediated apoptosis.

As previously proposed, the uAUG codon located in exon 1a may initiate the translation of ΔMYCN, a new isoform of the MYCN protein [4]. It has also been hypothesized, but not proven at this time, that this uAUG is able to translate a protein (GenBank: [AAG40001.1](#)) [5]. We now demonstrated the existence of this 11.8 kDa protein and named it MYCNOT: MYCN Overlap Transcript. Even if uORFs appear to apply to as much as 50% of the transcriptome [6,7,23], only few papers have reported their translation in *in-vitro* experiments using chimeric tag-fusion proteins [23]. This is particularly true for the MYC gene, to which MYCN is related, which presents several uORFs: one uORF in exon 1 conducts to the translation of a protein of unknown function called MYCHEX1 [33]. To our knowledge, only one team, using two-dimensional chromatography and mass spectrometry analysis, has been able to prove the existence of eight proteins encoded by uORFs in human K562 and HEK293 cell lines [34]. Moreover, these small proteins (< 20 kDa) were not subject to rapid proteasome degradation suggesting functional activity [34]. The MYCNOT protein, which has no homology with MYCN, other known proteins or protein motifs, shares exon 1a-encoded amino acids with the putative ΔMYCN previously described [4]. Because MYCNOT is translated in foetal but not adult brain, we may hypothesize that it plays a role during neural development. Our

results need further analysis to determine the functions of MYCNOT and might provide new insights into MYCN regulation.

Conclusion

The MYCN gene is transcribed into two major mRNAs which are differently translated. Higher levels of protein were seen with MYCN^{Δ1b} mRNA. Existence of an upstream ORF was shown to have positive cis-regulatory activity on translation from MYCN but not from MYCN^{Δ1b} mRNA. The high MYCN dosage obtained with MYCN^{Δ1b} mRNA translation induces, in transfected SH-EP neuroblastoma cells, a weak but significant antiapoptotic effect after serum deprivation that was not observed with low MYCN expression.

Here, we show the translation of a new protein of unknown function from the upstream AUG of MYCN^{Δ1b} mRNA.

In conclusion, further studies are now needed to explore the exact relationship between alternative splicing of exon 1b, MYCN dosage, and the mechanism, either extrinsic or intrinsic, underlying apoptosis.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RB carried out the design of the study and manuscript writing and participated in plasmids preparation, cell culture and Western blotting experiments. SWV participated to design of the study and manuscript writing. CL participated to cell culture and Western blotting experiments. CDB participated to plasmids preparation, cell culture and Western blotting experiments. GT participated to cell culture and Western blotting experiments. ACJ participated to cell culture and Western blotting experiments. EG participated to cell culture and Western blotting experiments. AP participated to design of the study and manuscript writing. All authors read and approved the final manuscript."

Acknowledgements

The authors are grateful to Dr. A. Denuzière for statistical assistance and to Mrs M.-D. Reynaud for editorial assistance. This work was supported by INCa Projet Libre 2006-2008 PL 06-76.

References

- Schwab M, Varmus HE, Bishop JM, Grzeschik KH, Naylor SL, Sakaguchi AY, et al.: **Chromosome localization in normal human cells and neuroblastomas of a gene related to c-myc.** *Nature* 1984, **308**:288-291.
- Stanton LW, Bishop JM: **Alternative processing of RNA transcribed from NMYC.** *Mol Cell Biol* 1987, **7**:4266-4272.
- Ibson JM, Rabbitts PH: **Sequence of a germ-line N-myc gene and amplification as a mechanism of activation.** *Oncogene* 1988, **2**:399-402.

4. van Bokhoven H, Celli J, van Reeuwijk J, Rinne T, Glaudemans B, van Beusekom E, et al.: **MYCN haploinsufficiency is associated with reduced brain size and intestinal atresias in Feingold syndrome.** *Nat Genet* 2005, **37**:465-467.
5. Jopling CL, Willis AE: **N-myc translation is initiated via an internal ribosome entry segment that displays enhanced activity in neuronal cells.** *Oncogene* 2001, **20**:2664-2670.
6. Kochetov AV, Ahmad S, Ivanisenko V, Volkova OA, Kolchanov NA, Sarai A: **uORFs, reinitiation and alternative translation start sites in human mRNAs.** *FEBS Lett* 2008, **582**:1293-1297.
7. Calvo SE, Pagliarini DJ, Mootha VK: **Upstream open reading frames cause widespread reduction of protein expression and are polymorphic among humans.** *Proc Natl Acad Sci USA* 2009, **106**:7507-7512.
8. Landry JR, Mager DL, Wilhelm BT: **Complex controls: the role of alternative promoters in mammalian genomes.** *Trends Genet* 2003, **19**:640-648.
9. Schwab M, Alitalo K, Klempnauer KH, Varmus HE, Bishop JM, Gilbert F, et al.: **Amplified DNA with limited homology to myc cellular oncogene is shared by human neuroblastoma cell lines and a neuroblastoma tumour.** *Nature* 1983, **305**:245-248.
10. Sawai S, Shimono A, Wakamatsu Y, Palmes C, Hanaoka K, Kondoh H: **Defects of embryonic organogenesis resulting from targeted disruption of the N-myc gene in the mouse.** *Development* 1993, **117**:1445-1455.
11. Knoepfler PS, Cheng PF, Eisenman RN: **N-myc is essential during neurogenesis for the rapid expansion of progenitor cell populations and the inhibition of neuronal differentiation.** *Genes Dev* 2002, **16**:2699-2712.
12. Okubo T, Knoepfler PS, Eisenman RN, Hogan BL: **Nmyc plays an essential role during lung development as a dosage-sensitive regulator of progenitor cell proliferation and differentiation.** *Development* 2005, **132**:1363-1374.
13. Wakamatsu Y, Watanabe Y, Nakamura H, Kondoh H: **Regulation of the neural crest cell fate by N-myc: promotion of ventral migration and neuronal differentiation.** *Development* 1997, **124**:1953-1962.
14. Schwab M, Varmus HE, Bishop JM: **Human N-myc gene contributes to neoplastic transformation of mammalian cells in culture.** *Nature* 1985, **316**:160-162.
15. Weiss WA, Aldape K, Mohapatra G, Feuerstein BG, Bishop JM: **Targeted expression of MYCN causes neuroblastoma in transgenic mice.** *EMBO J* 1997, **16**:2985-2995.
16. Seeger RC, Brodeur GM, Sather H, Dalton A, Siegel SE, Wong KY, et al.: **Association of multiple copies of the N-myc oncogene with rapid progression of neuroblastomas.** *N Engl J Med* 1985, **313**:1111-1116.
17. Pession A, Tonelli R: **The MYCN oncogene as a specific and selective drug target for peripheral and central nervous system tumors.** *Curr Cancer Drug Targets* 2005, **5**:273-283.
18. Schweigerer L, Ledoux D, Fleischmann G, Barritault D: **Enhanced MYCN oncogene expression in human neuroblastoma cells is associated with altered FGF receptor expression and cellular growth response to basic FGF.** *Biochem Biophys Res Commun* 1991, **179**:1449-1454.
19. Fulda S, Lutz W, Schwab M, Debatin KM: **MycN sensitizes neuroblastoma cells for drug-triggered apoptosis.** *Med Pediatr Oncol* 2000, **35**:582-584.
20. Paffhausen T, Schwab M, Westermann F: **Targeted MYCN expression affects cytotoxic potential of chemotherapeutic drugs in neuroblastoma cells.** *Cancer Lett* 2007, **250**:17-24.
21. Cui H, Li T, Ding HF: **Linking of N-Myc to death receptor machinery in neuroblastoma cells.** *J Biol Chem* 2005, **280**:9474-9481.
22. Ross RA, Spengler BA, Biedler JL: **Coordinate morphological and biochemical interconversion of human neuroblastoma cells.** *J Natl Cancer Inst* 1983, **71**:741-747.
23. Iacono M, Mignone F, Pesole G: **uAUG and uORFs in human and rodent 5'untranslated mRNAs.** *Gene* 2005, **349**:97-105.
24. Cobbold LC, Spriggs KA, Haines SJ, Dobbins HC, Hayes C, de Moor CH, et al.: **Identification of internal ribosome entry segment (IRES)-trans-acting factors for the Myc family of IRESs.** *Mol Cell Biol* 2008, **28**:40-49.
25. Spriggs KA, Cobbold LC, Jopling CL, Cooper RE, Wilson LA, Stoneley M, et al.: **Canonical initiation factor requirements of the Myc family of internal ribosome entry segments.** *Mol Cell Biol* 2009, **29**:1565-1574.
26. Watatani Y, Ichikawa K, Nakanishi N, Fujimoto M, Takeda H, Kimura N, et al.: **Stress-induced translation of ATF5 mRNA is regulated by the 5'-untranslated region.** *J Biol Chem* 2008, **283**:2543-2553.
27. Pendleton LC, Goodwin BL, Solomonson LP, Eichler DC: **Regulation of endothelial argininosuccinate synthase expression and NO production by an upstream open reading frame.** *J Biol Chem* 2005, **280**:24252-24260.
28. Park EH, Lee JM, Pelletier J: **The Tie2 5' untranslated region is inhibitory to 5' end-mediated translation initiation.** *FEBS Lett* 2006, **580**:1309-1319.
29. Martin MM, Buckenberger JA, Knoell DL, Strauch AR, Elton TS: **TGF-beta(1) regulation of human AT(1) receptor mRNA splice variants harboring exon 2.** *Mol Cell Endocrinol* 2006, **249**:21-31.
30. Krummheuer J, Johnson AT, Hauber I, Kammler S, Anderson JL, Hauber J, et al.: **A minimal uORF within the HIV-1 vpu leader allows efficient translation initiation at the downstream env AUG.** *Virology* 2007, **363**:261-271.
31. Hurlin PJ: **N-Myc functions in transcription and development.** *Birth Defects Res C Embryo Today* 2005, **75**:340-352.
32. Jasty R, van Golen C, Lin HJ, Solomon G, Heidelberger K, Polverini P, et al.: **Bcl-2 and M-Myc coexpression increases IGF-IR and features of malignant growth in neuroblastoma cell lines.** *Neoplasia* 2001, **3**:304-313.
33. Nanbru C, Prats AC, Droogmans L, Defrance P, Huez G, Kruys V: **Translation of the human c-myc P0 tricistronic mRNA involves two independent internal ribosome entry sites.** *Oncogene* 2001, **20**:4270-4280.
34. Oyama M, Kozuka-Hata H, Suzuki Y, Semba K, Yamamoto T, Sugano S: **Diversity of translation start sites may define increased complexity of the human short ORFeome.** *Mol Cell Proteomics* 2007, **6**:1000-1006.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/9/445/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

