



HAL
open science

The AP-1 binding sites located in the pol gene intragenic regulatory region of HIV-1 are important for virus infectivity

Nathalie Vandenhoudt, Stéphane de Walque, Benoît van Driessche, Laurence Colin, Valérie Martinelli, Allan Guiguen, Caroline Vanhulle, Arsène Burny, Georges Herbein, Olivier Rohr, et al.

► To cite this version:

Nathalie Vandenhoudt, Stéphane de Walque, Benoît van Driessche, Laurence Colin, Valérie Martinelli, et al.. The AP-1 binding sites located in the pol gene intragenic regulatory region of HIV-1 are important for virus infectivity. *Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts*, Sep 2009, Montpellier, France. pp.P89, 10.1186/1742-4690-6-S2-P89 . inserm-00663598

HAL Id: inserm-00663598

<https://inserm.hal.science/inserm-00663598>

Submitted on 27 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Poster presentation

Open Access

The AP-1 binding sites located in the *pol* gene intragenic regulatory region of HIV-1 are important for virus infectivity

Nathalie Vandenhoudt*¹, Stéphane de Walque¹, Benoît Van Driessche¹, Laurence Colin¹, Valérie Martinelli¹, Allan Guiguen¹, Caroline Vanhulle¹, Arsène Burny¹, Georges Herbein², Olivier Rohr³ and Carine Van Lint¹

Address: ¹Laboratory of Molecular Virology, Institut de Biologie et de Médecine Moléculaires (IBMM), Université Libre de Bruxelles (ULB), 6041 Gosselies, Belgium, ²Department of Virology, University of Franche-Comte, St-Jacques Hospital, 25030 Besançon, France and ³INSERM U575, Virology Institute, 67000 Strasbourg, France

* Corresponding author

from *Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts* Montpellier, France. 21-23 September 2009

Published: 24 September 2009

Retrovirology 2009, **6**(Suppl 2):P89 doi:10.1186/1742-4690-6-S2-P89

This abstract is available from: <http://www.retrovirology.com/content/6/S2/P89>

© 2009 Vandenhoudt et al; licensee BioMed Central Ltd.

We have previously identified three AP-1 binding sites in the *pol* gene of human immunodeficiency virus type 1 (HIV-1) and shown that short oligonucleotides containing these sites functioned as phorbol ester-inducible enhancers (Van Lint et al., 1991, *J. Virol.*, 65:7066-7072). These sites are located in a region, called fragment 5103, exhibiting a phorbol ester-inducible enhancing activity on the viral thymidine kinase promoter in HeLa cells. In this study, we have further characterized each of the AP-1 binding sites and have shown that transcription factors c-Fos, JunB and JunD interacted *in vitro* with these motifs. For each site, we have identified mutations abolishing AP-1 factor binding without altering the underlying amino acid sequence of the HIV-1 reverse transcriptase. By transient transfection assays, we have demonstrated that the intragenic AP-1 binding sites were entirely responsible for the PMA-dependent transcriptional activity of fragment 5103. Moreover, this PMA-stimulated activity of fragment 5103 was inhibited by a dominant-negative A-Fos mutant provided the AP-1 sites were not mutated. Finally, we have investigated the biological significance of the intragenic AP-1 binding sites in HIV-1 replication and have shown that these sites are important for viral infectivity.