



HAL
open science

Elevated soluble Flt1 mediates an anti-angiogenic state in patients with ANCA-associated vasculitis

Sandrine Le Roux, Ruth Pepper, Alexandre Dufay, Mélanie Néel, Noël Lamandé, Marie Rimbart, Régis Josien, Mohamed Hamidou, Maryvonne Hourmant, H Terence Cook, et al.

► **To cite this version:**

Sandrine Le Roux, Ruth Pepper, Alexandre Dufay, Mélanie Néel, Noël Lamandé, et al.. Elevated soluble Flt1 mediates an anti-angiogenic state in patients with ANCA-associated vasculitis. 5th European Workshop on Immune-Mediated Inflammatory Diseases, Dec 2010, Sitges-Barcelona, Spain. pp.P13, 10.1186/1479-5876-8-S1-P13 . inserm-00663873

HAL Id: inserm-00663873

<https://inserm.hal.science/inserm-00663873v1>

Submitted on 27 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

Elevated soluble Flt1 mediates an anti-angiogenic state in patients with ANCA-associated vasculitis

S Le Roux^{1,2,3†}, R Pepper^{4†}, A Dufay², M Néel^{1,2,3}, N Lamandé⁵, M Rimbert^{1,6}, R Josien^{1,3,6}, M Hamidou^{3,7}, M Hourmant^{1,2,3}, H T Cook⁸, B Charreau^{1,2,3}, E Larger⁵, A Salama^{4*}, F Fakhouri^{1,2,3*}

From 5th European Workshop on Immune-Mediated Inflammatory Diseases Sitges-Barcelona, Spain. 1-3 December 2010

Introduction

Anti-neutrophil cytoplasm antibody-associated vasculitides (AAV) represent a group of necrotizing small vessel vasculitides that include Wegener's granulomatosis, microscopic polyangiitis and Churg-Strauss-Syndrome. To date, little is known regarding endothelial cell survival and vessel regeneration in patients with AAV, despite the increasingly recognised role of vascular endothelial growth factor (VEGF) in mediating vessel repair.

Aim

Assess the role of sFlt1 in patients with AAV.

Material and methods

Were included 40 patients with PR3-AAV during active disease (n=20) or remission (n=20) and 23 patients with MPO-AAV during active disease (n=10) or remission (n=13). Eighteen additional PR3-AAV patients had paired serum samples drawn at the onset of the disease and at three months of follow-up.

Results

Serum levels of soluble Flt1 (sFlt1), a potent inhibitor of VEGF, are significantly increased during the acute phase of PR3-AAV (2592 ± 11347 pg/ml) and MPO-AAV (476 ± 4258 pg/ml) compared to controls (118 ± 269 pg/ml). sFlt1 levels decreased during disease remission but remained increased compared to controls in patients with PR3-AAV. sFlt1 serum levels correlated with serum levels of C5a, an anaphylatoxin released following complement activation. Serum from patients with acute

PR3-AAV induced the release of sFlt1 by human monocytes *in vitro*, but failed to induce a similar effect on endothelial cells. Pre-treatment of monocytes with an anti-C5a receptor blocking antibody attenuated sFlt1 release. Serum from patients with acute AAV induced a disruption of blood flow in the chicken chorioallantoic membrane assay and this effect was prevented by incubating patients' serum with an excess of human VEGF.

Conclusion

Our data indicate that a complement mediated-increase in sFlt1 occurs during acute AAV which leads to an "anti-angiogenic" state that hinders endothelial repair. "Pro-angiogenic" therapies, which would include complement activation inhibitors, may enhance endothelial repair during AAV and thus reduce renal vascular scarring.

Author details

¹INSERM UMR 643, Nantes, France. ²CHU Nantes, Institute of Transplantation Urology Nephrology, Nephrology Dept., Nantes, France. ³Université de Nantes, Faculté de Médecine, Nantes, France. ⁴Centre for Nephrology, University College London, Royal Free Hospital, London, UK. ⁵INSERM 833, Collège de France, Paris, France. ⁶CHU de Nantes, Immunology Laboratory, Nantes, France. ⁷CHU de Nantes, Internal medicine Dept., Nantes, France. ⁸Centre for Complement and Inflammation Research, Imperial College, London, UK.

Published: 25 November 2010

doi:10.1186/1479-5876-8-S1-P13

Cite this article as: Le Roux *et al.*: Elevated soluble Flt1 mediates an anti-angiogenic state in patients with ANCA-associated vasculitis. *Journal of Translational Medicine* 2010 **8**(Suppl 1):P13.

† Contributed equally

¹INSERM UMR 643, Nantes, France

⁴Centre for Nephrology, University College London, Royal Free Hospital, London, UK

Full list of author information is available at the end of the article