



**HAL**  
open science

## **P19-25. A vaccine model to prevent the depletion of uninfected bystander CD4 cells during HIV infection**

Vincent Vieillard, Hugues Fausther-Bovendo, Roger Le Grand, Patrice Debre

### ► To cite this version:

Vincent Vieillard, Hugues Fausther-Bovendo, Roger Le Grand, Patrice Debre. P19-25. A vaccine model to prevent the depletion of uninfected bystander CD4 cells during HIV infection. *AIDS Vaccine* 2009, Oct 2009, Paris, France. pp.P345, 10.1186/1742-4690-6-S3-P345 . inserm-00663919

**HAL Id: inserm-00663919**

**<https://inserm.hal.science/inserm-00663919>**

Submitted on 27 Jan 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Poster presentation

Open Access

## PI9-25. A vaccine model to prevent the depletion of uninfected bystander CD4 cells during HIV infection

V Vieillard\*<sup>1</sup>, H Fausther-Bovendo<sup>1</sup>, R Le Grand<sup>2</sup> and P Debre<sup>1</sup>

Address: <sup>1</sup>INSERM UMR945, Paris, France and <sup>2</sup>Neuro Virologie, CEA-DSV/DRM, Fontenay-aux-Roses, France

\* Corresponding author

from AIDS Vaccine 2009  
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, 6(Suppl 3):P345 doi:10.1186/1742-4690-6-S3-P345

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P345>

© 2009 Vieillard et al; licensee BioMed Central Ltd.

### Background

The CD4 depletion in the chronic phase of HIV infection is mostly due to the loss of uninfected cells. We recently showed that the expression of NKp44L, a cellular ligand of an activating NK receptor was exclusively expressed on non-infected CD4 cells from HIV-infected patients. NKp44L is specially induced by the highly conserved 3S motif of the HIV-1 gp41 envelope protein. In this study we sought to determine whether the loss of uninfected bystander CD4 cells could be prevented by an anti-3S vaccine.

### Methods

Ten cynomolgus macaques were primed/boosted in IFA with 3S-KLH or free KLH, as control. Sixteen weeks after the last immunization, all animals were challenged IV with the SHIV162P3. Peripheral and lymphoid lymphocyte samples were periodically tested.

### Results

We discovered that 1) During HIV infection, uninfected CD4 cells exclusively express NKp44L; 2) In a macaque model of SHIV162P3 infection, a 3S-KLH immunization significantly decreases NKp44L expression on CD4 cells, NK cells cytotoxicity and apoptosis, when compared to infected control group immunized with free KLH. Interestingly, the CD4 cells level of immunized animals remains stable, while it decreased in the control group.

### Conclusion

These results emphasize the deleterious role of NK cells on the depletion of uninfected bystander CD4 cells. They

also present new opportunities for HIV vaccine strategies, which should subsequently inhibits and/or delay the disease evolution to AIDS.