Supplementary data

Table 7 Phospholipid concentrations of milk lipid extracts from different animal species by ³¹P NMR spectrum.

PL classes	Concentration ^a (µg/ml)				
	Cow	Camel	Mare	Human	
LPA	0 ± 0	0 ± 0	0.66 ± 0.73	0.29 ± 0.66	
LPE	0.32 ± 0.90^{a}	$0 \pm 0^{\rm b}$	$4.37 \pm 2.48^{\circ}$	$3.68 \pm 2.23^{\circ}$	
PA	0 ± 0^{a}	3.60 ± 3.64^{b}	2.29 ± 2.49^{ab}	0.89 ± 1.43^{ab}	
CL	0 ± 0	2.25 ± 6.36	0 ± 0	0.11 ± 0.39	
EPLAS	10.16 ± 8.18^{a}	30.15 ± 18.48^{b}	$3.21 \pm 2.56^{\circ}$	29.91 ± 11.08^{b}	
PE	65.99 ± 33.37^{a}	124.51 ± 41.34^{b}	$14.80 \pm 5.00^{\circ}$	50.76 ± 23.56^{a}	
SM	44.01 ± 22.42^{a}	112.76 ± 44.32^{b}	$17.80 \pm 3.95^{\circ}$	80.47 ± 23.14^{b}	
PS	23.35 ± 14.10^{a}	$43.19 \pm 22.30^{\rm b}$	$6.76 \pm 4.14^{\circ}$	23.60 ± 9.62^{a}	
LPC	0 ± 0^{a}	0 ± 0^{ac}	$1.01 \pm 0.91^{\rm b}$	0.76 ± 1.18^{bc}	
PI	$9.72 \pm 7.25^{\circ}$	20.71 ± 8.32^{b}	$5.62 \pm 1.57^{\circ}$	11.58 ± 4.67^{a}	
aaPC	0 ± 0^{a}	4.77 ± 3.62^{ab}	0.78 ± 0.43^{ab}	$1.17 \pm 2.43^{\rm b}$	
PC	59.95 ± 29.21^{a}	79.78 ± 29.61^{a}	$17.00 \pm 3.44^{\rm b}$	64.66 ± 23.80^{a}	
unknown	3.99 ± 0.77	0 ± 0	0 ± 0	0.36 ± 0.72	
Total PL	213.50 ± 110.90^{a}	419.32 ± 144.58^{b}	$73.72 \pm 13.10^{\circ}$	268.10 ± 87.59^{a}	
PL% of total lipid	0.57 ± 0.32^{a}	$1.00 \pm 0.42^{\rm b}$	1.15 ± 0.35^{b}	0.55 ± 0.27^{a}	

^a Results are means \pm SD of 15 samples for cow, 8 samples for camel and mare, 22 samples for human milk; the means within each row labelled by different letters are significantly different (Mann-Whitney test, P < 0.05).

Table 8Relative proportion of phospholipid classes in the milk from different animal species.

PL classes	Relative proportion ^a (mol% of total PLs)				
	Cow	Camel	Mare	Human	
LPA	0 ± 0	0 ± 0	1.66 ± 1.92	0.21 ± 0.46	
LPE	0.25 ± 0.73^{a}	0 ± 0^{a}	9.22 ± 3.90^{b}	$2.52 \pm 1.60^{\circ}$	
PA	0 ± 0^{a}	$0.80 \pm 0.75^{\rm b}$	3.58 ± 3.86^{ab}	0.32 ± 0.50^{ab}	
CL	0 ± 0	0.37 ± 1.04	0 ± 0	0.05 ± 0.15	
EPLAS	4.91 ± 2.53^{a}	7.01 ± 1.82^{a}	4.19 ± 2.96^{a}	$11.49 \pm 2.04^{\rm b}$	
PE	31.82 ± 2.88^{a}	$30.79 \pm 2.72^{\rm b}$	$19.65 \pm 5.41^{\circ}$	$18.83 \pm 3.37^{\circ}$	
SM	20.02 ± 1.38^{a}	26.14 ± 5.81^{b}	$22.66 \pm 4.55^{\text{b}}$	$29.33 \pm 2.28^{\circ}$	
PS	9.97 ± 2.81	10.84 ± 7.36	8.24 ± 4.44	8.42 ± 2.34	
LPC	0 ± 0^{a}	0 ± 0^{a}	2.04 ± 1.64^{b}	$0.55 \pm 0.88^{\circ}$	
PI	3.72 ± 1.87^{a}	4.22 ± 0.71^{a}	6.18 ± 0.91^{b}	3.67 ± 1.10^{a}	
aaPC	0 ± 0^{a}	0.45 ± 0.71^{ab}	0.23 ± 0.44 ab	$0.39 \pm 0.70^{\rm b}$	
PC	28.68 ± 2.73^{a}	19.14 ± 1.31^{b}	$22.35 \pm 3.22^{\circ}$	$24.05 \pm 3.01^{\circ}$	
unknown	0.63 ± 1.30	0.24 ± 0.45	0 ± 0	0.18 ± 0.55	

^a Results are means \pm SD of 15 samples for cow, 8 samples for camel and mare, 22 samples for human milk; the means within each row labelled by different letters are significantly different (Mann-Whitney test, P < 0.05).

Table 9Lipid globules size of the milk from different animal species.

Milk species	Diameter ^a (µm)		
	$d_{(4,3)}$	$d_{(v0,5)}$	
Cow	3.36 ± 1.39^{a}	3.02 ± 1.25^{a}	
Camel	1.57 ± 0.42^{b}	1.16 ± 0.69^{b}	
Mare	$1.51 \pm 0.37^{\rm b}$	0.44 ± 0.16^{b}	

^a Results are means \pm SD of 15 samples for cow, 8 samples for camel and mare; the means within each column labelled by different letters are significantly different (Mann-Whitney test, P < 0.05).