



**HAL**  
open science

## **NAMPT/Visfatin expression by inflammatory monocytes mediates arthritis pathogenesis by promoting IL-17-producing T cells**

Jessy Pr sumey, Gabriel Courties, Pascale Louis-Plence, Virginie Escriou, Daniel Scherman, Yves-Marie Pers, Hans Yssel, J r me P ne, Diego Kyburz, Steffen Gay, et al.

### ► To cite this version:

Jessy Pr sumey, Gabriel Courties, Pascale Louis-Plence, Virginie Escriou, Daniel Scherman, et al.. NAMPT/Visfatin expression by inflammatory monocytes mediates arthritis pathogenesis by promoting IL-17-producing T cells. 7th European Workshop on Immune-Mediated Inflammatory Diseases, Nov 2012, Noordwijk aan Zee, Netherlands. pp.P48. inserm-00758223

**HAL Id: inserm-00758223**

**<https://inserm.hal.science/inserm-00758223>**

Submitted on 28 Nov 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin e au d p t et   la diffusion de documents scientifiques de niveau recherche, publi s ou non,  manant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv s.



POSTER PRESENTATION

Open Access

# NAMPT/Visfatin expression by inflammatory monocytes mediates arthritis pathogenesis by promoting IL-17-producing T cells

Jessy Présumev<sup>1,2</sup>, Gabriel Courties<sup>1,2</sup>, Pascale Louis-Plence<sup>1,2</sup>, Virginie Escriou<sup>3,4,5,6</sup>, Daniel Scherman<sup>3,4,5,6</sup>, Yves-Marie Pers<sup>1,2,7</sup>, Hans Yssel<sup>1,2</sup>, Jérôme Pène<sup>1,2</sup>, Diego Kyburz<sup>8</sup>, Steffen Gay<sup>8</sup>, Christian Jorgensen<sup>1,2,7</sup>, Florence Apparailly<sup>1,2,7\*</sup>

From 7th European Workshop on Immune-Mediated Inflammatory Diseases Noordwijk aan Zee, the Netherlands. 28-30 November 2012

## Background

Nicotinamide phosphoribosyltransferase (NAMPT)/PBEF/Visfatin exerts multiple functions and has been implicated in the pathogenesis of rheumatoid arthritis. The expression of NAMPT is increased during inflammation and is identified as a novel mediator of innate immunity. To gain insight into its role in arthritis and given that NAMPT induces IL-6 expression that is critical for Th17 lymphocytes, we hypothesized that NAMPT-stimulated production of IL-6 by monocytes might in turn promote Th17 cells.

## Materials and methods

siRNA uptake and NAMPT expression were determined in Ly6C<sup>high</sup> and Ly6C<sup>low</sup> monocyte subsets following intravenous injection of siRNA against NAMPT (siNAMPT) or non-targeting siRNA (siCT) formulated with the DMA-PAP cationic liposome into mice. Mice with established collagen-induced arthritis (CIA) were treated weekly after disease onset with siNAMPT or siCT and clinical features were assessed. T helper cell frequencies, cytokine production and percentage of IL-6-producing Ly6C<sup>high</sup> monocytes were analyzed. Using a coculture system consisting of purified CD14<sup>+</sup> monocytes and autologous CD4<sup>+</sup> T cells, NAMPT and cytokine production, as well as the percentage of IL-17-producing CD4<sup>+</sup> T cells were determined following transfection of CD14<sup>+</sup> monocytes with siCT or siNAMPT.

## Results

Upon intravenous injection, siRNA was preferentially engulfed by Ly6C<sup>high</sup> monocytes and siRNA-mediated silencing of NAMPT expression in Ly6C<sup>high</sup> monocytes reduced IL-6 production by these cells, mitigated Th17 cell expansion, and inhibited inflammatory features and CIA progression. Moreover, NAMPT-RNAi-silenced CD14<sup>+</sup> monocytes were found to reduce the percentage of IL-17-producing CD4<sup>+</sup> T cells.

## Conclusions

Taken together, our results show that the expression of NAMPT in Ly6C<sup>high</sup> monocytes promotes Th17 cells. Our findings provide new mechanistic insight into the action of NAMPT in arthritis and demonstrate the utility of targeting disease-causing genes in Ly6C<sup>high</sup> monocytes for therapeutic intervention in arthritis.

## Author details

<sup>1</sup>INSERM, U844, Montpellier, France. <sup>2</sup>University of Medicine, Montpellier, France. <sup>3</sup>INSERM, U1022, UFR des Sciences Pharmaceutiques et Biologiques, Paris, France. <sup>4</sup>CNRS, UMR8151, Paris, France. <sup>5</sup>University of Pharmacy Paris Descartes, Paris, France. <sup>6</sup>Ecole Nationale Supérieure de Chimie de Paris, Paris, France. <sup>7</sup>University Hospital of Montpellier, Clinical Dept. for osteoarticular diseases, Montpellier, France. <sup>8</sup>Center of Experimental Rheumatology, University Hospital Zurich and Zurich Center of Integrative Human Physiology, Zurich, Switzerland.

Published: 28 November 2012

doi:10.1186/1479-5876-10-S3-P48

**Cite this article as:** Présumev et al.: NAMPT/Visfatin expression by inflammatory monocytes mediates arthritis pathogenesis by promoting IL-17-producing T cells. *Journal of Translational Medicine* 2012 **10**(Suppl 3):P48.

<sup>1</sup>INSERM, U844, Montpellier, France  
Full list of author information is available at the end of the article