

Fast functional imaging of multiple brain regions in intact zebrafish larvae using Selective Plane Illumination Microscopy

Raphaël Candelier, Thomas Panier, Sebastián Romano, Raphaël Olive, Thomas Pietri, Germán Sumbre, Georges Debrégeas

▶ To cite this version:

Raphaël Candelier, Thomas Panier, Sebastián Romano, Raphaël Olive, Thomas Pietri, et al.. Fast functional imaging of multiple brain regions in intact zebrafish larvae using Selective Plane Illumination Microscopy. the Twenty Second Annual Computational Neuroscience Meeting: CNS*2013, Jul 2013, Paris, France. pp.P97, 10.1186/1471-2202-14-S1-P97. inserm-00842318

HAL Id: inserm-00842318 https://inserm.hal.science/inserm-00842318

Submitted on 8 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION



Open Access

Fast functional imaging of multiple brain regions in intact zebrafish larvae using Selective Plane Illumination Microscopy

Raphaël Candelier^{1*}, Thomas Panier¹, Sebastián Romano^{2,3,4,5}, Raphaël Olive¹, Thomas Pietri^{2,3,4,5}, Germán Sumbre^{2,3,4,5}, Georges Debrégeas¹

From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013 Paris, France. 13-18 July 2013

The optical transparency and the small dimensions of zebrafish at the larval stage make it a vertebrate model of choice for brain-wide *in-vivo* functional imaging. However, current point-scanning imaging techniques, such as two-photon or confocal microscopy, impose a strong limit on acquisition speed which in turn sets the number of neurons that can be simultaneously recorded [1]. At 5 Hz, this number is of the order of one thousand, *i.e.* approximately 1-2% of the brain. We demonstrate that this limitation can be greatly overcome by using Selective-Plane Illumination Microscopy (SPIM) [2-4]. Zebrafish larvae expressing the genetically encoded calcium indicator GCaMP3 were illuminated with a scanned laser

sheet and imaged with a camera whose optical axis was oriented orthogonally to the illumination plane. This optical sectioning approach was shown to permit functional imaging of most of the brain volume of 5-9 day old larvae with single-cell resolution. The spontaneous activity of up to 5000 neurons was recorded at 20 Hz for 20-60 min. By rapidly scanning the specimen in the axial direction, the activity of 25000 individual neurons from 5 different z-planes (approximately 30% of the entire brain) could be simultaneously monitored at 4 Hz. Compared to point-scanning techniques, this imaging strategy thus yields a ~20-fold increase in data throughput (number of recorded neurons times acquisition rate) without



correlated activity.

* Correspondence: raphael.candelier@upmc.fr

¹CNRS / UPMC Univ. Paris 06, FRE 3231, Laboratoire Jean Perrin LJP, F-75005, Paris, France

Full list of author information is available at the end of the article



© 2013 Candelier et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

compromising the signal-to-noise ratio. The extended field of view offered by the SPIM method allowed us to directly identify large scale ensembles of neurons, spanning several brain regions (see Figure 1), that displayed correlated activity and were thus likely to participate in common neural processes.

Author details

¹CNRS / UPMC Univ. Paris 06, FRE 3231, Laboratoire Jean Perrin LJP, F-75005, Paris, France. ²Ecole Normale Supérieure, Institut de Biologie de l'ENS, IBENS, Paris, F-75005 France. ³Inserm, U1024, Paris, F-75005 France. ⁴CNRS, UMR 8197, Paris, F-75005 France. ⁵IBENS, ENS, Paris, France.

Published: 8 July 2013

References

- Christine Grienberger, Arthur Konnerth: Imaging calcium in neurons. Neuron 2012, 73(5):862-885.
- 2. Michael Weber, Jan Huisken: Light sheet microscopy for real-time developmental biology. *Curr Opin Genet Dev* 2011, 21(5):566-572.
- Jerome Mertz: Optical sectioning microscopy with planar or structured illumination. Nature Methods 2011, 8(10):811-819, October.
- Raju Tomer, Khaled Khairy, Philipp JKeller: Shedding light on the system: studying embryonic development with light sheet microscopy. Curr Opin Genet Dev 2011, 21(5):558-565.

doi:10.1186/1471-2202-14-S1-P97

Cite this article as: Candelier *et al.*: Fast functional imaging of multiple brain regions in intact zebrafish larvae using Selective Plane Illumination Microscopy. *BMC Neuroscience* 2013 14(Suppl 1):P97.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

BioMed Central

Submit your manuscript at www.biomedcentral.com/submit