

Measuring cognitive change in subjects with prodromal Alzheimer's disease.

Thibault Mura, Cécile Proust-Lima, Hélène Jacqmin-Gadda, Tasnime N. Akbaraly, Jacques Touchon, Bruno Dubois, Claudine Berr

▶ To cite this version:

Thibault Mura, Cécile Proust-Lima, Hélène Jacqmin-Gadda, Tasnime N. Akbaraly, Jacques Touchon, et al.. Measuring cognitive change in subjects with prodromal Alzheimer's disease.. Journal of Neurology, Neurosurgery and Psychiatry, 2014, 85 (4), pp.363-70. 10.1136/jnnp-2013-305078. inserm-00854469

HAL Id: inserm-00854469 https://inserm.hal.science/inserm-00854469v1

Submitted on 27 Aug 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Measuring cognitive change in subjects with prodromal Alzheimer's

disease

T Mura*, MD, PhD ^{1, 2, 3, 4}, C Proust-Lima, PhD ^{5, 6}, H Jacqmin-Gadda PhD ^{5, 6}, TN Akbaraly

PhD ^{1,2,7}, J. Touchon, MD, PhD^{8,2}, B Dubois, MD, PhD ⁹, C Berr, MD, PhD ^{1,2,8}.

¹ INSERM, U1061, Neuropsychiatrie: Recherche Epidémiologique et Clinique, 34093

Montpellier, Cedex 5, France

² Université Montpellier I, 34095 Montpellier, Cedex 5, France

³ Département d'Information Médicale & Centre d'Investigation Clinique, CHRU Montpellier,

34093 Montpellier, France

⁴ INSERM, CIC 1001, Montpellier, France

⁵ INSERM U897, Equipe de Biostatistique, Centre de Recherche en Epidémiologie et

Biostatistique, F-33076 Bordeaux, France

⁶ Université Bordeaux Segalen, ISPED, F-33076 Bordeaux, France

⁷ Department of Epidemiology and Public Health, University College London, London,

United Kingdom.

⁸ CMRR Languedoc Roussillon, service de Neurologie, CHRU Montpellier,

34093Montpellier, France

⁹ INSERM-UPMC UMRS 975, Institut de la Mémoire et de la Maladie d'Alzheimer, ICM,

APHP, Salpétrière Hospital, University Paris 6, Paris, France

Corresponding author: Thibault Mura

INSERM U1061, Hôpital La Colombière, 39 Avenue Charles Flahault, BP 34493, 34093


Montpellier, Cedex 5, France.

Phone: 33 (0)4 67 33 23 28, Fax: 33 (0)4 67 33 23 35

email: t-mura@chu-montpellier.fr


1

Figure 1: Conceptualization of the nonlinear mixed model involving a latent process to model cognition from several neuropsychological tests.




- (a) A linear mixed model describes the change over time in the latent cognitive process and evaluates the common effects of covariates on this latent cognitive trajectory
- (b) Test-specific measurement models relate <u>each administration of the</u> psychometric tests with the latent cognitive process, by accounting for and describing the metrological properties of the tests and test-specific associations with covariates.
- (c) Overall effect of a covariate on each specific test is calculated by adding together the effect of the covariate on the latent cognitive process (a) and the test-specific effect (b).

Figure 2: Diagram mapping the administration of the neuropsychological tests and the occurrence of AD during the three-year follow-up (FU) of the study.



^{*} In the event of a suspected conversion, the patient underwent an additional neuropsychological evaluation 6 months later.

Figure 3: Mean annual change for each neuropsychological test according to the occurrence of AD during the follow-up (in latent cognitive process units).




Mean annual change with 95% confidence interval for each neuropsychological test (in latent cognitive process unit) for a 71.8 year-old woman with a low level of education.

*denotes a significant difference (adjusted for age, sex and level of education) between Prodromal-AD and MCI Non-AD (p<0.05), ** for p<0.01


Baddeley Mü was not represented in this figure because of its high level of individual variability; this test did not significantly change over time in any group and was not different between groups.

Figure 4: Metrological properties of the thirteen neuropsychological scores used in the study


Neuropsychological tests with high sensitivity to changes due to prodromal-AD*



Neuropsychological tests with medium sensitivity to changes due to prodromal-AD*



Neuropsychological tests with low sensitivity to changes due to prodromal-AD*



^{*}according to the previous results display in figure 3