



HAL
open science

Cerebral hemodynamic and ventilatory responses to hypoxia, hypercapnia, and hypocapnia during 5 days at 4,350 m.

Thomas Rupp, François Esteve, Pierre Bouzat, Carsten Lundby, Stéphane Perrey, Patrick Levy, Paul Robach, Samuel Verges

► **To cite this version:**

Thomas Rupp, François Esteve, Pierre Bouzat, Carsten Lundby, Stéphane Perrey, et al.. Cerebral hemodynamic and ventilatory responses to hypoxia, hypercapnia, and hypocapnia during 5 days at 4,350 m.. *Journal of Cerebral Blood Flow and Metabolism*, 2014, 34 (1), pp.52-60. 10.1038/jcbfm.2013.167 . inserm-00873387

HAL Id: inserm-00873387

<https://inserm.hal.science/inserm-00873387v1>

Submitted on 24 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LEGENDS TO FIGURES IN ONLINE DATA SUPPLEMENT

Figure 1 online supplement. Hypoxic responses: Ventilatory (V_E , panel A), TCD (TCD CVR, panel B) and NIRS (oxyhemoglobin HbO_2 , panel C; tissue oxygenation index TOI, panel D; deoxyhemoglobin HHb, panel E; total hemoglobin $HbTot$, panel F) responses to isocapnic hypoxia at sea level (SL), at day 1 (D1) and 5 (D5) at altitude. Changes in V_E , TCD CVR and NIRS variables are expressed per % of change in arterial O_2 saturation. * significantly different from SL; + significantly different from D1.

Figure 2 online supplement. Individual hypoxic responses: Ventilatory (V_E , panel A), TCD (TCD CVR, panel B) and NIRS (oxyhemoglobin HbO_2 , panel C; tissue oxygenation index TOI, panel D; deoxyhemoglobin HHb, panel E; total hemoglobin $HbTot$, panel F) responses to isocapnic hypoxia at sea level (SL), at day 1 (D1) and 5 (D5) at altitude. Changes in V_E , TCD CVR and NIRS variables are expressed per mmHg of change in end-tidal O_2 partial pressure. Whiskers under and above the mean values (large horizontal lines) represent the 95% confident interval. * significantly different from SL; + significantly different from D1.

Figure 3 online supplement. Individual hypercapnic responses: Ventilatory (V_E , panel A), TCD (TCD CVR, panel B) and NIRS (oxyhemoglobin HbO_2 , panel C; tissue oxygenation index TOI, panel D; deoxyhemoglobin HHb, panel E; total hemoglobin $HbTot$, panel F) responses to hypercapnia (expressed per mmHg of change in end-tidal CO_2 partial pressure) at sea level (SL), at day 1 (D1) and 5 (D5) at altitude. Whiskers under and above the mean values (large horizontal lines) represent the 95% confident interval. * significantly different from SL; + significantly different from D1.

Figure 4 online supplement. Individual hypocapnic responses: TCD (TCD CVR, panel A) and NIRS (oxyhemoglobin HbO₂, panel B; tissue oxygenation index TOI; panel C; deoxyhemoglobin HHb, panel D; total hemoglobin HbTot, panel E) responses to hypocapnia (expressed per mmHg of change in end-tidal CO₂ partial pressure) at sea level (SL), at day 1 (D1) and 5 (D5) at altitude. Whiskers under and above the mean values (large horizontal lines) represent the 95% confident interval.







