

**Table 2: Clinical characteristics at baseline of the whole cohort**

|                              | UHDRS |      |      |          | UHDRS-FAP |      |      |          |
|------------------------------|-------|------|------|----------|-----------|------|------|----------|
|                              | N     | Mean | SD   | Min/Max  | N         | Mean | SD   | Min/Max  |
| Motor score                  | 53    | 82.5 | 14.8 | [0-124]  | 68        | 24.6 | 8.1  | [0-48]   |
| Behavioural score            | 46    | 14.9 | 9.8  | [0-88]   | 69        | 2.4  | 1.5  | [0-8]    |
| Cognitive score (raw Stoop)  | 42    | 44.9 | 41.6 | [0->660] | 33        | 56.0 | 43.9 | [0->455] |
| Cognitive score (Stroop log) |       |      |      |          | 33        | 33.6 | 23.2 | [0->80]  |
| Somatic score                |       |      |      |          | 64        | 9.4  | 6.0  | [0-32]   |
| Independence Scale           | 53    | 42.3 | 16.6 | [10-100] |           |      |      |          |
| Functional Assessment        | 48    | 44.7 | 4.8  | [25-50]  |           |      |      |          |
| TFC                          | 53    | 1.7  | 1.4  | [0-13]   |           |      |      |          |

N= number of patients, SD= standard deviation, TFC= Total Functional Capacity;  
UHDRS = Unified Huntington Disease Rating Scale; UHDRS-FAP = UHDRS For  
Advanced Patients