

Sex and age-related differences in performance in a 24-hour ultra-cycling draft-legal event - a cross-sectional data analysis

Lara Pozzi, Beat Knechtle, Patrizia Knechtle, Thomas Rosemann, Romuald Lepers, Christoph Alexander Rüst

► To cite this version:

Lara Pozzi, Beat Knechtle, Patrizia Knechtle, Thomas Rosemann, Romuald Lepers, et al.. Sex and age-related differences in performance in a 24-hour ultra-cycling draft-legal event - a cross-sectional data analysis. BMC Sports Science, Medicine and Rehabilitation, 2014, 6 (1), pp.19. inserm-00998246

HAL Id: inserm-00998246 https://inserm.hal.science/inserm-00998246

Submitted on 30 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Open Access

Sex and age-related differences in performance in a 24-hour ultra-cycling draft-legal event – a cross-sectional data analysis

Lara Pozzi¹, Beat Knechtle^{1,2*}, Patrizia Knechtle², Thomas Rosemann¹, Romuald Lepers³ and Christoph Alexander Rüst¹

Abstract

Background: The purpose of this study was to examine the sex and age-related differences in performance in a draft-legal ultra-cycling event.

Methods: Age-related changes in performance across years were investigated in the 24-hour draft-legal cycling event held in Schötz, Switzerland, between 2000 and 2011 using multi-level regression analyses including age, repeated participation and environmental temperatures as co-variables.

Results: For all finishers, the age of peak cycling performance decreased significantly ($\beta = -0.273$, p = 0.036) from 38 ± 10 to 35 ± 6 years in females but remained unchanged ($\beta = -0.035$, p = 0.906) at 41.0 ± 10.3 years in males. For the annual fastest females and males, the age of peak cycling performance remained unchanged at 37.3 ± 8.5 and 38.3 ± 5.4 years, respectively. For all female and male finishers, males improved significantly ($\beta = 7.010$, p = 0.006) the cycling distance from 497.8 ± 219.6 km to 546.7 ± 205.0 km whereas females ($\beta = -0.085$, p = 0.987) showed an unchanged performance of 593.7 ± 132.3 km. The mean cycling distance achieved by the male winners of 960.5 ± 51.9 km was significantly (p < 0.001) greater than the distance covered by the female winners with 769.7 ± 65.7 km but was not different between the sexes (p > 0.05). The sex difference in performance for the annual winners of 19.7 ± 7.8% remained unchanged across years (p > 0.05). The achieved cycling distance decreased in a curvilinear manner with advancing age. There was a significant age effect (F = 28.4, p < 0.0001) for cycling performance where the fastest cyclists were in age group 35–39 years.

Conclusion: In this 24-h cycling draft-legal event, performance in females remained unchanged while their age of peak cycling performance decreased and performance in males improved while their age of peak cycling performance remained unchanged. The annual fastest females and males were 37.3 ± 8.5 and 38.3 ± 5.4 years old, respectively. The sex difference for the fastest finishers was ~20%. It seems that women were not able to profit from drafting to improve their ultra-cycling performance.

Keywords: Cycling, Master athletes, Sex difference, Ultra-endurance

* Correspondence: beat.knechtle@hispeed.ch

¹Institute of General Practice and Health Services Research, University of Zurich, Zurich, Switzerland

²Gesundheitszentrum St. Gallen, St. Gallen, Switzerland

Full list of author information is available at the end of the article



© 2014 Pozzi et al.; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

Background

The most traditional endurance and ultra-endurance sports are swimming, cycling, running, and triathlon as a combination of them. In recent years, several studies reported an increased participation in ultra-endurance performances – defined as an endurance performance of six hours and longer [1] – such as ultra-running [2-4], ultracycling [5-8] and ultra-triathlon [9,10]. For several of these ultra-endurance events, an increased participation and an improvement in performance of master athletes older than 35 years [11] have been observed [5,12-14].

Several recent studies analysed also the influence of age and sex on triathlon performance [9,15-17]. There were also studies focusing on the influence of age and sex on running performance [18-20], however, only a few studies investigated other endurance disciplines such as swimming [21,22] or cycling [23,24]. Cycling as a non-weightbearing activity represents an interesting model because it can be performed even in older ages [7] because of its non-technical and its non-weight-bearing character [25].

Age has been reported as an important predictor variable in ultra-endurance athletes such as ultra-marathoners [26]. An age-related decline in endurance performance is inevitable even if master athletes tend nowadays to improve their performance [12,13,18]. Endurance performance starts to decline after the age of ~55 years independently of the physical activity [25]. However, there seemed to be differences in the age-related performance decline regarding the different endurance disciplines. Ransdell et al. demonstrated that the age-related decline in endurance performance was exponential after the age of ~55 years for both sexes in swimming, cycling and running [25]. However, the age-related decline was less pronounced in swimming and cycling compared to running when master competitors in running, swimming and cycling were investigated [25]. For triathletes, Bernard et al. [27] and Lepers et al. [28] showed a less pronounced age-related decline in endurance performance in cycling compared to running and swimming. In contrast to Ransdell et al. [25], Baker and Tang [29] reported for sprint cycling that female and male master record performances decreased with advancing age in a similar manner as for swimming, rowing, weightlifting, triathlon and running. Balmer et al. [23] showed that cycling performance declined in an indoor 16.1-km time-trial with increasing age.

Apart from the age-related performance decline, the age at which peak performance is achieved would be of interest for endurance athletes such as cyclists to plan their career. Cycling races can be held without drafting such as cycling time trials [30], mountain bike cycling races [31], ultra-endurance cycling races [7,8] or cycling time trials in long-distance triathlons [9,15], or with drafting such as traditional road cycling races. Regarding ultra-cycling, a recent study investigating the age trends from 2001 to 2012 in a 720 km ultra-cycling race reported that the fastest female and male performance was achieved at the age of 35.9 ± 9.6 and 38.7 ± 7.8 years, respectively [7]. In a 120 km mountain bike cycling race held between 1994 and 2012, the age of the fastest athletes was lower than in a 720 km ultra-cycling race where the fastest females achieved the fastest race times at the age of $30.7 \pm$ 5.0 years and males at the age of 27.1 ± 3.4 years [31]. These recent studies investigating performance in ultracycling used data from races where drafting was forbidden such as the 120 km mountain bike ultra-cycling race 'Swiss Bike Master' [31], the 720 km 'Swiss Cycling Marathon' as a gualifier for the 'Race across America' (RAAM) [7], and the 'RAAM' itself [8]. These studies showed that the participation in females was low [5,31], the performance decreased in males compared to females [31] but improved in females compared to males [7] across years, and the sex difference in performance remained unchanged [8] or decreased [7,31]. The sex difference in performance for the fastest finishers was at $\sim 20 \pm 10\%$ in these races [5,7,8].

Drafting during cycling permits reducing average power output, oxygen consumption (VO₂) and heart rate and therefore to improve performance [32]. Furthermore, it results in a reduction in frontal resistance and reduced energy cost at a given submaximal intensity [33]. However, no data exist about the age of peak ultra-cycling performance in a draft-legal cycling race in contrast to a non-drafting ultracycling performance [7]. Performance between females and males in a draft-legal ultra-cycling race might also be different as it has been shown for non-drafting ultra-cycling races [7]. Drafting might enhance female performance as it has been shown for ultra-swimming. For example, in the 34-km 'English Channel Swim' from Dover (Great Britain) to Calais (France) where athletes have to cover the distance alone, the fastest men were faster than the fastest women [34]. However, in the 46-km 'Manhattan Island Marathon Swim' where drafting is allowed, the best women were ~12-14% faster than the best men [35]. To date, no study investigated the age and sex of finishers in an ultra-cycling performance where drafting is allowed. If males and females were allowed to ride together with drafting, the sex difference in cycling performance might be lower compared to non-drafting conditions.

The aims of the study were to investigate the participation trends, and the sex and age-related differences in performance in ultra-cycling in a 24-hour draft-legal ultraendurance cycling event held in Schötz, Switzerland, between 2000 and 2011. Considering existing findings in ultra-swimming, we hypothesized for a draft-legal cycling race that the sex difference in ultra-cycling performance would be lower compared to previous observations in ultra-endurance running or ultra-endurance cycling performance where drafting was not allowed. Pozzi et al. BMC Sports Science, Medicine, and Rehabilitation 2014, 6:19 http://biomedcentral.com/2052-1847/6/19



Methods

Ethics

This study was approved by the Institutional Review Board of St. Gallen, Switzerland, with waiver of the requirement for informed consent given that the study involved the analysis of publicly available data.

The race

To test our hypothesis, the age at the time of the competition and the achieved cycling distance (km) of all male and female solo riders at the 24-hours cycling race Schötz ('24 Stunden Schötz') were analysed from 2000 to 2011. Since 2000, the '24 Stunden Schötz' has been held each year in the city of Schötz, Switzerland, in the first weekend of august. The last edition was held in 2011. Many athletes used this race to prepare for the 'Race Across AMerica' (RAAM). Several winners of the '24 Stunden Schötz' won later the 'RAAM'. The flat circuit extended over a distance of 9,888 m. In each lap, the athletes had to overcome 35 m difference in altitude. Solo and team riders were competing in the same field. The laps were counted electronically, drafting was allowed and solo riders were generally drafting behind team riders. The athletes had the opportunity to be assisted during the race by a personal support crew.

Data collection and data analysis

The data set from this study was obtained from the race website of the '24 Stunden Schötz' (www.24stundenrennen.ch) and from the race director for data of earlier years where the age of the athletes was missing. Between 2000 and 2011, a total of 916 finishers (i.e. 83 females and 833 males) completed the race. The 24-hour cycling performance was expressed in km. The age at the time of the competition and the cycling performances of all male and female competitors were analysed from 2000 to 2011. The magnitude of the sex difference was examined by calculating the percent difference between the female and male winner for each year. The effect of age on the 24-hour cycling performance was only analysed in males because the number of female finishers in the different age groups was too small for an accurate data analysis (Figure 1A). Because the age of both the annual male winners and all annual male finishers did not significantly change across the years, we pooled the data of the 12 years for the ten fastest males for each age group. The age groups distinguish the categories for each period of 5 years as follows: < 25 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years, 50-54 years, 55-59 years, and 60–64 years. Therefore, the best top ten performances of the athletes in nine age groups during the studied period were considered.

Statistical analyses

Each set of data was tested for normal distribution using D'Agostino and Pearson omnibus normality test and for homogeneity of variances using Levene's test prior to statistical analyses. Trends in participation were analysed using linear regression whereas results were tested for linearity with run's test. Single and multi-level regression analyses were used to investigate changes in performance and age of the finishers. A hierarchical regression model avoided the impact of a cluster-effect on results where a particular athlete finished more than once. Regression analyses of performance were corrected for age of athletes to prevent a misinterpretation of the 'age-effect' as a 'time-effect' since age is an important predictor variable in ultraendurance performance [26]. Regression models were also corrected with environmental temperatures (i.e. lowest and the highest temperature during every race) since environmental conditions such as extreme heat impairs endurance [30,36] and ultra-endurance performance [37,38]. Historical weather data with air temperature for this analysis were provided by "MeteoSchweiz" (www.meteoschweiz.admin.ch) (Table 1). The performance of the top ten athletes per age group were compared to the performance of the fastest age group using one-way analysis of variance (ANOVA) with Dunnett post-hoc analysis. Statistical analyses were performed using IBM SPSS Statistics (Version 22, IBM SPSS, Chicago, IL, USA) and GraphPad Prism (Version 6.01, GraphPad Software, La Jolla, CA, USA). Significance was accepted at p < 0.05 (two-sided for *t*-tests). Data in the text and figures are given as mean ± standard deviation (SD).

Table 1 Daily maximum (T max) and daily minimum (T min) temperatures on race days

Year	T max (°C)	T min (°C)
2000	20	18
2001	27	20
2002	20	17
2003	29	21
2004	26	18
2005	22	16
2006	15	10
2007	24	16
2008	25	19
2009	22	15
2010	20	16
2011	19	15

Weather data were acquired from 'MeteoSchweiz' (www.meteoschweiz.admin.ch).

Results

Participation trends and age groups

A total of 916 finishers (*i.e.* 83 females and 833 males) completed the race between 2000 and 2011. The annual number of finishers over the history of the event is shown in Figure 1A. The annual number of males decreased across years whereas the annual number of females remained unchanged. Between 2000 and 2011, the annual number of finishers was 70 ± 14 (range: 44–88) for males and 7 ± 2 (range: 4–11) for females, respectively. Females accounted on average for $9.2 \pm 3.0\%$ of the field over the 12-years period. The age distribution of both female and male finishers during the 12-years period is displayed in Figure 1B. Most of the male finishers were ranked in age group 35–39





years whereas most of the female finishers were in the age groups 25–29 and 35–39 years. Cyclists older than 40 years of age represented ~53% of the male and ~35% of the female finishers. Expressed in percent of all finishers, most of the male finishers were ranked in age group 25–29 years and most of the female finishers in age group 35–39 years (Figure 1C).

The age of the cyclists

The age of the finishers for each sex is shown in Figure 2A for all annual female and male finishers and in Figure 2B for annual female and male winners. For all female and male annual finishers, the age of peak cycling performance decreased significantly in females from 38 ± 10 years (2000) to 35 ± 6 years (2011) (Table 2). In males, however, the age of peak cycling performance remained unchanged at 41.0 ± 10.3 years (Table 2). For the annual fastest females and males, the age of peak cycling performance remained unchanged at 37.3 ± 8.5 and 38.3 ± 5.4 years, respectively.

The performance of the cyclists

The achieved cycling distance (km) of all finishers for both sexes is presented in Figure 3A and for the annual winners in Figure 3B. Overall males improved their cycling distance significantly from 497.8 ± 219.6 km (2000) to 546.7 ± 205.0 km (2011) whereas overall females showed an unchanged performance over time of 593.7 ± 132.3 km (Table 3). The mean cycling distance covered by the male winners (960.5 ± 51.9 km) was significantly (p < 0.001) greater than the mean distance covered by the female winners (769.7 ± 65.7 km) (Figure 3B) but did not significantly change across the years for both sexes (Table 3).

The sex difference in cycling performance

The sex difference in cycling performance for the winners during the 2000–2011 period is shown in Figure 4. The mean sex difference was $19.7 \pm 7.8\%$ (range: 7.7-33.0%) and did not significantly change across years (Table 4).

The age-related changes in male cycling performances

The mean age-related change for the achieved cycling distance in the ten fastest males for each age group throughout the 2000–2011 period is shown in Figure 5. The achieved cycling distance decreased in a curvilinear manner with advancing age. There was a significant age effect (F = 28.4, p < 0.0001) for the cycling performance. The fastest cyclists were in age group 35–39 years. Cyclists younger than 34 years and cyclists older than 45 years were significantly slower than cyclists aged from 35–44 years.

Discussion

The main findings were that females accounted for $\sim 9\%$ of the field, the sex difference for the annual winners

was unchanged at ~20%, the performance in females remained unchanged while their age of peak cycling performance decreased and the performance in males improved while their age of peak cycling performance remained unchanged, and the annual fastest females and males were 37.3 ± 8.5 and 38.3 ± 5.4 years old, respectively.

Participation trends regarding women and master athletes

Females accounted on average for ~9% of the field over the 12-years period. This percentage is in line with recent findings for non-drafting ultra-cycling races. By comparison, female finishers accounted for ~11% in both the 'RAAM' and the 'Furnace Creek 508' as a qualifier for the 'RAAM' [5]. In another qualifier for the 'RAAM', the 'Swiss Cycling Marathon', female participation was at ~3% [5]. In other ultra-endurance sports disciplines such as running, female participation was higher. By comparison, in a 161 km ultra-marathon, female participation reached ~20% in 2004 [2]. Ultra-running seems to have a higher popularity than ultra-cycling regarding female participation.

Ultra-cycling seems to be of interest for master athletes. In the present study, cyclists older than 40 years of age

Table 2 Multi-level regression analyses for change in age across years (Model 1) with correction for multiple finishes (Model 2), daily lowest air temperature (Model 3) and daily highest air temperature (Model 4) for the annual fastest and all finishers

Model	β	SE (β)	Stand. β	т	Р
		All male	finishers		
1	0.018	0.256	0.008	0.069	0.945
2	0.018	0.256	0.008	0.069	0.945
3	0.003	0.282	0.001	0.009	0.993
4	-0.035	0.297	-0.015	-0.119	0.906
		All femal	e finishers		
1	-0.241	0.106	0.079	2.284	0.023
2	-0.241	0.106	0.079	2.284	0.023
3	-0.205	0.094	-0.076	-2.193	0.029
4	-0.273	0.129	-0.073	-2.106	0.036
	Ar	nnual fastes	t male finisher	s	
1	0.245	0.472	0.162	0.519	0.615
2	0.245	0.472	0.162	0.519	0.615
3	0.088	0.490	0.058	0.179	0.862
4	-0.191	0.464	-0.126	-0.410	0.691
	Anı	nual fastest	female finishe	rs	
1	-1.042	0.669	-0.442	-1.558	0.150
2	-1.042	0.669	-0.442	-1.558	0.150
3	-0.917	0.725	-0.389	-1.265	0.238
4	-0.585	0.723	-0.248	-0.809	0.439



represented ~53% of male finishers and ~35% of female finishers. Similar findings have been reported for nondrafting races. In two qualifiers for the 'RAAM', the 'Swiss Cycling Marathon' and the 'Furnace Creek 508', ~46% of the finishers were between 35 and 49 years of age [5]. These findings in ultra-cycling differ to findings in other ultra-endurance sports disciplines such as running. In a 161 km ultra-marathon, runners older than 40 years of age account for 65-70% of the finishers [2]. In ultra-running, Hoffman *et al.* [2] showed that the increase in the overall participation in the 161 km ultra-marathons held in North America was mostly due to the increasing number of master and female athletes . In contrast to ultra-cycling, ultra-running seems to have a higher popularity among master

Table 3 Multi-level regression analyses for change in performance across years (Model 1) with correction for multiple finishes (Model 2), age of athletes with multiple finishes (Model 3), daily lowest air temperature (Model 4) and daily highest air temperature (Model 5) for the annual fastest and all finishers

Model	β	SE (β)	Stand. β	т	Р
		All male	e finishers		
1	5.614	2.222	0.087	2.527	0.012
2	5.614	2.222	0.087	2.527	0.012
3	5.452	2.229	0.085	2.446	0.015
4	6.424	2.317	0.100	2.773	0.006
5	7.010	2.538	0.109	2.762	0.006
		All fema	le finishers		
1	-0.193	4.315	-0.005	-0.045	0.964
2	-0.193	4.315	-0.005	-0.045	0.964
3	-0.176	4.335	-0.005	-0.041	0.968
4	0.632	4.773	0.016	0.132	0.895
5	-0.085	5.027	-0.002	-0.017	0.987
	An	nual fastest	female finishe	rs	
1	8.004	5.176	0.439	1.546	0.153
2	8.004	5.176	0.439	1.546	0.153
3	5.506	5.784	0.302	0.952	0.366
4	6.581	5.945	0.361	1.107	0.300
5	7.074	6.152	0.388	1.150	0.283
	Ar	nnual fastes	t male finisher	s	
1	-3.330	4.429	-0.231	-0.752	0.469
2	-3.330	4.429	-0.231	-0.752	0.469
3	-2.098	3.993	-0.146	-0.525	0.612
4	-2.745	4.296	-0.191	-0.639	0.541
5	-3.347	4.657	-0.233	-0.719	0.493

athletes. This could be due to the fact that only a small and a less expensive equipment is needed for ultrarunning compared to ultra-cycling [6]. This makes the access to running easier in general because it is easier and less expensive to start training and racing.

The sex difference in ultra-cycling performance

The mean cycling distance covered by the male winners was significantly greater than the distance covered by the female winners but did not significantly change across the years for both sexes. Similar findings of an unchanged sex difference in ultra-cycling performance have been reported for the 'RAAM' from 1982–2012 [8]. However, in the 720 km 'Swiss Cycling Marathon' held between 2001 and 2012 [7] and the 120 km 'Swiss Bike Masters' held between 1994 and 2009 [31], the sex difference in performance decreased. These disparate findings might be explained by the different lengths of the races (*i.e.* ~4,800 km in the

'RAAM', 720 km in the 'Swiss Cycling Marathon', 120 km in the 'Swiss Bike Masters' and ~600 km in the present race) and the different lengths of the investigated time periods (*i.e.* 31 years in the 'RAAM', 12 years in the 'Swiss Cycling Marathon', 16 years in the 'Swiss Bike Masters, and 12 years in the present race). The changes in cycling equipment (*e.g.* weight and aerodynamics of the bike) during a 12-year period might be too small to influence ultracycling performance.

It was hypothesized that the sex difference in ultracycling performance would be lower compared to previous observations in ultra-cycling. Interestingly, the mean sex difference in performance between male and female winners was ~20%. This value is close to the 23% sex difference found in the 540 km cycling split in a Triple-Iron triathlon where drafting was not allowed [9]. Also in nondrafting ultra-cycling races, the sex difference in performance for the fastest cyclist was at ~20% in the 'RAAM' $(25.0 \pm 11.9\%)$, in the 'Swiss Cycling Marathon' $(27.8 \pm$ 9.4%) and in the 'Furnace Creek 508' (18.4 ± 13.9%) [5]. For the annual three fastest in the 'RAAM', the sex difference remained unchanged at $24.6 \pm 3.0\%$ [8]. For the annual three fastest in the 'Swiss Cycling Marathon', the sex difference decreased from $35.0 \pm 9.5\%$ in 2001 to $20.4 \pm$ 7.7% in 2012 [7].

A special aspect regarding the ultra-endurance cycling performance in this study was drafting. Individual riders have the possibility to cycle behind the team riders and the teams were consistently changing their riders so a high average cycling speed can be sustained over the whole 24 hours for the leaders. Hausswirth and Brisswalter [32] showed that drafting in cycling permits to decrease average power output, VO₂ and heart rate and therefore improve the performance in long-duration cycling events. Riders can cycle at a higher velocity with lower relative intensity while drafting [33]. Because drafting allows cycling at a lower VO_2 [32], drafting might be an advantage for individuals with a lower VO₂max such as master and female athletes to achieve a better performance than in a nondrafting race. However, subjects with higher VO₂ (i.e. males) can also sustain higher velocities.

Obviously, females could not profit from drafting in this 24-hour ultra-cycling draft-legal event. Limiting factors for ultra-endurance performance in females are their smaller muscle mass [39], their lower peak values $inVO_2$ [40] and their higher body fat [39] compared to males. Females present a disadvantage compared to males because they have a higher percent of body fat, even if they were equally well-trained [40]. In an Ironman triathlon, for example, the lower body fat was associated with a faster Ironman race time for males [41]. In long-distance triathletes, a lower percent of body fat is related to a better race performance. In female triathletes, the percent of body fat has a strong and positive relationship with total race time and

both running and cycling split times. In males, however, body fat is positively related to total race time and running split time [42]. In ultra-runners, Hoffmann *et al.* [43] showed in a 161 km ultra-marathon that percent body fat was lower for finishers than for non-finishers and that percent body fat was significantly associated with race times in males.

Another explanation for the lower performance in females compared to males might be the power-to-weight ratio (PWR). Women have a lower body mass and might benefit from lower body mass in climbing. However, the main factor accounting for sex differences in peak and mean power output during cycling is the skeletal muscle mass of the lower extremities [44]. Since female athletes have a lower muscle mass [39], the power in cycling might be more limited by the skeletal muscle mass than by body mass. Additionally, laps in the '24 Stunden Schötz' were flat where no climbing was needed.

In this study, we included the daily highest and the daily lowest air temperatures as co-variables to investigate a

Table 4 Multi-level regression analyses for change in sex difference across years (Model 1) with correction for daily lowest air temperature (Model 2) and daily highest air temperature (Model 3) for the annual fastest finishers

Model	β	SE (β)	Stand. β	Т	Р		
Annual fastest finishers							
1	-1.103	0.585	-0.512	-1.887	0.089		
2	-1.209	0.635	-0.561	-1.903	0.089		
3	-1.106	0.695	-0.513	-1.590	0.146		

potential influence of environmental temperatures on performance since it has been reported for both runners [37,38] and cyclists [30,36] that air temperature has an influence on performance. However, performance and sex difference in performance was not influenced. This might be due to the rather moderate maximum temperatures of $23 \pm 4^{\circ}$ C (range $15-29^{\circ}$ C) and minimum temperatures of $17 \pm 3^{\circ}$ C (range $10-21^{\circ}$ C). Cycling performance was impaired at air temperatures of > 30°C in a 20 km time trial [36].

The age-related decline in ultra-cycling performance

For all female and male finishers, the age of peak cycling performance decreased in females from 38 ± 10 years to 35 ± 6 years whereas in males, it remained unchanged at 41.0 ± 10.3 years. For the annual fastest females and males, the age of peak cycling performance remained unchanged at 37.3 ± 8.5 and 38.3 ± 5.4 years, respectively. The 24-hour cycling performance decreased in a curvilinear manner with advancing age and the best cycling performances were obtained in the age group 35-39 years for males. Cyclists younger than 34 years and cyclists older than 45 years were significantly slower than cyclists aged from 35-44 years.

The findings for the age of peak cycling performance were very similar to the findings in the 720 km 'Swiss Cycling Marathon' where the fastest males and females were at the age of 35.9 ± 9.6 and 38.7 ± 7.8 years, respectively [7]. These findings for ultra-cyclists are comparable to the findings for ultra-marathoners competing in the 'Western State 100-Mile Endurance Run' where the fastest race times in



Pozzi et al. BMC Sports Science, Medicine, and Rehabilitation 2014, 6:19 http://biomedcentral.com/2052-1847/6/19 males were not significant different between the age groups <30 years, 30–39 years and 40–49 years [14]. Our data are also consistent with the finding that the decline in cycling performance in triathlon was not significant until the age of ~55 years [27]. However, the age-related decline in performance in 24-h road cycling appears to start later compared with a 120 km endurance mountain cycling race where a significant decline in performance was observed after the age of ~35 years [24].

The main physiological mechanism leading to a decline in performance with increasing age in endurance sports seems to be a progressive reduction of maximum oxygen uptake (VO_2max) [45]. However, also other physiological factors such as a lower maximum strength of the lower limb muscles and a decrease in skeletal muscle mass in general [29], a slower rate of force development and transmission, and a reduction in elastic energy storage and recovery in tendons could cause a decrease in cycling performance with increasing age [46]. Cyclists younger than 34 years and cyclists older than 45 years were significantly slower than cyclists at the age of 35-44 years. This could be due to the fact that younger athletes have a lack of experience in preparation, nutrition and competing in ultra-endurance races compared to older athletes. The importance of previous experience was reported for ultramarathoners where the personal best marathon time was the only predictor variable for race performance in a 24hour run [47]. Also for ultra-endurance cyclists, previous personal best time was an important predictor variable [48]. Additionally, older athletes were training more and for a longer time regarding their whole life than younger athletes, who maybe just started to compete. This is in line with the finding that both personal best time and training volume were associated with race time in a mountain bike ultra-marathon [48].

Limitations of the study

A limitation of this study is that we did not consider the training of the cyclists, their diet or their improvement in equipment. In female Ironman triathletes, training volume was related to overall race time [39] and in male ultraendurance cyclists, both training and nutrition were related to performance [6]. Carbohydrate loading before running a marathon influences performance because it allows a runner of a given aerobic capacity and leg muscle distribution to run at greater speeds without 'hitting the wall', succumbing to the failure mode associated with the exhaustion of glycogen reserves and the form supplemental carbohydrate is consumed can influence the effectiveness of midrace fuelling [49]. Also, the increased cycling distance in male cyclists could be due to an improvement in equipment [50].

Conclusion

To summarize, performance in females remained unchanged while their age of peak cycling performance decreased and performance in males improved while their age of peak cycling performance remained unchanged. For the annual winners, the mean cycling distance covered by males was significantly greater than the distance covered



by females but did not significantly change across the years for both sexes. The annual fastest females and males were 37.3 ± 8.5 and 38.3 ± 5.4 years old, respectively. The sex difference for the fastest finishers was ~20% in this 24-h cycling draft-legal event and very similar to other reports from races where drafting was not allowed. It seems that women were not able to profit from the possibility of drafting to improve their ultra-cycling performance.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

LP drafted the manuscript, BK and PK collected the data, and CAR and RL performed the statistical analyses, BK, CAR, TR and RL helped in drafting the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank the race director for providing us missing data of the athletes.

Author details

¹Institute of General Practice and Health Services Research, University of Zurich, Zurich, Switzerland. ²Gesundheitszentrum St. Gallen, St. Gallen, Switzerland. ³INSERM U1093, University of Burgundy, Faculty of Sport Sciences, Dijon, France.

Received: 3 February 2014 Accepted: 9 May 2014 Published: 15 May 2014

References

- Zaryski C, Smith DJ: Training principles and issues for ultra-endurance athletes. Curr Sports Med Rep 2005, 4:165–170.
- 2. Hoffman MD, Ong JC, Wang G: Historical analysis of participation in 161 km ultramarathons in North America. *Int J Hist Sport* 2010, 27:1877–1891.
- Cejka N, Rüst CA, Lepers R, Onywera V, Rosemann T, Knechtle B: Participation and performance trends in 100-km ultra-marathons worldwide. J Sports Sci 2013, Epub ahead of print.
- Rüst CA, Knechtle B, Rosemann T, Lepers R: Analysis of performance and age of the fastest 100-mile ultra-marathoners worldwide. *Clinics (Sao Paulo)* 2013, 68:605–611.
- Shoak MA, Knechtle B, Knechtle P, Rüst CA, Rosemann T, Lepers R: Participation and performance trends in ultracycling. Open Access J Sports Med 2013, 4:41–51.
- Knechtle B, Knechtle P, Rüst CA, Rosemann T, Lepers R: Finishers and nonfinishers in the 'Swiss Cycling Marathon ' to qualify for the 'Race Across America'. J Strength Cond Res 2011, 25:3257–3263.
- Zingg M, Knechtle B, Rüst CA, Rosemann T, Lepers R: Age and gender difference in non-drafting ultra-endurance cycling performance - the 'Swiss Cycling Marathon'. Extrem Physiol Med 2013, 2:18.
- Rüst CA, Knechtle B, Rosemann T, Lepers R: Men cross America faster than women-the "Race Across America" from 1982 to 2012. Int J Sports Physiol Perform 2013, 8:611–617.
- Knechtle B, Knechtle P, Lepers R: Participation and performance trends in ultra-triathlons from 1985 to 2009. Scand J Med Sci Sports 2011, 21:e82–e90.
- Rüst CA, Knechtle B, Knechtle P, Lepers R, Rosemann T, Onywera V: European athletes dominate performances in Double Iron ultra-triathlons – a retrospective data analysis from 1985 to 2010. Eur J Sport Sci 2014, 14(Suppl 1):S39–S50.
- 11. Reaburn P, Dascombe B: Endurance performance in master athletes. Eur Rev Aging Phys Act 2008, 5:31–42.
- Lepers R, Rüst CA, Stapley P, Knechtle B: Relative improvements in endurance performance with age: Evidence from 25 years of Hawaii Ironman racing. *Age (Dordr)* 2013, 35:953–962.
- Knechtle B, Rüst CA, Rosemann T, Lepers R: Age-related changes in 100-km ultra-marathon running performances. Age (Dordr) 2012, 34:1033–1045.
- Hoffman MD, Wegelin JA: The Western States 100-Mile Endurance Run: participation and performance trends. *Med Sci Sports Exerc* 2009, 41:2191–2198.

- Lepers R, Maffiuletti NA: Age and gender interactions in ultra-endurance performance: insight from triathlon. *Med Sci Sports Exerc* 2011, 43:134–139.
- Sultana F, Brisswalter J, Lepers R, Hausswirth C, Bernard T: Effects of age and gender on Olympic triathlon performances. Sci Sports 2008, 23:130–135.
- Lepers R, Knechtle B, Stapley PJ: Trends in triathlon performance: Effects of sex and age. Sports Med 2013, 43:851–863.
- 18. Lepers R, Cattagni T: Do older athletes reach limits in their performance during marathon running? *Age (Dordr)* 2012, **34**:773–781.
- Hoffman MD: Performance trends in 161-km ultramarathons. Int J Sports Med 2010, 31:31–37.
- Leyk D, Erley O, Ridder D, Leurs M, Rüther T, Wunderlich M, Sievert A, Baum K, Essfeld D: Age-related changes in marathon and half-marathon performances. Int J Sports Med 2007, 28:513–517.
- Donato AJ, Tench K, Glueck DH, Seals DR, Eskurza I, Tanaka H: Declines in physiological functional capacity with age: a longitudinal study in peak swimming performance. J Appl Physiol 2003, 94:764–769.
- 22. Eichenberger E, Knechtle B, Knechtle P, Rüst CA, Rosemann T, Lepers R: No gender difference in peak performance in ultra-endurance swimming performance Analysis of the 'Zurich 12-h Swim' from 1996 to 2010. *Chin J Physiol* 2012, **55**:346–351.
- Balmer J, Bird S, Davison R: Indoor 16.1-km time-trial performance in cyclists aged 25–63 years. J Sports Sci 2008, 26:57–62.
- Haupt S, Knechtle B, Knechtle P, Rüst CA, Rosemann T, Lepers R: The agerelated performance decline in ultra-endurance mountain biking. *Res Sports Med* 2013, 21:146–158.
- Ransdell LB, Vener J, Huberty J: Master athletes: an analysis of running, swimming and cycling performance by age and gender. J Exerc Sci Fit 2009, 7:61–73.
- Knechtle B, Knechtle P, Rosemann T, Lepers R: Predictor variables for a 100-km race time in male ultra-marathoners. *Percept Mot Skills* 2010, 111:681–693.
- Bernard T, Sultana F, Lepers R, Hausswirth C, Brisswalter J: Age-related decline in Olympic triathlon performance: effect of locomotion mode. *Exp Aging Res* 2010, 36:64–78.
- Lepers R, Sultana F, Bernard T, Hausswirth C, Brisswalter J: Age-related changes in triathlon performances. Int J Sports Med 2010, 31:251–256.
- Baker AB, Tang YQ: Aging performance for masters records in athletics, swimming, rowing, cycling, triathlon, and weightlifting. *Exp Aging Res* 2010, 36:453–477.
- Abbiss CR, Burnett A, Nosaka K, Green JP, Foster JK, Laursen PB: Effect of hot versus cold climates on power output, muscle activation, and perceived fatigue during a dynamic 100-km cycling trial. J Sports Sci 2010, 28:117–125.
- Gloor RU, Knechtle B, Knechtle P, Rüst CA, Haupt S, Rosemann T, Lepers R: Sex-related trends in participation and performance in the 'Swiss Bike Masters' from 1994–2012. Percept Mot Skills 2013, 116:640–654.
- Hausswirth C, Brisswalter J: Strategies for improving performance in long duration events. Olympic distance triathlon. Sports Med 2008, 38:881–891.
- Bentley DJ, Millet GP, Vleck VE, McNaughton LR: Specific aspects of contemporary triathlon. Implications for physiological analysis and performance. Sports Med 2002, 32:345–359.
- Fischer G, Knechtle B, Rüst CA, Rosemann T: Male swimmers cross the English Channel faster than female swimmers. Scand J Med Sci Sports 2013, 23:e48–e55.
- Knechtle B, Rosemann T, Lepers R, Rüst CA: Women outperform men in ultra-distance swimming - the 'Manhattan Island Marathon Swim' from 1983 to 2013. Int J Sports Physiol Perform 2014, Epub ahead of print.
- Peiffer JJ, Abbiss CR: Influence of environmental temperature on 40 km cycling time-trial performance. Int J Sports Physiol Perform 2011, 6:208–220.
- 37. Wegelin JA, Hoffman MD: Variables associated with odds of finishing and finish time in a 161-km ultramarathon. *Eur J Appl Physiol* 2011, 111:145–153.
- Parise CA, Hoffman MD: Influence of temperature and performance level on pacing a 161 km trail ultramarathon. Int J Sports Physiol Perform 2011, 6:243–251.
- Knechtle B, Wirth A, Baumann B, Knechtle P, Rosemann T, Senn O: Differential correlations between anthropometry, training volume and performance in male and female Ironman triathletes. J Strength Cond Res 2010, 24:2785–2793.
- Knechtle B, Müller G, Willmann F, Kotteck K, Eser P, Knecht H: Fat oxidation in men and women endurance athletes in running and cycling. Int J Sports Med 2004, 25:38–44.

- Knechtle B, Wirth A, Rüst CA, Rosemann T: The relationship between anthropometry and split performance in recreational male Ironman triathletes. Asian J Sports Med 2011, 2:23–30.
- Landers GJ, Blanksby BA, Ackland TR, Smith D: Morphology and performance of world championship triathletes. Ann Hum Biol 2000, 27:387–400.
- Hoffman MD, Lebus DK, Ganong AC, Casazza GA, Van Loan M: Body composition of 161-km ultramarathoners. Int J Sports Med 2010, 31:106–109.
- Perez-Gomez J, Rodriguez GV, Ara I, Olmedillas H, Chavarren J, González-Henriquez JJ, Dorado C, Calbet JA: Role of muscle mass on sprint performance: gender differences? Eur J Appl Physiol 2008, 102:685–694.
- Tanaka H, Seals DR: Endurance exercise performance in masters athletes: age-associated changes and underlying physiological mechanisms. *J Physiol* 2008, 586:55–63.
- Arampatzis A, Degens H, Baltzopoulos V, Rittweger J: Why do older sprinters reach the finish line later? Exerc Sport Sci Rev 2011, 39:18–22.
- Knechtle B, Wirth A, Knechtle P, Zimmermann K, Kohler G: Personal best marathon performance is associated with performance in a 24-h run and not anthropometry or training volume. *Br J Sports Med* 2009, 43:836–839.
- Knechtle B, Knechtle P, Rosemann T, Senn O: Personal best time and training volume, not anthropometry, is related to race performance in the "Swiss Bike Masters' Mountain Bike Ultramarathon". J Strength Cond Res 2011, 25:1312–1317.
- 49. Rapoport BI: Metabolic factors limiting performance in marathon runners. *PLoS Comput Biol* 2010, 6:e1000960.
- Nielsen H, Lejeune T: Bicycle shock absorption systems and energy expended by the cyclist. Sports Med 2004, 34:71–80.

doi:10.1186/2052-1847-6-19

Cite this article as: Pozzi *et al.*: **Sex and age-related differences in performance in a 24-hour ultra-cycling draft-legal event – a cross-sectional data analysis.** *BMC Sports Science, Medicine, and Rehabilitation* 2014 **6**:19.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

) BioMed Central

Submit your manuscript at www.biomedcentral.com/submit