



HAL
open science

Title: Personality Profiles in Young Adults with Disordered Eating Behavior

Patrick Raynal, Tiffany Melioli, Henri Chabrol

► **To cite this version:**

Patrick Raynal, Tiffany Melioli, Henri Chabrol. Title: Personality Profiles in Young Adults with Disordered Eating Behavior. *Eating Behaviors*, 2016, 22, pp.119-123. inserm-01330903

HAL Id: inserm-01330903

<https://inserm.hal.science/inserm-01330903v1>

Submitted on 13 Jun 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Behaviors

Elsevier Editorial System(tm) for Eating

Manuscript Draft

Manuscript Number:

Title: Personality Profiles in Young Adults with Disordered Eating Behavior

Article Type: Full Length Article

Keywords: Subclinical eating disorders; Personality; Eating behaviors; Cluster classification

Corresponding Author: Dr. Patrick Raynal,

Corresponding Author's Institution:

First Author: Patrick Raynal

Order of Authors: Patrick Raynal; Tiffany Melioli; Henri Chabrol

December 3rd, 2015

To the Editor of Eating Behaviors

Dear Pr. Miller,

Please find enclosed a manuscript entitled "Personality Profiles in Subclinical Eating Disorders".

We would like to submit this work for publication as a full-length paper in *Eating Behaviors* (word counts: abstract: 179; text + references: 3497).

As possible referees, I suggest:

Jennifer D. Slane

VISN 4 Mental Illness Research, Education and Clinical Center, Pittsburgh VA, USA

E-mail: jennifer.slane@va.gov

Kelly L. Klump

Michigan State University, Dept of Psychology, East Lansing, MI, USA

E-mail: klump@msu.edu

Michael W. MacGregor

University of Saskatchewan, Saskatoon, Canada

E-mail: michael.macgregor@usask.ca

I thank you very much in advance for the reviewing process.

Sincerely,

Patrick Raynal, Ph.D.

Senior research scientist

Highlights

- 101 college students with disturbed eating behavior were classified on personality disorder traits
- Schizotypal, autistic, obsessional, borderline and cyclothymic traits were used
- 3 distinct clusters were obtained: high-traits, low-traits and schizotypal-borderline-cyclothymic
- Symptoms of depression, suicidal ideas, obsessive-compulsion and trait anger varied between groups

Running head: PERSONALITY IN DISORDERED EATING BEHAVIOR

Personality Profiles in Young Adults with Disordered Eating Behavior

Patrick Raynal^a, Tiffany Melioli^a, Henri Chabrol^a

^aCentre d'Etudes et de Recherches en Psychopathologie et Psychologie de la

Santé

Université de Toulouse, France

Corresponding author:

P. Raynal, Laboratoire CERPPS, Université de Toulouse-Jean Jaurès, 5 allées Antonio Machado, 31058 Toulouse, France

Phone number : 33 561 22 52 90

Fax number : 33 561 25 70 93

E-mail address: patrick.raynal@inserm.fr

Abstract

Personality traits are closely related to eating disorders (ED) and might be involved in their maintenance and development. Nevertheless little is known regarding the association between personality traits and disordered eating in subclinical populations. College students answered questionnaires assessing disordered eating behaviors (DEB) and the following personality disorder (PD) traits: schizotypal, autistic, obsessional, borderline and cyclothymic. Participants with DEB ($n=101$, 87% women) displayed significantly higher scores for several variables including schizotypy, cyclothymic, borderline and obsessional traits compared to other participants ($n=378$). Cluster analysis in the DEB subsample led to the identification of three groups: 1) a cluster with a high level of traits (HT); 2) a cluster scoring high on schizotypal, borderline and cyclothymic traits (SBC); 3) a cluster with a low level of traits (LT). Symptoms of depression, suicidal ideations, trait anger and obsessive-compulsive symptoms were higher in the HT and the SBC clusters compared to the LT cluster. Given that two thirds of participants suffering from DEB appeared to display a morbid personality profile, taking into account PD traits of individuals with DEB appears of prime importance.

1. Introduction

Personality features might play an important role in both etiology and outcome of eating disorders (ED) (MacGregor & Lamborn, 2014). Personality disorders (PD) are common among individuals suffering from ED with a comorbidity reaching 60 % according to several studies (Cassin & von Ranson, 2005; Wonderlich & Mitchell, 2001). Furthermore, individuals suffering from anorexia nervosa (AN) are frequently affected by obsessive-compulsive PD (Young, Rhodes, Touyz, & Hay, 2013). Regarding individuals with bulimic symptoms, literature data have found a high prevalence of the traits associated with borderline PD (including impulsivity and sensation seeking) (Atiye, Miettunen, & Raevuori-Helkamaa, 2014; MacGregor & Lamborn, 2014; von Lojewski & Abraham, 2014).

While previous ED classifications were mainly based on the presence or absence of obsessive-compulsive or borderline PD traits, more recent studies have introduced larger comorbid personality psychopathology, leading to the description of three personality profiles, observed either in adolescents or in adults: undercontrolled, overcontrolled and low psychopathology profiles which could be associated to distinct illness course and treatment response (Lavender et al., 2013; Turner et al., 2014; Wildes & Marcus, 2013).

Other comorbidities of ED have been explored including autism-spectrum disorders. Recent studies have reported 20% of autism-spectrum traits among individuals with ED (Huke, Turk, Saeidi, Kent, & Morgan, 2013; Westwood et al., 2015). In addition, the presence of bipolar and schizophrenia-spectrum disorders were also described (Kouidrat, Amad, Lalau, & Loas, 2014; Remberk, Bazynska, Krempa-Kowalewska, & Rybakowski, 2014).

A major issue to better understand ED etiology lies in the relationship between clinical and subthreshold ED. Subthreshold ED have a high prevalence among adolescents and young women, representing an important source of physical and psychological impairments including symptoms of anxiety and depression (Chamay-Weber, Narring, & Michaud, 2005; Touchette et al., 2011). Nevertheless the relationship between threshold and subthreshold ED remains unclear, with a lack of consensus regarding whether subthreshold ED represent a less severe form of ED or a transitional phase to fullblown ED. Due to this lack of consensus, the characteristics of patients who will develop clinical ED are still unclear and surprisingly, to date, only a few studies focused on personality features associated to subthreshold ED. To our knowledge, only one study has explored the personality profiles of women with subthreshold ED, using an instrument assessing 18 dimensions of personality disorders. They have found the three personality clusters already emphasized in ED, suggesting that the same personality traits may contribute to both ED and subthreshold ED (Perkins, Slane, & Klump, 2013).

Following a similar approach, the aim of this study was to identify a typology of young adults with DEB based on personality disorder traits. Contrary to prior study, we focused on traits associated to ED and not limited to personality disorders (including traits reflecting the association to autistic, bipolar and schizophrenia spectra, that are obsessive-compulsive traits, borderline traits, autistic traits, cyclothymic traits and schizotypal traits).

2. Methods

2.1. Participants

Participants were recruited through social networks and official website of several French universities. Only those who answered >95% of the items were included in this study. The sample is thus composed of 479 individuals (96 males; 383 females) aged between 18 and 25 (mean age of males = 21 ±2.3; mean age of females = 20.7 ±1.9; $p=.32$). The study followed the ethical guidelines of the Helsinki Declaration and its procedures were approved by the ethics committee of the research ward. Participants provided their willingness to participate through a consent form and no compensation was offered. The questionnaires were anonymous.

2.2. Measures

2.2.1. Eating behavior

Disordered eating was assessed using the French translation of Eating Attitudes Test-26 (EAT-26) (Garner, Olmsted, Bohr, & Garfinkel, 1982; Leichner, Steiger, Puentes-Neuman, Perreault, & Gottheil, 1994) which assesses symptoms of ED through 26 items scored on a 6-point scale, ranging from 1 (never) to 6 (always). An example item is "I avoid eating when I am hungry." A cutoff score of 20 has been identified as indicating DEB (Garner et al., 1982).

2.2.2. Personality traits

2.2.2.1 Schizotypal traits

Schizotypal traits were assessed using the Schizotypal Personality Questionnaire-Brief (SPQ-B) (Raine & Benishay, 1995), a self-administered scale including 22 dichotomous items. Each item (e.g., "People sometimes find me aloof and distant") is scored 0/1 (no/yes).

2.2.2.2. Cyclothymic traits

The *Temperament Evaluation of Memphis, Pisa, Paris and San Diego autoquestionnaire* (TEMPS-A) (Akiskal et al., 2005) was used to measure cyclothymic traits, using the relevant subscale (12 items; e.g., "My ability to think varies greatly from sharp to dull for no apparent reason"). Items were scored 0 (no) or 1 (yes).

2.2.2.3. Obsessional personality traits

Obsessional traits were measured using the French version of the Obsessional personality disorder scale of the *Personality Diagnostic Questionnaire-4* (Bouvard, 2002; Hyler, Rieder, Williams, Spitzer, & Lyons, 1988). This scale contained 8 items (e.g., "I feel that my standards and ethics are higher than those of my peers") scored 0 (false) or 1 (true).

2.2.2.4. Borderline personality traits

Borderline PD traits were assessed using the nine relevant items of the *Personality Diagnostic Questionnaire-4* (Bouvard, 2002; Hyler et al., 1988). Items were scored from 1 (totally false) to 7 (totally true).

2.2.2.5. Autistic traits

Autistic traits were measured using the Autism spectrum Quotient (AQ) (Baron-Cohen, Wheelwright, Skinner, Martin, & Clubley, 2001) under its 10-item version (e.g., "When I am reading a story, I find it difficult to work out the characters intentions"). Items were scored 0 [no] or 1 [yes], except for items 2-6 and 9 that were scored inversely (Booth et al., 2013).

2.2.3. Psychopathological symptoms

2.2.3.1. Depression symptoms and suicidal ideation

Depression was evaluated using The *Centre for Epidemiological Studies-Depression* scale (CES-D) (Radloff, 1977) under its 10-item version (e.g., "My sleep has not been good"). Responses were made on a 4-point Likert scale, ranging from 0 to 3. A short suicidal ideation scale was added (Chabrol, Rodgers, & Rousseau, 2007), containing 3 items (e.g., "I felt life was not worth living") that were scored similarly as CES-D.

2.2.3.2. State-trait anger

State-trait anger was assessed using the second part of the *State-trait anger expression inventory-2* (STAXI-2) (Spielberger, 1988) under its French version (Borteyrou, Bruchon-Schweitzer, & Spielberger, 2008). This scale contained 8 items (e.g., "I get easily angry") scored 1 (almost never) to 4 (almost always).

2.2.3.3. Obsessive-compulsive symptoms

Obsessive-compulsive symptoms were assessed using the French version of the *Obsessive Compulsive Inventory-Revised* (OCI-R) (Foa et al., 2002; Zermatten, Van der Linden, Jermann, & Ceschl, 2006). It contains 18 items (e.g., "I check things more often than necessary") scored from 0 ("Not at all") to 4 ("Extremely").

2.2.4. Substance use

Cannabis use was measured with the *Cannabis Use Disorder Identification Test-Revised* (CUDIT-R) (Adamson et al., 2010). Alcohol consumption was assessed using the first item of the Alcohol Use Disorders Identification Test (AUDIT) (Saunders, Aasland, Babor, Delafuente, & Grant, 1993).

All these questionnaires have demonstrated adequate reliability and validity (Adamson et al., 2010; Akiskal et al., 2005; Baron-Cohen et al., 2001; Bouvard, 2002; Foa et al., 2002; Garner et al., 1982; Hylter et al., 1988; Radloff, 1977; Raine & Benishay, 1995; Saunders et al., 1993; Spielberger, 1988; Zermatten et al., 2006).

2.2.5. Personal information

This study collected general information regarding the participants (age, sex, academic degree and academic results during the past semester). The variable “academic results” was obtained by asking students their overall grade for the last semester (Failed; Passed with grade $\geq 10/20$ and $< 12/20$; Passed with grade $\geq 12/20$ and $< 14/20$; Passed with grade $\geq 14/20$ and $< 16/20$; Passed with grade $\geq 16/20$).

Cronbach alphas and score ranges are presented in Table 1. The obsessive-compulsive personality trait and the autistic trait scales showed suboptimal Cronbach's alphas ranging between 0.50 and 0.70, an acceptable level of consistency for short scales (Schmitt, 1996).

3. Results

Comparison of participants with and without DEB

Participants scoring above 20 on the EAT-26 ($n=101$, 87.3% female, 12.7% male) were compared to the rest of participants (considered as the control group; $n=378$, 77.2% female, 21.8% male). The proportion of female was higher in the group scoring above 20 (the DEB group) compared with the control group ($p=0.018$, Fisher's exact test). The DEB group had higher scores on all personality traits and psychopathological symptoms (Table 1). Cohen's d indicated large size effects.

Cluster analysis

Cluster analysis was conducted among the DEB group in two steps to generate personality profiles, based on personality traits (schizotypal, cyclothymic, borderline, obsessional and autistic) converted to z-scores. Absence of multicollinearity was evaluated through an analysis of the correlations between the variables selected for the cluster analysis (all variables had tolerance values > .40). In the first step, a hierarchical cluster analysis was conducted (Ward's method with squared Euclidean distance). Based on the dendrogram and the aggregation curve, a three-cluster solution was identified. In the second step, K-means clustering was used to assign individuals to one of the identified clusters. A discriminant analysis showed clear differences between clusters (Wilks' $\lambda = 0.145$, $p < 0.001$) with 97.3% of cases correctly classified. Data revealed a first group that was well below the mean of DEB group on all traits (Figure 1; Table 2). This cluster was thus termed the "Low Traits" cluster (LT, $n = 35$ [34.7%], women 94.3%, men 5.7%). A second group was characterized with high scores on schizotypal, borderline and cyclothymic traits and low scores on autistic and obsessional dimensions; it was thus called the "Schizotypal, Borderline and Cyclothymic" cluster (SBC, $n = 33$ [32.7%], women 84.8%, men 15.2%). The last cluster was characterized by scores that were at least a half standard deviation above the mean on all dimensions – schizotypal, autistic, cyclothymic, borderline, obsessional – and was thus named "High Traits" cluster (HT, $n = 33$ [32.7%], women 81.8%, men 18.2%). Interestingly autistic traits were the most elevated traits in this cluster, with AQ-10 scores twice higher than in other clusters. No difference in sex ratio could be observed between the different clusters (p range = 0.11-0.5, Fisher's exact test).

Cluster group differences in level of disturbed eating behaviors, psychopathological variables, frequency of cannabis and alcohol uses, and academic

results were tested using an analysis of variance. Compared to the LT cluster, the HT and SBC clusters participants displayed higher scores on depressive symptoms, suicidal ideas, obsessive-compulsive symptoms and state-trait anger (Table 2). In addition, individuals from the HT group showed a lower level of academic degree in comparison with the LT cluster.

4. Discussion

The aim of this study was to identify a typology of young adults with DEB based on personality disorder traits. To our knowledge, this study is the second to explore personality profiles among young adults with DEB, based on PD traits. Our classification showed convergences and some differences with the previous study aiming at exploring the PD trait profiles of women with DEB (Perkins et al., 2013). Perkins et al.'s study (2013) described three clusters (Adaptive, Dysregulated, Rigid) with two clusters closely matching those identified in our study. The Adaptive cluster corresponds to LT as the group with lowest traits. The Dysregulated group, displaying the highest levels of all traits with the exception of intimacy problems, appeared concordant with HT. Interestingly, autistic traits were the highest component of HT, and differences in autistic traits between HT and other clusters were large (1.5 SD). This finding thus provides further support to the notion of an association between a sub-group of ED and autistic traits (Courty et al., 2013; Oldershaw, Treasure, Hambrook, Tchanturia, & Schmidt, 2011). This study also suggests to pay special attention to autistic traits among participants suffering from DEB.

The Rigid cluster may appear, at first sight, relatively different from SBC. Indeed SBC was composed of individuals with high cognitive and emotional dysregulation (as suggested by high schizotypal, borderline and cyclothymic traits) and low rigidity (as

indicated by low autistic and obsessional traits). Nevertheless, the Rigid cluster showed some dysregulation traits as it displayed a level of affective instability higher than the Adaptive cluster. Furthermore, the compulsivity level in Rigid was lower than in Dysregulated, suggesting that obsessiveness was somewhat limited in Rigid cluster, as it appeared in SBC (Perkins et al., 2013).

At the symptom level, the comparison between our results and Perkins et al.'s study is limited by the difference in study variables. Our clusters did not differ from each other on DEB scores, whereas differences appeared in some disordered eating dimensions in Perkins et al.'s report. In our study, the most striking difference between clusters was the prominence of obsessional-compulsive symptoms, with a score twice higher in HT and SBC than in LT, while all other variables showed more modest inter-cluster variations (Table 2). Considering that HT and SBC altogether represent two thirds of individuals in our sample, these data indicate that obsessive-compulsive symptoms are a very frequent comorbidity of DEB.

This study is not exempt from limitations. Firstly, the small sample size and the low proportion of men may restrict the generalizability of the present finding. Secondly, self-report measures were used, which can induce some bias in data collection. Nevertheless, this study suggests that the classification of DEB in function of PD traits could be extended to men, considering that no difference in sex ratio could be observed between clusters. Moreover, personality traits of the autistic, schizotypal and bipolar spectra may be relevant to the classification of DEB or subclinical ED. As a conclusion, this study emphasizes the importance of taking into account PD traits in DEB. Two thirds of DEB group appeared to display a morbid personality profile which might represent a transitional phase to clinical ED. Further studies are needed to better understand the role and implication of personality profile in the etiology of ED.

References

- Adamson, S. J., Kay-Lambkin, F. J., Baker, A. L., Lewin, T. J., Thornton, L. et al. (2010). An improved brief measure of cannabis misuse: The Cannabis Use Disorders Identification Test-Revised (CUDIT-R). *Drug and Alcohol Dependence, 110*, 137-143.
- Akiskal, H. S., Mendlowicz, M. V., Jean-Louis, G., Rapaport, M. H., Kelsoe, J. R. et al. (2005). TEMPS-A: validation of a short version of a self-rated instrument designed to measure variations in temperament. *Journal of Affective Disorders, 85*, 45-52.
- Atiye, M., Miettunen, J., & Raevuori-Helkamaa, A. (2014). A Meta-Analysis of Temperament in Eating Disorders. *European Eating Disorders Review, 23*, 89-99.
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The autism-spectrum quotient (AQ): Evidence from Asperger Syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders, 31*, 603-603.
- Booth, T., Murray, A. L., McKenzie, K., Kuenssberg, R., O'Donnell, M. et al. (2013). Brief Report: An Evaluation of the AQ-10 as a Brief Screening Instrument for ASD in Adults. *Journal of Autism and Developmental Disorders, 43*, 2997-3000.
- Borteyrou, X., Bruchon-Schweitzer, M., & Spielberger, C. D. (2008). The French adaptation of the STAXI-2, C.D. Spielberger's State-trait anger expression inventory. *Encephale-Revue De Psychiatrie Clinique Biologique Et Therapeutique, 34*, 249-255.
- Bouvard, M. (2002). Questionnaires et échelles d'évaluation de la personnalité. *Masson. Paris.*
- Cassin, S. E., & von Ranson, K. M. (2005). Personality and eating disorders: A decade in review. *Clinical Psychology Review, 25*, 895-916.
- Chabrol, H., Rodgers, R., & Rousseau, A. (2007). Relations between suicidal ideation and dimensions of depressive symptoms in high-school students. *Journal of Adolescence, 30*, 587-600.
- Chamay-Weber, C., Narring, F., & Michaud, P. A. (2005). Partial eating disorders among adolescents: A review. *Journal of Adolescent Health, 37*, 417-427.
- Courty, A., Maria, A. S., Lalanne, C., Ringuenet, D., Vindreau, C. et al. (2013). Levels of autistic traits in anorexia nervosa: a comparative psychometric study. *Bmc Psychiatry, 13*, 222.

- Foa, E. B., Huppert, J. D., Leiberg, S., Langner, R., Kichic, R. et al. (2002). The obsessive-compulsive inventory: Development and validation of a short version. *Psychological Assessment, 14*, 485-496.
- Garner, D. M., Olmsted, M. P., Bohr, Y., & Garfinkel, P. E. (1982). The Eating Attitudes Test - Psychometric Features and Clinical Correlates. *Psychological Medicine, 12*, 871-878.
- Huke, V., Turk, J., Saeidi, S., Kent, A., & Morgan, J. F. (2013). Autism Spectrum Disorders in Eating Disorder Populations: A Systematic Review. *European Eating Disorders Review, 21*, 345-351.
- Hylner, S. E., Rieder, R. O., Williams, J. B., Spitzer, J. H., & Lyons, M. (1988). The Personality Diagnostic Questionnaire: Development and preliminary results. *Journal of Personality Disorders, 2*, 229-237.
- Kouidrat, Y., Amad, A., Lalau, J. D., & Loas, G. (2014). Eating disorders in schizophrenia: implications for research and management. *Schizophrenia Research and Treatment, 2014*, 791573.
- Lavender, J. M., Wonderlich, S. A., Crosby, R. D., Engel, S. G., Mitchell, J. E. et al. (2013). Personality-based subtypes of anorexia nervosa: Examining validity and utility using baseline clinical variables and ecological momentary assessment. *Behaviour Research and Therapy, 51*, 512-517.
- Leichner, P., Steiger, H., Puentes-Neuman, G., Perreault, M., & Gottheil, N. (1994). Validation of the French Version of the Eating Attitudes Test (Eat-26) for Francophone Quebecers. *Canadian Journal of Psychiatry-Revue Canadienne De Psychiatrie, 39*, 49-54.
- MacGregor, M. W., & Lamborn, P. (2014). Personality Assessment Inventory profiles of university students with eating disorders. *Journal of Eating Disorders, 2*, 20.
- Oldershaw, A., Treasure, J., Hambrook, D., Tchanturia, K., & Schmidt, U. (2011). Is Anorexia Nervosa a Version of Autism Spectrum Disorders? *European Eating Disorders Review, 19*, 462-474.
- Perkins, P. S., Slane, J. D., & Klump, K. L. (2013). Personality clusters and family relationships in women with disordered eating symptoms. *Eating Behaviors, 14*, 299-308.
- Radloff, L. S. (1977). The CES-D Scale: A self-report depression scale for research in the general population. *Applied Psychological Measurements, 1*, 385-401.

- Raine, A., & Benishay, D. (1995). The SBQ-B: A brief screening instrument for schizotypal personality disorder. *Journal of Personality Disorders, 9*, 346-355.
- Remberk, B., Bazynska, A. K., Krempa-Kowalewska, A., & Rybakowski, F. (2014). Adolescent insanity revisited: Course and outcome in early-onset schizophrenia spectrum psychoses in an 8-year follow-up study. *Comprehensive Psychiatry, 55*, 1174-1181.
- Saunders, J. B., Aasland, O. G., Babor, T. F., Delafuente, J. R., & Grant, M. (1993). Development of the Alcohol-Use Disorders Identification Test (Audit) - Who Collaborative Project on Early Detection of Persons with Harmful Alcohol-Consumption. *Addiction, 88*, 791-804.
- Schmitt, N. (1996). Uses and abuses of coefficient alpha. *Psychological Assessment, 8*, 350-353.
- Spielberger, C. D. (1988). *Manuel for the state-trait anger expression inventory (STAXI): PAR 2000*.
- Touchette, E., Henegar, A., Godart, N. T., Pryor, L., Falissard, B. et al. (2011). Subclinical eating disorders and their comorbidity with mood and anxiety disorders in adolescent girls. *Psychiatry Research, 185*, 185-192.
- Turner, B. J., Claes, L., Wilderjans, T. F., Pauwels, E., Dierckx, E. et al. (2014). Personality profiles in Eating Disorders: Further evidence of the clinical utility of examining subtypes based on temperament. *Psychiatry Research, 219*, 157-165.
- von Lojewski, A., & Abraham, S. (2014). Personality factors and eating disorders: Self-uncertainty. *Eating Behaviors, 15*, 106-109.
- Westwood, H., Eisler, I., Mandy, W., Leppanen, J., Treasure, J. et al. (2015). Using the Autism-Spectrum Quotient to Measure Autistic Traits in Anorexia Nervosa: A Systematic Review and Meta-Analysis. *Journal of Autism and Developmental Disorders, Nov 5 [Epub ahead of print]*, 1-15.
- Wildes, J. E., & Marcus, M. D. (2013). Alternative methods of classifying eating disorders: Models incorporating comorbid psychopathology and associated features. *Clinical Psychology Review, 33*, 383-394.
- Wonderlich, S., & Mitchell, J. E. (2001). The role of personality in the onset of eating disorders and treatment implications. *Psychiatric Clinics of North America, 24*, 249-258.

- Young, S., Rhodes, P., Touyz, S., & Hay, P. (2013). The relationship between obsessive-compulsive personality disorder traits, obsessive-compulsive disorder and excessive exercise in patients with anorexia nervosa: a systematic review. *Journal of Eating Disorders, 1*, 16.
- Zermatten, A., Van der Linden, M., Jermann, F., & Ceschl, G. (2006). Validation of a French version of the Obsessive-Compulsive Inventory-Revised in a non-clinical sample. *European Review of Applied Psychology-Revue Europeenne De Psychologie Appliquee, 56*, 151-155.

Table 1
Comparison of the disordered eating behavior group (DEB)
with the control group (t-test).

<i>Variables</i>	N=479			Control n=378		DEB n=101		<i>t</i>	<i>Cohen's d</i>
	<i>α</i>	<i>Range</i>	<i>M.</i>	<i>S.D.</i>	<i>M.</i>	<i>S.D.</i>			
Schizotypy	0.78	0-21	8.01	4.25	10.12	4.58	-4.39*	0.48	
Autistic traits	0.53	0-8	3.06	1.59	3.33	1.74	-1.46	0.16	
Cyclothymic traits	0.77	0-12	4.31	3.15	6.64	3.12	-6.64*	0.74	
Borderline traits	0.82	9-60	29.28	10.08	39.50	12.45	-8.62*	0.90	
Obsessional traits	0.58	0-8	3.58	1.54	4.02	1.52	-2.59*	0.29	
Depression	0.69	1-25	12.16	3.90	15.02	3.83	-6.59*	0.74	
Suicidal ideas	0.87	0-9	0.65	1.37	1.71	2.38	-5.79*	0.55	
Obsess.-compulsive symptoms	0.83	0-64	17.00	10.49	22.66	11.92	-4.70*	0.50	
State-trait anger	0.86	8-32	18.05	5.07	20.63	5.66	-4.44*	0.48	
Academic degree	n.a.	1-6	2.60	1.30	2.57	1.35	0.21	0.02	
Academic results	n.a.	1-5	3.03	1.03	3.05	1.16	-1.14	0.02	
Alcohol use	n.a.	0-4	1.27	1.16	1.35	1.22	-0.65	0.07	
Cannabis use	0.87	0-34	3.10	4.55	3.00	4.65	-0.20	0.02	

* $p < 0.05$

n.a.: not applicable

Table 2

Typology of 101 subjects with DEB based on the following traits: schizotypal, autistic, cyclothymic, borderline and obsessional;
Cluster comparison using ANOVA and post-hoc test

<i>Variables</i>	Cluster:						
	LT <i>n</i> =35 34.65%		SBC <i>n</i> =33 32.67%		HT <i>n</i> =33 32.67%		<i>F</i>
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	
Schizotypal traits	6.03 ^{ab}	3.34	11.33 ^a	3.40	13.06 ^b	3.72	37.85*
Autistic traits	2.54 ^a	1.44	2.33 ^b	0.96	5.09 ^{ab}	1.18	53.06*
Cyclothymic traits	3.34 ^{ab}	1.91	8.85 ^a	1.86	7.94 ^b	2.18	75.80*
Borderline traits	28.49 ^{ab}	8.20	43.30 ^a	9.372	47.09 ^b	11.06	36.03*
Obsessional traits	3.43 ^a	1.61	3.88 ^b	1.53	4.79 ^{ab}	1.14	7.71*
Eating behavior	36.71	14.29	31.48	10.43	34.12	12.22	1.50
Depression	12.06 ^{ab}	3.11	15.91 ^a	3.51	17.00 ^b	2.87	22.89*
Suicidal ideas	0.51 ^{ab}	1.46	1.91 ^a	2.58	2.82 ^b	2.45	9.41*
Obsess.-compulsive symptoms	13.57 ^{ab}	8.48	25.85 ^a	11.451	29.24 ^b	9.59	23.81*
State-trait anger	17.69 ^{ab}	5.30	21.58 ^a	5.51	22.67 ^b	5.08	8.36*
Alcohol	1.23	1.21	1.61	1.25	1.21	1.19	1.11
Cannabis	1.40	2.88	3.00	5.30	2.79	6.78	0.96
Academic degree	3.06 ^a	1.23	2.36	1.38	2.24 ^a	1.32	3.82*
Academic results	3.40	1.03	2.79	1.11	2.94	1.27	2.67

Note: Means with the same superscript letter are significantly different from each other (Tukey post-hoc test)

* $p < 0.05$




Figure 1. Classification of individuals from the DEB group: Three-cluster solution based on scores for the dimensions indicated on the x-axis.

SBC : cluster with high level of Schizotypal, Borderline and Cyclothymic traits

HT : High Traits cluster

LT : Low Traits cluster

Author disclosures

- **Role of funding sources:** This study received no funding.
- **Contributors:** PR and HC designed the study and conducted the statistical analyses. PR wrote the first draft of the manuscript. HC and TM contributed to and have approved the final manuscript.
- **Conflict of Interest:** The authors report no conflict of interest.
- **Acknowledgments:** PR is supported by Inserm.