



Nutrition and mortality in the elderly over 10 years of follow-up: the Three-City study

Flavie Letois, Thibault Mura, Jacqueline Scali, Laure-Anne Gutierrez,
Catherine Feart, Claudine Berr

► To cite this version:

Flavie Letois, Thibault Mura, Jacqueline Scali, Laure-Anne Gutierrez, Catherine Feart, et al.. Nutrition and mortality in the elderly over 10 years of follow-up: the Three-City study: Nutrition and mortality. British Journal of Nutrition, 2016, 116 (5), pp.882-9. 10.1017/S000711451600266X . inserm-01355378

HAL Id: inserm-01355378

<https://inserm.hal.science/inserm-01355378>

Submitted on 23 Aug 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BRITISH JOURNAL
of **NUTRITION**



CAMBRIDGE
UNIVERSITY PRESS

Nutrition and mortality in the elderly over 10 years of follow-up, the 3-city study

Journal:	<i>British Journal of Nutrition</i>
Manuscript ID	BJN-RA-15-1368.R2
Manuscript Type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Letois, Flavie; Centre Hospitalier Regional Universitaire de Montpellier, CIC Mura, Thibault; Centre Hospitalier Regional Universitaire de Montpellier, DIM Scali, Jacqueline; Inserm U1061, Affective disorders: bio-environmental risk and resilience Gutierrez, Laure-Anne; Inserm U1061, Cognitive ageing Féart, Catherine; INSERM U897, Nutrition Berr, Claudine; INSERM U1061, Cognitive ageing
Keywords:	mortality, dietary habits, fruits and vegetables, olive oil, fish
Subject Category:	Dietary Surveys and Nutritional Epidemiology

SCHOLARONE™
Manuscripts

Nutrition and mortality in the elderly over 10 years of follow-up, the 3-city study

Flavie Letois^{ab}, Thibault Mura^{a,c,d}, Jacqueline Scali^{c,d}, Laure-Anne Gutierrez^{a,c,d}, Catherine Féart^{e,f}, Claudine Berr^{a,c,d},

^aCHRU Montpellier, F-34000 France

^bINSERM, CIC 1411, F-34000 Montpellier, France

^cINSERM U1061, La Colombière Hospital, F-34093 Montpellier, France.

^dMontpellier University, F-34090 Montpellier, France

^eINSERM, ISPED, Centre INSERM U1219-Bordeaux Population Health, F-33000 Bordeaux, France.

^fUniv. Bordeaux, F- 33000 Bordeaux, France

Corresponding author

Dr Claudine BERR,
Inserm U1061, Hôpital La Colombière, F-34093 Montpellier Cedex 5, France.
Phone: 33 (0)4 99 61 45 66; Fax: 33 (0)4 99 61 45 79.
@address: claudine.berr@inserm.fr

Short title: Nutrition and mortality in the elderly

Key Words: mortality, dietary habits, fruits and vegetables, olive oil, regular fish consumption.

Abstract

In the last 20 years, many prospective cohort studies assessed the relationships between food consumption and mortality. Result interpretation is mainly hindered by the limited adjustment for confounders and, to a lesser extent, the small sample sizes. The aim of this study was to investigate the association between dietary habits and all-cause mortality in a multicentre prospective cohort that included non-institutionalized community-based elderly individuals (3-City Study). A brief Food Frequency Questionnaire was administered at baseline. Hazard ratios (HRs) and 95% confidence intervals (95%CI) for all-cause mortality were estimated relative to the consumption frequency of several food groups, using Cox proportional hazards models adjusted for sex, centre, socio-demographic characteristics and health status indicators. Among the 8937 participants (mean age: 74.2 y, 60.7% women), 2016 deaths were recorded during an average follow-up of nine years. The risk of death was significantly lower among subjects with the highest fruit and vegetable consumption (HR=0.90, 95%CI=0.82-0.99, $p=0.03$) and with regular fish consumption (HR=0.89, 95%CI=0.81-0.97, $p=0.01$). The benefit of olive oil use was found only in women (moderate olive oil use: HR=0.80, 95%CI=0.68-0.94, $p=0.007$; intensive use: HR=0.72, 95%CI=0.60-0.85, $p=0.0002$). Conversely, daily meat consumption increased the mortality risk (HR=1.12, 95%CI=1.01-1.24, $p=0.03$). No association was found between risk of death and diet diversity, use of various fats. These findings suggest that fruits/vegetables, olive oil and regular fish consumption have a beneficial effect on the risk of death, independently of the sociodemographic features and the number of medical conditions.

Nutrition and mortality in the elderly over 10 years of follow-up, the 3-city study

Introduction

The number of elderly people is progressively increasing. Besides medical progress, the life expectancy increase could be linked also to environmental and behavioural factors, some of which can be modified to achieve beneficial effects for health, particularly dietary habits ⁽¹⁾. The importance of diet has led public health policy-makers to develop specific prevention programmes. Moreover, large observational studies have examined the relationship between diet and all-cause mortality in elderly subjects using various food habit classifications that focus, for example, on specific foods or food consumption groups, or dietary patterns that combine multiple features. They showed that some dietary patterns (eating habits, including several food groups), such as the Mediterranean diet ^(2; 3; 4; 5; 6; 7; 8; 9; 10; 11; 12; 13), or “healthy” dietary groups defined by high consumption of fruits and vegetables and fish ^(14; 15; 16; 17), or greater consumption of olive oil with raw vegetables ⁽¹⁸⁾, have a protective effect. Similarly, some studies on consumption frequency found that high intake of fruits and/or vegetables ^(19; 20; 21), fish ^(20; 22) or coffee ^(23; 24; 25) has a beneficial effect on survival. However, few of these studies have considered the most relevant adjustment factors. Indeed, dietary habits and mortality are both influenced by many factors not only at the individual level (age, sex, health status, socio-economic, cultural and genetic factors), but also at the population level (for instance, food availability and accessibility in terms of cost).

Another way to assess the effect of diet on health is to consider the concept of food diversity. The advantage of studying the diversity of diets and of used fats compared with monotonous diets is that this allows evaluating the consumption of the most essential micronutrients and of different types of fatty acids (FA). This concept is part of the French National Health and Nutrition Programme ⁽²⁶⁾. Indeed, a varied diet, defined by the daily consumption of at least one unit of each essential food group, has been associated with a lower mortality risk ^(27; 28). However, few studies have focused on this approach.

Therefore, our main objective was to examine the associations between dietary habits and mortality in an elderly population. The data collected by the 10-year follow-up study of the large French population-based Three-City (3C) cohort allowed investigating whether some dietary habits (consumption of fruits, vegetables, olive oil, fish and caffeine, diet diversity and diversity of used

fats) are associated with reduced all-cause mortality in people aged >65 years, independently of previously identified major confounders.

Methods

Data were extracted from the database of the 3C study, a prospective cohort study of vascular risk factors of dementia⁽²⁹⁾. The present study was conducted according to the Declaration of Helsinki guidelines. The 3C study protocol was approved by the Consultative Committee for the Protection of Persons participating in Biomedical Research of the Kremlin-Bicêtre University Hospital (Paris, France). All participants signed a written informed consent.

Participants

A sample of community dwellers aged 65 years and older was selected in 1999–2000 from the electoral rolls of Bordeaux (South-West of France), Montpellier (South-East of France) and Dijon (North-East of France). To be eligible for recruitment into the study⁽²⁹⁾, persons had to be (1) living in these cities or their suburbs and registered on the electoral rolls, (2) aged 65 years and over, and (3) not institutionalized. The cohort size by center was set at 2,500 in Bordeaux, 5,000 in Dijon and 2,500 in Montpellier. A personal letter was sent to each potential subject inviting them to participate and including a brief description of the study protocol and an acceptance/refusal form. The spouse/partner was also invited to participate in the study if he/she met eligibility criteria. As electoral rolls are occasionally inaccurate (persons who have moved may still be registered at their previous address), no further correspondence was sent to persons who did not respond to the first letter, but an attempt was made to contact them by telephone. Twenty-four percent of the eligible persons selected on the electoral rolls ($n = 34,922$) could not be reached; among those contacted, the acceptance rate was 37%. A total of 9,693 persons were included; 7 persons aged less than 65 years were subsequently excluded. Participants who had subsequently refused to participate in the baseline medical interview ($n = 392$) were excluded from all analyses leading to an original sample of 9294 community dwellers.

The participants were administered standardized questionnaires and underwent clinical examinations at baseline and after 2, 4, 8 and 10 years. Baseline data were collected by standardized face-to-face interview. They included information on sociodemographic features, lifestyle (including a brief Food Frequency Questionnaire), current symptoms and complaints, medical

history, blood pressure, past and present consumption of tobacco, alcohol and drug use, anthropometric data, neuropsychological testing and blood sampling. For the current study, the following participants were excluded: 132 (1.4%) subjects with missing data for at least one dietary variable, 214 (2.3%) people with a diagnosis of dementia at baseline, and 11 (0.12%) individuals with no available vital statistics data. The analysis was thus carried out using a sample of 8,937 participants with a mean follow-up of 8.85 years.

Baseline dietary assessment

The brief Food Frequency Questionnaire (FFQ) was administered at baseline to assess the participants' dietary habits concerning nine broad food categories: (1) meat and poultry, (2) fish (including seafood), (3) eggs, (4) milk and dairy products, (5) cereals (including bread and starches), (6) raw fruits, (7) raw vegetables, (8) cooked fruits or vegetables, and (9) pulses. Consumption frequency was classified as follows: never, less than once per week, once per week, 2–3 times per week, 4–6 times per week and daily. The dietary habits of the whole sample have been described previously⁽³⁰⁾. Participants were also invited to indicate the dietary fats used at least once per week for dressing, cooking, or spreading among those included in the following list: butter, margarine, maize oil, peanut oil, sunflower or grapeseed oil, olive oil, blended oil, duck or goose fat, lard, 'Vegetaline'© (mainly saturated fat), colza oil, walnut oil, soybean oil. Caffeine consumption was calculated by multiplying the number of tea and coffee cups drunk per day by their estimated caffeine content (about 50mg/cup for tea and 100mg/cup for coffee, respectively)⁽³¹⁾.

Fruits/vegetables, fish, meat and olive oil were the four main food consumption groups of interest that were studied as categorical variables. Their consumption thresholds were based on the French National Nutrition and Health Programme guidelines, when applicable to the available data⁽²⁶⁾. Cooked fruits and vegetables, olive oil and caffeine consumption were investigated using previously defined categories^(31; 32; 33). Dietary habits included the intake of fish/seafood (less than twice per week versus more than twice a week), fruits and vegetables (less than one fruit and one vegetable, cooked or raw, per day versus at least one fruit and one vegetable, cooked or raw, per day, and less versus more than 4–6 servings of cooked fruits or vegetables per week), meat (less versus more than once per day), caffeine (<250mg/day, 250–375mg/day, >375mg/day). Three categories of olive oil consumption were defined: 'no use', 'moderate use' (for cooking or dressing alone), or 'intensive use' (for cooking and for dressing)^(32; 34). To calculate the Diet Diversity Score (DDS) (from 0 to 5), one point was assigned for each of the following food categories consumed at

least once per day: dairy products, meat, cereals, fruits and vegetables. Low diversity was defined by a DDS <3 and high diversity by a DDS ≥ 4 ⁽³⁵⁾. Fat diversity was evaluated as ≤ 3 different fats versus >3 fats. The threshold of three different fats corresponds to the median value of various fats used by the population included in the 3C cohort.

Baseline covariates

Socio-demographic information recorded at baseline included age, sex, centre, educational level (no or primary/middle/high school/university), occupation (white collar, employee, blue collar, housewife) and monthly income level ($<1,500$, $>1,500$ euros). Health behaviours were assessed based on smoking status (no, 0-10, 10-30 or >30 packets per year), intake of alcoholic beverages (0-24g/day, >24 g/day for women; 0-36g/day, >36 g/day for men).

Blood pressure and anthropometric data were measured in standardized conditions. Health-related variables included body mass index (BMI: <27 and ≥ 27 kg/m²) ⁽³⁶⁾, hypertension (defined as a systolic blood pressure >140 mm Hg or diastolic blood pressure >90 mm Hg, or use of antihypertensive drugs) ⁽³⁷⁾, diabetes (fasting blood glucose ≥ 7.0 mmol/l, or taking an anti-diabetic treatment), hypercholesterolemia (yes/no, self-reported) and the number of drugs (0, 1-4, ≥ 5). History of cardiovascular diseases (yes/no) and other chronic diseases (chronic respiratory disease, cancer, Parkinson's disease or hypothyroidism) were reported. Disability was estimated using the Instrumental Activities of Daily Living scale (IADL; yes (score >0 /no,). Depressive mood was assessed using the Center for Epidemiological Studies-Depression scale (CES-D) and was defined by a score <17 for men and <23 for women. Self-rated health was grouped in three categories (poor, average, fair) and self-assessment of correct diet in two classes (no /yes).

Physical activity was assessed differently in the three centres (self-reported or not and with different accuracy levels). Therefore, a common binary variable was considered for the three centres (none or low/regular). Regular physical activity was defined as doing some sport regularly or having at least one hour of leisure or household activity per day ⁽³⁸⁾. Given the important amount of missing data for this variable (1115 missing data for the Bordeaux centre, 11% of the whole sample), it was used only in sensitivity analyses.

Vital status

Mortality was ascertained from the civil registry by systematic request for all subjects not included in follow-up visits. The date of death was defined as the date of event and the date of the last follow-up or phone contact for the 10-year follow-up as the date of censoring. Follow-up included

the precise assessment of all causes of death that were coded according to the tenth revision of the International Classification of Diseases (ICD-10)⁽³⁹⁾. Mortality from cardiovascular disease (CVD) (ICD-10: I) and cancer (ICD-10: C), the two leading causes of death in aged populations, were considered for the analyses⁽³⁹⁾.

Statistical Analyses

Survival analysis was carried out using the Kaplan Meier method. Comparisons of baseline characteristics and dietary habits were performed with the log-rank test.

The associations between dietary habits, diet diversity and mortality risk were determined by using Cox proportional hazards regression with delayed entry, with age (in years) used as the time axis and left truncation at the age of study entry. The results of the survival analysis were expressed as hazard ratios (HRs) with 95% confidence intervals (CI).

In these analyses, the adjustments for covariates were performed in three steps. Univariate analysis was the basic model to test associations between dietary features and mortality (crude model). The second model (model 1) was adjusted for sex and study centre, educational level, occupation and income. Health behaviours and health factors, chosen based on literature data, were then added in the third model (model 2). Interactions between the different dietary habits and covariates included in model 2 were tested and stratified analyses were carried out accordingly.

For missing data, multiple imputations were used by generating five copies of the original dataset in which the missing values for the covariates considered in the analysis were replaced by values generated according to the Markov Chain Monte Carlo (MCMC) method and using the PROC MI procedure. Each imputed dataset was analysed as if it were complete to calculate the mean HRs and CI with the PROC MIANALYZE procedure⁽⁴⁰⁾. Imputation were performed for covariates included in analyses: education (missing data 0.17%), income (6.13%), occupation (0.29%), smoking (1.63%), alcohol (1.51%), history of cardiovascular disease (0.08%), , depression (1.28%), diabetes (5.65%), hypertension (0.06%), hypercholesterolemia (1.1%), dependence (0.69%)self-rated health (0.57%), number of chronic diseases (0.09%), self rated diet quality (2.38%).

Analyses were carried out using the SAS software (version 9.2).

Results

The baseline characteristics and dietary habits of the participants ($n=8937$) according to their vital status (dead/alive) at the end of the 10-year follow-up are reported in Table 1 and Table 2, respectively. Participants who died ($n=2016$) were significantly older at baseline than those still alive (mean age: 77 vs 73, respectively) and more often men, heavy smoker or alcohol consumers. They also had more often a history of cardiovascular or chronic diseases, more vascular risk factors and poor self-rated health. In these crude comparisons, mortality risk was higher among individuals with low income and blue collar workers. Conversely, survival was not influenced by the education level in this highly educated population. Daily consumption of fruits and vegetables, regular consumption of cooked vegetables/fruits, intensive olive oil use, wide fat diversity and bi-weekly fish consumption were more frequently reported by subjects still alive at the end of the 10-year follow-up. Caffeine consumption and DDS were not associated with survival.

Then, the associations between the 10-year mortality risk and dietary habits or diet diversity (at baseline) were assessed after multiple imputations for missing data (Table 3 and Table 4, respectively). In crude analyses, various well-known healthy food habits were significantly associated with better survival: eating at least one fruit and one vegetable (raw or cooked) per day, consumption of at least four cooked fruit/vegetable servings per week, fish at least twice per week, meat less than once per day, greater diet diversity and diversity in fat use.

In the partly (only sociodemographic covariates) and fully (sociodemographic and health-related covariates) adjusted models, consuming at least one fruit and vegetable per day and consuming more than four cooked fruit/vegetable servings per week remained significantly associated with a better survival compared with lower levels of consumption ($HR=0.90$, 95% $CI=0.82-0.99$, $p=0.03$ and $HR=0.80$, 95% $CI=0.70-0.90$, $p=0.0005$, respectively, in the fully adjusted models). Fish intake at least twice per week was also significantly associated with better survival ($HR=0.89$, 95% $CI=0.81-0.97$, $p=0.01$), while consuming meat more than once per day had a deleterious effect on survival in the fully adjusted models ($HR=1.12$, 95% $CI=1.01-1.24$, $p=0.03$) (Table 3).

For the two indicators of diet diversity (DDS and diversity in fats used), the significant associations observed in crude and partly adjusted analyses were no longer significant after adjustment also for health-related variables (Table 4).

229 As the interaction between sex and consumption of olive oil was significant, analyses were
230 stratified by sex (Supplementary Table 1). The use of olive oil was inversely correlated with the
231 mortality risk after adjustment for all covariates, but only in women (moderate olive oil use:
232 HR=0.80, 95%CI=0.68-0.94, p=0.007; intensive use: HR=0.72, 95%CI=0.60-0.85, p=0.0002).

233 Furthermore, sensitivity analyses were performed on the 5322 individuals with available
234 data on physical activity. In this sub-sample, the consumption of at least 4-6 servings of cooked
235 fruits or vegetables per week was again associated with a lower mortality risk at the end of the 10-
236 year follow-up, independently of physical activity (HR=0.75, 95% CI=0.62-0.90, p=0.002). No
237 association was observed between mortality risk and meat, fish or olive oil consumption
238 (Supplementary Table 2) or diet diversity (Supplementary Table 3).

239 Finally, there was no significant association between dietary habits/diversity and the main
240 causes of death (389 due to CVD and 542 to cancer) after controlling for sex, centre, educational
241 level, income and occupation (data not shown).

242

Discussion

The present study based on the 10-year follow-up of a French cohort of people aged 65 or older suggests that healthy dietary habits, such as daily consumption of fruits and vegetables, eating fish twice per week and regular use of olive oil (this only in women), are linked to better survival, independently of sociodemographic, health-related and lifestyle variables. Overall, higher diet diversity also is significantly associated with lower mortality risk.

A major difficulty when assessing food habits concerns the heterogeneity of the collected data. Indeed, each country has its consumption standards, tailored to cultural habits and the local availability of food resources. Moreover, the methods used to collect such data also are not homogeneous and consequently results are difficult to compare and generalize from one country to another. Nevertheless, the various eating habits can be standardized in dietary patterns (such as composite scores or consumption patterns of some preferred foods) or by quantitative measurements of food types. Studies on diet quality or adherence to a particular diet are used to determine the most beneficial approach; however, in the case of composite scores, the proper role of each score component cannot be identified. In the present study where food habits were evaluated with a short FFQ, we examined food groups that have been found to be associated with longer lifespan in previous studies using these different approaches. Our results are in agreement with the French National Nutrition and Health Programme guidelines that include the promotion of daily consumption of fruit and vegetables⁽²⁶⁾.

We observed a significant association between high consumption of fruits and vegetables and survival after adjusting for major confounders. This result is consistent with earlier studies^(19; 20; 21). In the EPIC study, a better adherence to a plant-based diet was associated with a reduced mortality risk in Southern Europe, but not in the UK nor in Germany, after controlling for all known risk factors⁽¹⁵⁾. Similarly, the “Olive oil and salad” pattern was associated with longevity in an Italian elderly cohort⁽¹⁸⁾.

Fruits and vegetables, which were not separate entities in our FFQ, are the main source of antioxidants that are the basis of the free radical theory of ageing⁽⁴¹⁾. According to this theory, a balance between free radicals and antioxidants increases life expectancy. Some authors hypothesize that plant foods promote the activation of immune functions or have protective properties^(16; 17). A recent literature review highlighted the effect of high concentration of polyphenols, carotenoids, folic acid and vitamin C on mortality⁽¹³⁾. High consumption of flavonoids, found mainly in fruits, could be associated with reduced mortality risk⁽⁴²⁾. The beneficial effects of fruits and vegetables

could be also explained by their fibre content that modulates LDL cholesterol level, insulin sensitivity and blood pressure ⁽⁴³⁾, all conditions associated with increased mortality risk. This hypothesis is supported by the PREDIMED study that demonstrated a significant association between lower risk of death and baseline higher fibre intake and fruit consumption ⁽⁴⁴⁾.

Our results are in agreement with the few studies that investigated fish consumption and risk of death ^(20; 22). A Japanese study found that high consumption of vegetables and fish in subjects older than 75 years is associated with better survival ⁽¹⁷⁾. Most of the published results on the positive association between fish and lower mortality have been carried out in countries with traditional high fish consumption (Japan and Scandinavia). Other studies included fish consumption in the diet scores (e.g., the Mediterranean Diet Score) or in “healthy” diet patterns ^(14; 16; 45) with contrasting results. Hoffman et al. did not find any significant association between these dietary patterns and mortality⁽⁴⁵⁾. Conversely, Anderson et al⁽¹⁶⁾ observed a positive link between survival and a healthy dietary pattern (consumption of low-fat dairy products, fruit, whole grains, poultry, fish and vegetables). The protective effect of fish consumption on health has been linked to the anti-inflammatory effects of the essential omega-3 fatty acids ^(13; 22).

Olive oil is the main source of fat in the traditional Mediterranean diet. Our study reported a beneficial effect only among women. This finding is in agreement with the findings in an elderly British cohort ⁽⁷⁾ where the Mediterranean style dietary pattern was associated with reduced mortality only in women. This sex-specific effect could be explained by the women’s longer life expectancy. Many studies support the beneficial effect of the Mediterranean diet on mortality ^(2; 3; 4; 5; 6; 7; 8; 12; 46). Masala et al. observed that the ‘Olive Oil & Salad’ dietary pattern (high consumption of olive oil) is inversely associated with all-cause mortality in the elderly ⁽¹⁸⁾. The PREDIMED study found that higher consumption of olive oil was associated with a specific reduction in the cardiovascular mortality risk in an elderly Mediterranean population ⁽⁴⁷⁾. The beneficial effect of olive oil could be explained by its high concentration in **mono**-unsaturated fatty acids and phenolic compounds that improves endothelium function and reduces oxidative stress, thereby promoting healthy aging and longevity ⁽¹³⁾.

On the other hand, we found that daily consumption of meat, as a broad category, had a negative effect on survival. Most previous studies showing a deleterious effect of meat consumption on lifespan ^(17; 45) also found that high meat consumption patterns were often associated with unhealthy dietary habits, such as high levels of fat consumption. However, in other studies, red meat consumption was not associated with increased mortality risk after correction for confounding

factors⁽¹⁶⁾. Comparisons with previous findings are limited by the different definitions of meat consumption groups. In some works red meat with high content of fatty acids was considered on its own, whereas poultry or lean meat was not included in the analysis. Anderson et al. suggested that high consumption of animal foods could increase the mortality risk, only if meat replaces protective plant foods in the diet⁽¹⁶⁾. The link between red meat and dietary fat could affect the lipid and lipoprotein profile and thus the cardiovascular disease risk⁽⁴⁵⁾.

The association between diet diversity and mortality did not remain significant after adjustment for confounders, unlike in other studies^(27; 28). However, comparisons are difficult because these studies used different methods of food collections and different food groups.

The data from our study should be interpreted in the light of the following limitations. First, the 3C sample is not representative of the French population. Indeed, compared with the whole French population, this cohort included more urban dwellers, in better health and living together⁽³⁰⁾. Therefore, extrapolation of our results to the general population of elderly should be made with caution. The FFQ did not record quantitative data (portions, grams), but only frequencies of consumption. Low frequency consumption does not necessarily mean lower energy intake, but simply suggests a lack of diversity in eating habits. Moreover, the FFQ did not allow calculating the total energy intake, as recommended to better control for confounding factors and to reduce extraneous variation⁽⁴⁸⁾. In a previous article exclusively on the population from the Bordeaux centre (N=1660) of the 3C study, daily energy intake was calculated in a sub-sample by using the 24-hour dietary recall technique. In this sub-sample, daily energy intake was not affected by the frequency of consumption of fish, or of fruits and vegetable (frequent vs less frequent), or by the use of omega-3 or omega-6 rich oils (regular vs non-regular)⁽⁴⁹⁾. Moreover, the baseline energy intake among the Bordeaux sub-sample was not significantly different between those who were dead (N=375) or still alive (N=1422) at the end of the 10-year follow-up (data not shown). This suggests that the energy intake is not a major confounder in the present analyses. According to the literature, mainly studies with the most accurate data on food consumption (in daily amount)^(3; 7; 15; 16; 18; 20; 21; 22; 43; 46) found significant associations between dietary patterns and mortality. Our data did not allow this level of detail. However, our results confirm the benefit of several food categories on survival. Moreover, in our study, the assessment of dietary intake covered only broad food groups. This limitation is important for the interpretation of the association between meat and survival, because the meat group included both white meat (lean, healthy) and red meat (fat, deleterious). The limitations the food questionnaire (FFQ) used in this study (few items, absence of validation, only baseline administration) could have led to non-differential misclassification bias.

However, previous studies using this questionnaire reported innovative results that are in agreement with the literature or have been successively confirmed by other independent analyses^(30; 34; 49; 50). Finally in our sensitivity analysis including physical activity, only the association with fruits and vegetables remains significant. But the variable physical activity did not rely on a detailed physical activities questionnaire and these analyzes are conducted in a limited sample (n=5273) reducing statistical power. The interpretation of this result is therefore limited.

Strengths of this study include the prospective population-based study design, with a large sample size, the completeness of the collected dietary data (data provided by more than 99% of subjects, possibly due to the use of a simple and short questionnaire) and the low number of drop-outs (0.1%). Many variables were investigated as potential confounders, such as socio-demographic factors, health behaviours, BMI, drug use, self-rated health status and diet. The FFQ used in the 3C study has already allowed highlighting many interesting associations between diet habits and diseases^(31; 33; 34; 49; 50; 51; 52; 53). Chrysohoou et al. insisted on the importance of taking into account socio-economic factors when considering the relationship between food habits and life expectancy, as healthy habits are associated with financial and educational status⁽¹³⁾. In our study, we considered simultaneously education, occupation and income.

In conclusion, this study on the associations between dietary habits and mortality in a large population-based elderly cohort brings reliable evidence that could be used for developing nutritional programmes. Specifically, a diet with a daily intake of fruits and vegetables, regular consumption of fish, as recommended by the French Nutrition and Health Programme⁽²⁶⁾, and regular use of olive oil promotes longevity among an elderly community-living population. Diet quality is an important component of a healthy lifestyle that has beneficial effects on survival⁽¹⁾. Additional studies should investigate the association between survival, diet and healthy behaviours, but this does not limit the importance of this simple public health message.

Acknowledgments

The 3C Study was conducted under a partnership agreement between the Institut National de la Santé et de la Recherche Médicale (INSERM), the Victor Segalen – Bordeaux II University and the Sanofi-Synthelabo Company. The Fondation pour la Recherche Médicale supported the preparation and initiation of the study. The 3C Study is also supported by the Caisse Nationale Maladie des Travailleurs Salariés, Direction Générale de la Santé, Conseils Régionaux of Aquitaine, Languedoc-Roussillon and Bourgogne, Fondation de France, Ministry of Research-INSERM Programme ‘Cohortes et collections de données biologiques’, Mutuelle Générale de l’Education Nationale, Institut de la longévité, Conseil Général de la Côte d’Or, Agence Nationale de la Recherche ANR PNRA 2006 and Longvie 2007 and Fonds de coopération scientifique Alzheimer (FCS 2009-2012).

Conflict of interest

None for F Letois, T Mura, J Scali, LA Gutierrez, C Berr.
C Féart received fees for conferences from Danone Research and Nutricia.

Authorship

F Letois: formulation of the research question, statistical analysis and interpretation of the data; drafting the manuscript.
T Mura; drafting/revising the manuscript
J Scali drafting/revising the manuscript
LA Gutierrez: statistical analysis, revising the manuscript
C Féart: drafting/revising the manuscript
C. Berr: data acquisition; analysis and interpretation of the data; study design; study supervision; drafting/revising the manuscript.

389 **Table 1 - Baseline socio-demographic and medical characteristics of the 3C cohort (N=8937)**
390 **subdivided according to their vital status (dead or alive) at the end of the 10-year follow-up**
391

	Alive N=6921 (77%)		Dead N=2016 (23%)	
	N (%)	mean (sd)	N (%)	mean (sd)
Age (years)		73 (4.9)		77 (6.1)
Women	4507 (65%)		941 (47%)	
Education				
No or primary	1747 (25%)		531(26%)	
Secondary	2474 (36%)		714 (36%)	
High school	1403 (20%)		389 (19%)	
University	1287 (19%)		377 (19%)	
Monthly income higher than 1500€	4148 (64%)		1153 (62%)	
Occupation				
White collar	2790 (40%)		815 (41%)	
Employee	2029 (29%)		535 (27%)	
Blue collar	1395 (20%)		496 (25%)	
Housewife	687 (10%)		164 (8%)	
Smoking				
Never	4436 (65%)		1047 (53%)	
0-10 packets/year	902 (13%)		274 (14%)	
10-29 packets/year	898 (13%)		308 (15%)	
> 30 packets/year	586 (9%)		340 (17%)	

Alcohol (≥ 24 g/d for women; ≥ 36 g/d for men)	643 (9%)	237 (12%)
History of cardiovascular diseases ^a	1752 (25%)	878 (44%)
BMI < 27 kg/m ²	4668 (68%)	1280 (65%)
Depression	879 (13%)	333 (17%)
Diabetes	568 (9%)	287 (15%)
Hypertension	5235 (76%)	1684 (84%)
Hypercholesterolemia	2405 (35%)	553 (28%)
Dependence ^b	431 (6%)	398 (20%)
Low/regular physical activity	2119 (34%)	373 (22%)
Self-rated health		
Poor	276 (4%)	182 (9%)
Fair	2337 (34%)	828 (41%)
Good	4275 (62%)	988 (49%)
Number of drugs		
None	651 (9%)	103 (5%)
1-4	3433 (50%)	752 (37%)
≥ 5	2837 (41%)	1161 (58%)
Number of chronic diseases ^c		
None	5850 (85%)	1567 (78%)
1	975 (14%)	396 (20%)
2 or more	91(1%)	50 (2%)

^a Coronary heart disease was reported by 671 (9.7%) of subjects alive and 398 (20%) of subjects dead at the end of the follow-up.

^b Assessed with the IADL scale (at least 1 impairment).

^c Chronic diseases included mainly respiratory diseases and dyspnoea (n=477, 5.8% of people still alive; n=285, 14.1% of subjects dead at the end of the follow-up), Parkinson’s disease (n=46, 0.66% of alive; n=59, 2.9% of dead subjects), and hyperthyroidism (n=6, 18.9% of alive; n=143, 7.1% of dead).

For Review Only

Table 2 - Baseline food habits of the 3C cohort (N=8937) subdivided according to their vital status (alive or dead) at the end of the 10-year follow-up

	Alive N=6921 (77%)	Dead N=2016 (23%)
	N (%)	N (%)
At least 1 fruit and 1 vegetable, cooked or raw /day	2782 (40%)	663 (33%)
Cooked fruits/vegetables: ≥ 4 -6 servings/week	6162 (89%)	1715 (85%)
Meat: ≥ 1 serving/day	1689 (24%)	558 (28%)
Fish: ≥ 2 servings/week	3569 (52%)	923 (46%)
Diversity diet score ≥ 4	4986 (72%)	1371 (68%)
Olive oil use		
None	1543 (23%)	644 (32%)
Moderate	2768 (40%)	755 (37%)
Intensive	2610 (38%)	617 (31%)
Different fats > 3	1523 (22%)	345 (17%)
Caffeine (mg/day)		
< 250	4781 (69%)	1489 (74%)
250-375	1441 (21%)	370 (18%)
> 375	699 (10%)	157 (8%)
Self- rated diet quality		
Bad	589 (9%)	171 (9%)
Good	6174 (91%)	1790 (91%)

403

404

405

406

407

408

409

410

	N	Crude	(95% CI)	p	Model 1 ^a	(95% CI)	p	Model 2 ^b	(95% CI)	p
At least 1 fruit and 1 vegetable, cooked or raw/day	8937	0.82	[0.75-0.90]	***	0.84	[0.77-0.93]	***	0.90	[0.82-0.99]	*
Cooked fruits or vegetables: ≥4-6/ week	8937	0.73	[0.65-0.83]	***	0.78	[0.69-0.89]	***	0.80	[0.70-0.90]	***
Meat: ≥1 serving/day	8937	1.22	[1.11-1.34]	***	1.16	[1.05-1.28]	**	1.12	[1.01-1.24]	*
Fish: ≥2 servings/week	8937	0.83	[0.76-0.91]	***	0.87	[0.80-0.96]	**	0.89	[0.81-0.97]	**

^aCox proportional model adjusted for sex, centre, education (no or primary/secondary/ high school/university), income (<1500 or >1500 euros/month), occupation (white collar/employee/blue collar/housewife)

^bCox proportional model adjusted for sex, centre, education, income, occupation and also smoking, alcohol consumption, history of cardiovascular diseases, BMI, depression, diabetes, hypertension, hypercholesterolemia, dependence (IADL scale), self-rated health, self-rated diet quality, number of drugs, number of chronic diseases.

*P<0.05; **P<0.01; ***P<0.001

411

412 **Table 4 - Association between 10-year overall mortality and diet diversity in the 3C elderly cohort (N=8937): crude and adjusted hazard ratios**
 413 **(HR) and 95% CI**

Diet Diversity	N	Crude	(95% CI)	p	Model 1 ^a	(95% CI)	p	Model 2 ^b	(95% CI)	p
Diversity diet score 4-5 vs 0-3	8937	0.89	[0.81-0.98]	*	0.90	[0.82-0.99]	*	0.94	[0.85-1.04]	ns
Various fats >3 vs ≤3	8937	0.84	[0.75-0.95]	**	0.86	[0.76-0.97]	*	0.90	[0.80-1.01]	ns

414 ^aCox proportional model adjusted for sex, centre, education (no or primary/secondary/ high school/university), income (<1500 or >1500

415 euros/monthly), occupation (white collar/employee/blue collar/housewife)

416 ^bCox proportional model adjusted for sex, centre, education, income, occupation and also smoking, alcohol consumption, history of cardiovascular

417 diseases, BMI, depression, diabetes, hypertension, hypercholesterolemia, dependence (IADL scale), self-rated health, self-rated diet quality, number of

418 drugs, number of chronic diseases

419 *P<0.05; **P<0.01; ***P<0.001; ns=not significant

References

1. King DE, Mainous AG, 3rd, Geesey ME (2007) Turning back the clock: adopting a healthy lifestyle in middle age. *Am J Med* **120**, 598-603.

2. Trichopoulou A, Kouris-Blazos A, Wahlqvist ML *et al.* (1995) Diet and overall survival in elderly people. In *BMJ*, 1995/12/02 ed., vol. 311, pp. 1457-1460.

3. Osler M, Schroll M (1997) Diet and mortality in a cohort of elderly people in a north European community. *Int J Epidemiol* **26**, 155-159.

4. Haveman-Nies A, de Groot L, Burema J *et al.* (2002) Dietary quality and lifestyle factors in relation to 10-year mortality in older Europeans: the SENECA study. *Am J Epidemiol* **156**, 962-968.

5. Knuops KT, de Groot LC, Kromhout D *et al.* (2004) Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. *JAMA* **292**, 1433-1439.

6. Sjogren P, Becker W, Warensjo E *et al.* (2010) Mediterranean and carbohydrate-restricted diets and mortality among elderly men: a cohort study in Sweden. *Am J Clin Nutr* **92**, 967-974.

7. Hamer M, McNaughton SA, Bates CJ *et al.* (2010) Dietary patterns, assessed from a weighed food record, and survival among elderly participants from the United Kingdom. *Eur J Clin Nutr* **64**, 853-861.

8. McNaughton SA, Bates CJ, Mishra GD (2012) Diet quality is associated with all-cause mortality in adults aged 65 years and older. *J Nutr* **142**, 320-325.

9. van den Brandt PA (2011) The impact of a Mediterranean diet and healthy lifestyle on premature mortality in men and women. *Am J Clin Nutr* **94**, 913-920.

10. Sofi F, Cesari F, Abbate R *et al.* (2008) Adherence to Mediterranean diet and health status: meta-analysis. *BMJ* **337**, a1344.

11. Roman B, Carta L, Martinez-Gonzalez MA *et al.* (2008) Effectiveness of the Mediterranean diet in the elderly. *Clin Interv Aging* **3**, 97-109.

12. Trichopoulou A, Orfanos P, Norat T *et al.* (2005) Modified Mediterranean diet and survival: EPIC-elderly prospective cohort study. *BMJ* **330**, 991.

13. Chrysoshoou C, Stefanadis C (2013) Longevity and diet. Myth or pragmatism? *Maturitas* **76**, 303-307.

14. Kumagai S, Shibata H, Watanabe S *et al.* (1999) Effect of food intake pattern on all-cause mortality in the community elderly: a 7-year longitudinal study. *J Nutr Health Aging* **3**, 29-33.

15. Bamia C, Trichopoulos D, Ferrari P *et al.* (2007) Dietary patterns and survival of older Europeans: the EPIC-Elderly Study (European Prospective Investigation into Cancer and Nutrition). *Public Health Nutr* **10**, 590-598.

16. Anderson AL, Harris TB, Tyllavsky FA *et al.* (2011) Dietary patterns and survival of older adults. *J Am Diet Assoc* **111**, 84-91.

17. Iimuro S, Yoshimura Y, Umegaki H *et al.* (2012) Dietary pattern and mortality in Japanese elderly patients with type 2 diabetes mellitus: does a vegetable- and fish-rich diet improve mortality? An explanatory study. *Geriatr Gerontol Int* **12 Suppl 1**, 59-67.

18. Masala G, Ceroti M, Pala V *et al.* (2007) A dietary pattern rich in olive oil and raw vegetables is associated with lower mortality in Italian elderly subjects. *Br J Nutr* **98**, 406-415.

19. Strandhagen E, Hansson PO, Bosaeus I *et al.* (2000) High fruit intake may reduce mortality among middle-aged and elderly men. The Study of Men Born in 1913. *Eur J Clin Nutr* **54**, 337-341.

20. Darmadi-Blackberry I, Wahlqvist ML, Kouris-Blazos A *et al.* (2004) Legumes: the most important dietary predictor of survival in older people of different ethnicities. *Asia Pac J Clin Nutr* **13**, 217-220.

21. Sahyoun NR, Jacques PF, Russell RM (1996) Carotenoids, vitamins C and E, and mortality in an elderly population. *Am J Epidemiol* **144**, 501-511.

22. Gillum RF, Mussolino M, Madans JH (2000) The relation between fish consumption, death from all causes, and incidence of coronary heart disease. the NHANES I Epidemiologic Follow-up Study. *J Clin Epidemiol* **53**, 237-244.

23. Fortes C, Forastiere F, Farchi S *et al.* (2000) Diet and overall survival in a cohort of very elderly people. *Epidemiology* **11**, 440-445.
24. Paganini-Hill A, Kawas CH, Corrada MM (2007) Non-alcoholic beverage and caffeine consumption and mortality: the Leisure World Cohort Study. *Prev Med* **44**, 305-310.
25. Freedman ND, Park Y, Abnet CC *et al.* (2012) Association of coffee drinking with total and cause-specific mortality. *N Engl J Med* **366**, 1891-1904.
26. (2011) French National Nutrition and Health Program 2011-2015 [Mo Health, editor].
27. Kant AK, Schatzkin A, Harris TB *et al.* (1993) Dietary diversity and subsequent mortality in the First National Health and Nutrition Examination Survey Epidemiologic Follow-up Study. *Am J Clin Nutr* **57**, 434-440.
28. Lee MS, Huang YC, Su HH *et al.* (2011) A simple food quality index predicts mortality in elderly Taiwanese. *J Nutr Health Aging* **15**, 815-821.
29. (2003) Vascular factors and risk of dementia: design of the Three-City Study and baseline characteristics of the study population. *Neuroepidemiology* **22**, 316-325.
30. Larrieu S, Letenneur L, Berr C *et al.* (2004) Sociodemographic differences in dietary habits in a population-based sample of elderly subjects: the 3C study. *J Nutr Health Aging* **8**, 497-502.
31. Ritchie K, Carriere I, de Mendonca A *et al.* (2007) The neuroprotective effects of caffeine: a prospective population study (the Three City Study). *Neurology* **69**, 536-545.
32. Berr C, Portet F, Carriere I *et al.* (2009) Olive oil and cognition: results from the three-city study. *Dement Geriatr Cogn Disord* **28**, 357-364.
33. Ritchie K, Artero S, Portet F *et al.* (2010) Caffeine, cognitive functioning, and white matter lesions in the elderly: establishing causality from epidemiological evidence. *J Alzheimers Dis* **20 Suppl 1**, S161-166.
34. Samieri C, Feart C, Proust-Lima C *et al.* (2011) Olive oil consumption, plasma oleic acid, and stroke incidence: the Three-City Study. *Neurology* **77**, 418-425.
35. Kant AK, Schatzkin A, Ziegler RG (1995) Dietary diversity and subsequent cause-specific mortality in the NHANES I epidemiologic follow-up study. *J Am Coll Nutr* **14**, 233-238.
36. Heiat A, Vaccarino V, Krumholz HM (2001) An evidence-based assessment of federal guidelines for overweight and obesity as they apply to elderly persons. *Arch Intern Med* **161**, 1194-1203.
37. Brindel P, Hanon O, Dartigues JF *et al.* (2006) Prevalence, awareness, treatment, and control of hypertension in the elderly: the Three City study. *J Hypertens* **24**, 51-58.
38. Feart C, Lorrain S, Ginder Coupez V *et al.* (2013) Adherence to a Mediterranean diet and risk of fractures in French older persons. *Osteoporos Int* **24**, 3031-3041.
39. Alperovitch A, Bertrand M, Jouglu E *et al.* (2009) Do we really know the cause of death of the very old? Comparison between official mortality statistics and cohort study classification. *Eur J Epidemiol* **24**, 669-675.
40. Yuan Y (2011) Multiple Imputation Using SAS Software. *Journal of Statistical Software* **45**.
41. Harman D (1972) Free radical theory of aging: dietary implications. *Am J Clin Nutr* **25**, 839-843.
42. Ivey KL, Hodgson JM, Croft KD *et al.* (2015) Flavonoid intake and all-cause mortality. *Am J Clin Nutr* **101**, 1012-1020.
43. Streppel MT, Ocke MC, Boshuizen HC *et al.* (2008) Dietary fiber intake in relation to coronary heart disease and all-cause mortality over 40 y: the Zutphen Study. *Am J Clin Nutr* **88**, 1119-1125.
44. Buil-Cosiales P, Zazpe I, Toledo E *et al.* (2014) Fiber intake and all-cause mortality in the Prevencion con Dieta Mediterranea (PREDIMED) study. *Am J Clin Nutr* **100**, 1498-1507.
45. Hoffmann K, Boeing H, Boffetta P *et al.* (2005) Comparison of two statistical approaches to predict all-cause mortality by dietary patterns in German elderly subjects. *Br J Nutr* **93**, 709-716.
46. Knuops KT, Groot de LC, Fidanza F *et al.* (2006) Comparison of three different dietary scores in relation to 10-year mortality in elderly European subjects: the HALE project. *Eur J Clin Nutr* **60**, 746-755.
47. Guasch-Ferre M, Hu FB, Martinez-Gonzalez MA *et al.* (2014) Olive oil intake and risk of cardiovascular disease and mortality in the PREDIMED Study. *BMC Med* **12**, 78.
48. Willett WC, Howe GR, Kushi LH (1997) Adjustment for total energy intake in epidemiologic studies. *Am J Clin Nutr* **65**, 1220S-1228S; discussion 1229S-1231S.

520 49. Barberger-Gateau P, Raffaitin C, Letenneur L *et al.* (2007) Dietary patterns and risk of dementia: the
521 Three-City cohort study. *Neurology* **69**, 1921-1930.

522 50. Berr C, Portet F, Carriere I *et al.* (2009) Olive Oil and Cognition: Results from the Three-City Study.
523 *Dementia and Geriatric Cognitive Disorders* **28**, 357-364.

524 51. Barberger-Gateau P, Jutand MA, Letenneur L *et al.* (2005) Correlates of regular fish consumption in
525 French elderly community dwellers: data from the Three-City study. *Eur J Clin Nutr* **59**, 817-825.

526 52. Feart C, Torres MJ, Samieri C *et al.* (2011) Adherence to a Mediterranean diet and plasma fatty acids:
527 data from the Bordeaux sample of the Three-City study. *Br J Nutr* **106**, 149-158.

528 53. Feart C, Samieri C, Rondeau V *et al.* (2009) Adherence to a Mediterranean diet, cognitive decline, and
529 risk of dementia. *JAMA* **302**, 638-648.

530

531

532

For Review Only

Supplementary Table 1 - Association between 10-year overall mortality and olive oil use stratified on sex in the elderly 3C Study cohort (8937): crude and adjusted hazard ratios (HR) and 95% CI

Olive oil use	N	Crude	(95% CI)	p	Model 1 ^a	(95% CI)	p	Model 2 ^b	(95% CI)	p
<i>All</i>	8937									
Moderate / None		0.79	[0.71-0.88]	***	0.84	[0.76-0.94]	**	0.89	[0.80-0.99]	*
Intensive / None		0.76	[0.68-0.84]	***	0.80	[0.71-0.90]	***	0.83	[0.74-0.63]	**
<i>Women</i>	5448									
Moderate / None		0.75	[0.64-0.87]	***	0.77	[0.64-0.90]	***	0.80	[0.68-0.94]	**
Intensive / None		0.67	[0.57-0.79]	***	0.69	[0.58-0.81]	***	0.72	[0.60-0.85]	***
<i>Men</i>	3489									
Moderate / None		0.87	[0.75-1.01]	ns	0.90	[0.78-1.05]	ns	0.97	[0.83-1.13]	ns
Intensive / None		0.87	[0.75-1.01]	ns	0.91	[0.78-1.06]	ns	0.94	[0.80-1.10]	ns

^a Cox proportional model adjusted for sex, centre, education (no or primary/middle/ high school/university), income (<1500, >1500 euros/month), occupation (white collar/employee/blue collar/housewife)

^b Cox proportional model adjusted for sex, centre, education, income, occupation, smoking, alcohol consumption, history of cardiovascular diseases, BMI, depression, diabetes, [hypertension](#), [hypercholesterolemia](#), dependence (IADL), self-rated health, self-rated diet quality, number of drugs, number of chronic diseases.

*P<0.05; **P<0.01; ***P<0.001; ns=not significant

Supplementary Table 2 - Association between 10-year overall mortality (event N=1033) and dietary habits in the subgroup of the elderly cohort of the 3C Study with data on physical activity (only Dijon and Montpellier; N=5273): crude and adjusted hazard ratios (HR) and 95% CI

Dietary habits	N	Crude	(95% CI)	p	Model 1 ^a	(95% CI)	p	Model 2 ^b	(95% CI)	p
At least 1 fruit and 1 vegetable cooked or raw/day	5273	0.83	[0.73-0.94]	**	0.85	[0.74-0.96]	*	0.90	[0.79-1.03]	ns
Cooked fruits or vegetables: ≥ 4-6 servings/week	5273	0.71	[0.59-0.85]	***	0.76	[0.63-0.91]	**	0.75	[0.62-0.90]	**
Meat: ≥1 serving/day	5273	1.10	[0.96-1.27]	ns	1.03	[0.89-1.19]	ns	1.01	[0.87-1.16]	ns
Fish: ≥2 servings/week	5273	0.85	[0.75-0.96]	**	0.88	[0.78-1.00]	*	0.92	[0.81-1.04]	ns
Olive oil use	5273									
Moderate / None§		0.85	[0.72-0.99]	*	0.89	[0.76-1.04]	ns	0.97	[0.83-1.15]	ns
Intensive / None§		0.81	[0.69-0.95]	**	0.85	[0.72-1.00]	*	0.90	[0.76-1.06]	ns

^a Cox proportional model adjusted for sex, centre, education (no or primary/middle/high school/university), monthly income (<1500, >1500 euros), occupation (white collar/employee/blue collar/housewife)

^b Cox proportional model adjusted for sex, centre, education, income, occupation, smoking, alcohol consumption, history of cardiovascular diseases, BMI, depression, diabetes, [hypertension](#), [hypercholesterolemia](#), dependence, self-rated health, self-rated diet quality, number of medication, number of chronic diseases and [physical activity](#)

§reference *P<0.05; **P<0.01; ***P<0.001; ns=not significant

Supplementary Table 3 - Association between 10-year overall mortality (event N=1033) and diet diversity in the subgroup of the elderly cohort of the 3C Study with data on physical activity (only Dijon and Montpellier; N=5273): crude and multivariate adjusted of hazard ratios (HR) and 95%

	N	Crude	(95% CI)	p	Model 1 ^a	(95% CI)	p	Model 2 ^b	(95% CI)	p
Diversity diet score 4-5 / 0-3	5273	0.90	[0.79-1.02]	ns	0.89	[0.78-1.02]	ns	0.92	[0.80-1.05]	ns
Various fats > 3/≤ 3	5273	0.86	[0.74-1.01]	ns	0.88	[0.75-1.02]	ns	0.90	[0.77-1.05]	ns

^a Cox proportional model adjusted for sex, centre, education (no or primary/middle/high school/university), monthly income (<1500, >1500 euros), occupation (white collar/employee/blue collar/housewife)

^b Cox proportional model adjusted for sex, centre, education, income, occupation, smoking, alcohol consumption, history of cardiovascular diseases, BMI, depression, diabetes, hypertension, hypercholesterolemia, dependence, self-rated health, self-rated diet quality, number of medication, number of chronic diseases and physical activity

*P<0.05; **P<0.01; ***P<0.001; ns=not significant