

13 C and 15 N natural isotope abundance reflects breast cancer cell metabolism

Illa Tea, Estelle Martineau, Ingrid Antheaume, Julie C Lalande, Caroline C Mauve, Francoise Gilard, Sophie C Barillé-Nion, Anneke C Blackburn, Guillaume Tcherkez

► To cite this version:

Illa Tea, Estelle Martineau, Ingrid Antheaume, Julie C
 Lalande, Caroline C Mauve, et al.. 13 C and 15 N natural isotope abundance reflects breast cancer cell metabolism. Scientific Reports, 2016, 6, pp.34251.
 10.1038/srep34251 .
 inserm-01385206

HAL Id: inserm-01385206 https://inserm.hal.science/inserm-01385206

Submitted on 21 Oct 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SCIENTIFIC REPORTS

Received: 11 May 2016 Accepted: 08 September 2016 Published: 28 September 2016

OPEN ¹³C and ¹⁵N natural isotope abundance reflects breast cancer cell metabolism

Illa Tea^{1,2}, Estelle Martineau^{1,3}, Ingrid Antheaume¹, Julie Lalande¹, Caroline Mauve⁴, Francoise Gilard⁴, Sophie Barillé-Nion⁵, Anneke C. Blackburn² & Guillaume Tcherkez⁶

Breast cancer is the most common cancer in women worldwide. Despite the information provided by anatomopathological assessment and molecular markers (such as receptor expression ER, PR, HER2), breast cancer therapies and prognostics depend on the metabolic properties of tumor cells. However, metabolomics have not provided a robust and congruent biomarker yet, likely because individual metabolite contents are insufficient to encapsulate all of the alterations in metabolic fluxes. Here, we took advantage of natural ¹³C and ¹⁵N isotope abundance to show there are isotopic differences between healthy and cancer biopsy tissues or between healthy and malignant cultured cell lines. Isotope mass balance further suggests that these differences are mostly related to lipid metabolism, anaplerosis and urea cycle, three pathways known to be impacted in malignant cells. Our results demonstrate that the isotope signature is a good descriptor of metabolism since it integrates modifications in C partitioning and N excretion altogether. Our present study is thus a starting point to possible clinical applications such as patient screening and biopsy characterization in every cancer that is associated with metabolic changes.

Medical applications of stable isotopes are now widespread, like the well-known ¹³C-urea breath assay for ulcer detection¹. This takes advantage of ¹³C-labelling and thus usually neglects differences in reaction rates between isotopic forms, because the isotopic signal used for diagnosing is far above small natural variations in ¹³C. By contrast, the use of isotopes at natural abundance exploits such subtle differences (referred to as isotope effects) to identify bottlenecks in metabolic pathways (rate-limiting steps) or the contribution of multiple elemental sources (mass balance), without the need to introduce expensive isotope tracers into the patient. Isotope effects in metabolism are mostly caused by enzymatic reactions that preferentially consume substrates containing either the light or the heavy isotope (isotopologues) and therefore, the natural isotope abundance in metabolites depends on metabolic fluxes and source substrates². For example, the natural 13 C abundance in respired CO₂ has been used to trace diet composition and substrate changes during exercise^{3,4}. In cancer biology, the use of natural variations in Cu and S stable isotopes in hepatocellular carcinoma has been attempted recently⁵. But to our knowledge, no study has looked at alterations of natural isotope abundance in breast cancer. Due to changes in primary C and N metabolism such as increased glycolysis, glutaminolysis and nucleotide synthesis⁶, important changes in ¹³C and ¹⁵N natural abundance can be anticipated.

To address this question, we examined the isotopic signature of intact breast cancer biopsies (mostly from invasive ductal carcinoma, IDC) and cultured breast cancer cell lines (Supplementary Tables S1 and S2) using elemental analysis coupled to isotope ratio mass spectrometry (EA-IRMS). This technique has been recently shown to be applicable to the biochemical analysis of cancerous cell lines7.

¹EBSI team, CEISAM, University of Nantes-CNRS UMR 6230, 2 rue de la Houssinière BP 92208, F-44322 Nantes, France. ²Cancer Metabolism and Genetics Group, The John Curtin School of Medical Research, The Australian National University, Canberra 2601 ACT, Australia. ³Spectromaitrise, CAPACITÉS SAS, 26 Bd Vincent Gâche, 44200 Nantes, France. ⁴Plateforme Métabolisme-Métabolome, Bâtiment 630 IPS2, Université Paris-Sud, 91405 Orsay cedex, France. ⁵CRCNA, UMR INSERM U892/CNRS 6299/Université de Nantes, Team 8 «Cell survival and tumor escape in breast cancers», Institut de Recherche en Santé de l'Université de Nantes, 8 quai Moncousu, BP 70721, 44 007 Nantes cedex 1, France. ⁶Research School of Biology, The Australian National University, Canberra 2601 ACT, Australia. Correspondence and requests for materials should be addressed to I.T. (email: illa.tea@anu.edu.au)



Figure 1. Breast cancer biopsies are naturally ¹³C-enriched and nitrogen-rich. (a) The natural abundance in ¹³C (δ^{13} C, vs. V-PDB) and ¹⁵N (δ^{15} N, vs. air N₂) differentiates control from tumor patient tissue samples (n = 23). One sample pair of healthy (adjacent) and cancerous tissues not characterized for receptors (u and u', respectively) and three triple negative tumor samples (t) are also shown. (b) Taken as a whole, the change in the ¹³C-abundance ($\Delta\delta$) in cancerous tissues in tumor/adjacent tissue pairs was significant (*P* < 0.01, n = 5). (c) ¹³C-abundance ($\Delta\delta$) was inversely correlated with the total lipid content (r² = 0.48, *P* < 0.03). (d) Regardless of receptor expression, cancerous tissues had a higher elemental content in N (*P* < 0.05). In (a,c), each datum is the average of 3 sub-samples. In (d), lower case letters stand for statistical classes, *P* < 0.05.

.....

Results

Isotope signature of breast cancer biopsies. Natural ¹³C and ¹⁵N abundance was found to discriminate normal and cancerous biopsies, the latter being significantly ¹³C-enriched by ~4‰ and tended to be (though not significantly) ¹⁵N-depleted (Fig. 1a,b). Triple negative tissues (ER- PR- HER2-) tended to be relatively ¹⁵N-depleted (Fig. 1a and Supplementary Table S1). The observed isotopic difference between cancerous and non-cancerous tissues can have been caused by either variation in chemical composition or isotope fractionations in metabolism.

First, there were alterations in total lipid content, which correlated to the ¹³C-abundance (Fig. 1c and Supplementary Figure S1). In fact, lipids are typically ¹³C-depleted components of human tissues⁸ and a systematic ¹³C-depletion has been observed in adipose tissue⁹. Here, cancerous tissues are on average less lipid-rich, reflecting the lower abundance of adipose cells in breast IDC^{10,11}. Cancerous tissues were also enriched in nitrogenous compounds, as shown by the higher N elemental content (Fig. 1d). This effect is due to either a lower cellular N turnover rate or a higher content in protein and total amino acids including arginine (Supplementary Figure S2). Second, metabolic fluxes were modified thereby modifying the net exchange with the extracellular fluid and thus the overall (apparent) cellular isotope fractionation (see below). It is highly unlikely that the isotopic difference between biopsies was due to a confounding effect of nutritional preferences amongst patients (such as the proportion of animal proteins or sugars from C₄-plants)¹²⁻¹⁴ because the comparison of paired normal and cancerous biopsy samples from the same patient still shows a significant difference for ¹³C (Fig. 1b). ¹³C and ¹⁵N abundance were also corrected so as to express them relative to a common elemental source (see Methods below). Moreover, female patients from which biopsies originated had the same geographical origin (western France) and thus important differences between nutritional habits or general isotope composition of body matter are improbable¹⁵.



Figure 2. Cancerous cell lines are naturally ¹³C-enriched and ¹⁵N-depleted. When compared to source C and N used for growth, the natural isotopic enrichment (ε ¹³C and ε ¹⁵N) differentiates the control immortalized line (MCF10A) and the breast tumor ascites-derived line (ZR75-1) from other adenocarcinoma lines (**a**). The isotope enrichment in lipids parallels that in total organic matter, and cancerous lines contain more (¹³C-enriched) C₄-acids (**b**). Despite some variations, transcript quantitation of key enzymes shows a generally lower expression in cancer cells compared to MCF10A cells of enzymes of the urea cycle, lipid synthesis or the anaplerotic pathway (gene abbreviations in Supplementary Table S5), suggesting that metabolite excretion in addition to biosynthesis is involved in the isotopic differences (**c**).

Metabolic alterations in breast cancer biopsies. Targeted metabolomics analyses by GC-MS were then carried out to identify major metabolic alterations. Normal tissues were clearly differentiated from cancerous tissues in both unsupervised (PCA) and supervised (orthogonal partial least square-discriminant analysis, OPLS-DA) analysis (Supplementary Figures S3 and S4), with ethanolamine, methionine and arginine having the highest weight in such a differentiation (Supplementary Figures S3 and S4). Other discriminating metabolites included carbohydrates (such as fructose) and putrescine, and these metabolites have also been found to be significant elsewhere^{16,17}. While the effect on free ethanolamine reflected the general reorchestration of lipid metabolism (urea cycle), thereby causing changes in N excretion and thus, by mass balance, in ¹⁵N-abundance and N elemental content.

Isotopic signature of cell lines. Potentially, the isotopic signature in biopsies may have been influenced by non-controlled epidemiological and clinical factors (e.g., patient age, pharmacological treatments, etc.). Therefore, we also examined different cell lines cultured under isotopically-controlled conditions: one immortalized, non-cancerous cell line (MCF10A) and six breast cancer cell lines (ZR75-1, MCF7, SKBR3, MDA-MB-231, MDA-MB-468 and Cal51, see Supplementary Table S2). The use of cultured cells further allowed us to carry out isotopic compound-specific analyses as well as targeted gene-expression, metabolomics and EA-IRMS analyses (see Methods) on the same samples.

With the exception of ZR75-1 (non-pleural origin, ER+), all other breast cancer cell lines were ¹⁵N-depleted (Fig. 2a). The ¹⁵N-abundance in DNA, proteins and total soluble metabolites paralleled that in total organic matter (Supplementary Figure S5). The ¹³C-abundance in non-cancerous cells was found to be lower than in cancerous cells (Fig. 2a). It did not correlate to the lipid content but rather to the ¹³C-abundance in lipids and loosely, to the content in C₄ organic acids (Fig. 2b) which are known to be ¹³C-enriched¹⁹.

Primary metabolism in cell lines. The endometabolome (i.e., intracellular metabolome) was analyzed to gain information on primary C and N metabolism (summarized in Supplementary Figure S6a). Some metabolites such as arginine, urea and glucose were significantly different between cancer and normal cell lines. Close correlations or anti-correlations were obtained between ¹³C- or ¹⁵N-abundance and key metabolites or metabolic ratios such as urea/arginine and fumarate/aspartate (Fig. 3a). As such, cancer cell lines could be differentiated from the MCF10A non-cancerous cells using these ratios (Fig. 3a).

The present results on both metabolome and isotopes suggest that three metabolic pathways are at the origin of the isotopic differences between cell lines: the urea cycle (arginine metabolism), the anaplerotic carbon fixation, and lipid synthesis. Transcript levels of a selection of genes encoding key enzymes of these pathways was examined by qPCR and in most cases, these genes were less expressed in cancer cell lines (Fig. 2c). Thus, rather than substantial changes in enzyme abundance, the alteration of isotope abundance was caused by changes in metabolic fluxes and/or net exchange of metabolites (absorption, excretion and accumulation) with the extracellular medium.

Isotopic efflux in cell lines. We then examined the metabolic composition of the extracellular medium (exometabolome, Supplementary Figure S6b) to determine the preferential distribution (extracellular vs. intracellular) of metabolites and compute isotopic fluxes (isofluxes). The extracellular-to-intracellular difference and





the ¹⁵N-abundance were found to be best anti-correlated with arginine, β -alanine, ethanolamine and spermidine (Supplementary Figure S6c,d). Amongst these, arginine had the highest value of the extracellular-to-intracellular difference (Fig. 3b,c) and was the most significantly different between cancer and normal cell lines (*P* < 0.005) and thus, appeared to be the best candidate to explain the ¹⁵N-depletion (Fig. 3d). There was little difference in the ¹⁵N-abundance of arginine itself amongst cell lines, although the non-cancerous (MCF10A) and ER lines (ZR75-1 and MCF7) synthesized ¹⁵N-enriched arginine (by a few ‰). Still, in all cases, arginine was much more ¹⁵N-depleted than the isotopic influx (isotope composition of substrates consumed by cells from supplied culture medium) or extracellular urea (Fig. 3d). As a result, arginine efflux from cells represented a very substantial part of the isotopic efflux to the extracellular medium in both ZR75-1 and MCF10A lines but was modest in others (Fig. 3e). That is, cancerous lines were naturally ¹⁵N-depleted (Fig. 3b) because of the proportionally lower excretion of ¹⁵N-depleted arginine. The higher metabolic ratio urea-to-arginine then comes as no surprise: it reflected the prevalence of arginine metabolism via the urea cycle (Fig. 3a).

Discussion

We showed isotopic differences between healthy and cancer tissues or cells, and provided evidence that these differences are related to urea, anaplerotic C fixation and/or lipid metabolism, and that the balance in nitrogen metabolism is modified in breast adenocarcinoma tissues or cells compared to normal breast tissues or cells thereby altering the natural ¹⁵N-abundance (Fig. 4). The consumption of glutamine (glutaminolysis) is a major pathway in cancerous cells²⁰ that involves deamidation by glutaminase, thereby producing ammonia which is partly recycled and excreted in the form of urea. Here, source glutamine is naturally 15 N-depleted (δ^{15} N = -6.5‰) and 14 N/ 15 N isotope effects in both deamidation and the urea cycle²¹⁻²³ yield 15 N-depleted arginine. The efflux (excretion) of arginine does not counterbalance the N influx and this imbalance is compensated for by intracellular build-up and recycling into polyamines and β -alanine (Fig. 4) thereby causing a general ¹⁵N-depletion of organic matter. In addition to acid metabolism (C_4 -acids and lactate, Fig. 2b, Supplementary Table S3), the urea cycle incorporates bicarbonate (naturally ¹³C-enriched)²⁴ via the action of carbamoyl phosphate synthase and thus, arginine build-up contributes to the enrichment of ¹³C in cells. This appears indeed to be the best explanation considering known isotope effects in metabolism^{6,21,25-29} (Supplementary Table S4). In fact, the natural ¹³C-enrichment cannot come from a change in total lipid content (as in tumors) since lipid content variation was small (Supplementary Figure S7) and furthermore, the isotope composition in lipids followed that in total organic matter (Fig. 2b).

Quite noticeably, the isotopic pattern seen in tumor biopsies was similar to that found in cell lines, that is, with a tendency to be ¹⁵N-depleted and ¹³C-enriched (Fig. 1a). This pattern is unlikely to come from an effect of nutrition, but rather explained by the specific primary C and N metabolism in tumor tissues, as explained above (Fig. 1). The metabolic origin of the ¹³C-enrichment was different to that in cultured cells (a decrease in total lipid content in tumors, Supplementary Figure S7 and refs 18,30). This is not surprising since cultured cell metabolism is different from that of cells *in situ*. By contrast, the mechanism explaining the lower ¹⁵N-abundance was similar: a net influx of ¹⁵N-depleted nitrogen associated with any metabolism. In cancerous cultured cells, there is an increased activity of the urea cycle and non-quantitative excretion of ¹⁵N-depleted arginine. In tumors *in situ*, it is believed



Figure 4. Major metabolic pathways are responsible for changes in the natural ¹³C and ¹⁵N abundance in cancerous cultured cells. Glutamine (Gln) is a major N and C source from which N is removed via hydrolysis (glutaminase **①**) and the urea cycle (**②**). These reactions are both fractionating against ¹⁵N thereby yielding ¹⁵N-depleted urea and arginine (Arg). Therefore, build-up and recycling rather than excretion of Arg (dotted arrow) tend to deplete cancer cells of ¹⁵N. Cells are also ¹³C-enriched due to the fixation of bicarbonate by carbamoyl-phosphate (CP) synthesis (to feed the urea cycle **③**) and the anaplerotic pathway (**④**), as well as a lower ¹³C content in non-structural lipids (dotted arrow). Lipids that are ¹³C-depleted come from the natural ¹³C-depletion in acetyl-CoA (Ac-CoA) (inherited from naturally ¹³C-depleted C-atom positions in glucose) and the isotope effect of pyruvate dehydrogenase^{45,46} (**⑤**). Amine acceptors are denoted as 'A' and oxaloacetate (OAA) converted to aspartate (Asp) is provided as an example. 2OG, 2-oxoglutarate; Pyr, pyruvate; Lact, lactate. The potential contribution of respiration (CO₂ loss) and lactate excretion to the natural ¹³C-abundance is described in Supplementary Table S4.

that cells are arginine-dependent to sustain polyamine metabolism³¹, and the auxotrophy of tumors for arginine has been exploited in the development of anticancer therapies³². Therefore cells capture naturally ¹⁵N-depleted arginine from their microenvironment, thereby leading to a ¹⁵N-depletion of cellular organic matter. Such a mechanism agrees with the well-known increase in polyamine metabolism³³ and the fact that urea and glutamine (glutaminase activity) are amongst the best urinary and tumor biomarkers for breast cancer, respectively^{34,35}. Also, differential expression of arginase, a key enzyme of the urea cycle, has been reported in breast cancer cell lines^{36,37}.

The natural abundance in ¹³C and ¹⁵N thus reflects primary metabolism of cancer cells and thus, offers potential opportunities for development of biomarkers for clinical research. Consistently, healthy and cancerous samples were found to fall in corresponding regions of the isotopic landscape (Fig. 1a), and normal and cancerous cell lines could be differentiated (Fig. 2a). The ability of natural ¹³C- or ¹⁵N-abundance to further differentiate tumor grade and severity remains to be explored. In addition, there is a potential use the natural isotope abundance to distinguish between different breast cancer subtypes, as they have been shown to have different metabolic phenotypes³⁸. Also, isotopic measurements would be best adapted to clinical settings if measured in plasma metabolites rather than in biopsies and thus, future work on both the accuracy and applicability of such plasma isotopic markers is required. For example, ¹⁵N natural abundance in plasma arginine could be exploited as a marker provided the metabolic activity of the tumor has a detectable effect on the total circulating arginine pool. That isotopic signature of tumorous breast tissues is due to alterations in major pathways of primary metabolism highlights the potential relevance of our findings to other cancer types which also display changes in metabolic fluxes.

Methods

Human Tissues. Fresh human mammary samples were obtained from chemotherapy naive patients with invasive carcinoma after surgical resection at the Institut de Cancérologie de l'Ouest – René Gauducheau, Nantes, France. Informed consent was obtained from patients to use their surgical specimens and clinicopathological data for research purposes, as required by the French Committee for the Protection of Human Subjects (CCPPRB). Ouest IV – Nantes CCPPRB approved use of tumor tissues for this study (6 May 2013: no. 357/2013). The collection of tumors was approved by the French Minister of higher education and research (no. AC-2008-141). This study did not require additional ethical approval. Amongst 17 breast cancer patients, 12 breast cancer samples and 5 pairs of breast cancer and adjacent non-cancerous tissues were obtained. One normal breast sample was obtained from reduction mammoplasty. For tumors, hormonal receptors, histological type and grade information were available, except that a paired of tissue #22/#19 (Supplementary Table S1, u and u' in Fig. 1a) did not have receptor status available.

Cultured cells. Six human breast cancer cell lines and one non-cancerous cell line (Supplementary Table S2) were used and grown as monolayer cultures (10^7 cells/75 m² flask). The cancer cells were grown in no-glucose

no-glutamine DMEM supplemented with 10% FBS and 1% 10,000 units/ml penicillin, 10,000 units/ml streptomycin at 37 °C in a humidified atmosphere of 5% CO_2 . The non-cancerous cell line (MCF10A) was grown in DMEM/F-12 (no-glucose no-glutamine) supplemented with 5% horse serum, 0.01% EGF (Epidermal Growth Factor), 0.005% cholera toxin, 0.1% insulin, 0.005% hydrocortisone, and 1% HEPES. L-glutamine and D-glucose (4 and 25 mM, respectively) were added separately to culture mediums to ensure these nutrients were isotopically identical for all cells. The same cells were grown several times to test the biological variability. As soon as cells were confluent (~10⁷ cells), cell counting and extraction were done using triplicate cell culture flasks. One flask was used for cell counting, after harvesting with trypsin, using trypan blue staining and microscopic hemocytometry. For other analyses carried out on the other flasks, cells were then prepared as an aqueous (isotopes) or methanolic (metabolomics) extract without using trypsin as explained below.

Extraction and separation of studied fractions. *Human tissues.* Human fresh tissues were lyophilised. Lipids were extracted from the dried human tissues (10 mg) with hexane (1 mL). After centrifugation (4000 rpm, 15 min, 4 °C), the pellet was kept for metabolomics and the lipids were recovered by evaporation of hexane under a stream of N₂ gas at 40 °C.

Cultured cells. After culture, cells were washed twice with PBS, and 3 mL of cold water was added to the flask. Cells were detached from the flask using a cell scraper. The lysed cell preparation was kept at -20 °C for 48 h and then lyophilized. Different fractions (lipids, total soluble fraction (TSF), protein and DNA) were extracted and separated from the dry cell material. To obtain a lipid fraction, lyophilized cells (40 mg) were ground and extracted twice under agitation with 2 mL cyclohexane for 5 min at 20 °C. After centrifugation (4000 rpm, 5 min, 20 °C), the lipid fraction (4 mL of cyclohexane) was dried under a stream of N₂ gas at 40 °C and the dry residue stored at -20 °C until required. To obtain a total soluble fraction, the pellet (insoluble in cyclohexane) obtained after centrifugation (4000 rpm, 5 min, 20 °C), the total soluble fraction was dried under a stream of N₂ gas at 40 °C and the dry cell after centrifugation (4000 rpm, 5 min, 20 °C), the total soluble fraction was dried under a stream of N₂ gas at 40 °C and the dry cell after centrifugation (4000 rpm, 5 min, 20 °C), the total soluble fraction was dried under a stream of N₂ gas at 40 °C and the dry residue stored at -20 °C until required. The pellet (insoluble in methanol/water) consisted of a protein fraction.

For DNA purification, lyophilised cells (15 mg) were reconstituted with 1 mL water and extracted with an equal volume of phenol/chloroform in an ultrasound bath for 30 min. After centrifugation (4000 rpm, 5 min, 4° C), the aqueous phase was filtered to remove proteins and the DNA was precipitated with an equal volume of isopropanol (1 mL). After gentle agitation, the DNA fragments were pelleted by centrifugation (4000 rpm, 5 min, 4° C), washed twice with ethanol (70%) to remove impurities, and dried under a stream of N₂ gas at 40 °C.

¹³C and ¹⁵N abundance by elemental-analysis coupled to isotope ratio mass spectrometry (EA-IRMS). Subcellular fractions (see above) were lyophilized for EA-IRMS analysis, in order to detect their ¹³C and ¹⁵N isotopes. From each sample ~0.7 mg was weighed with 10^{-5} g precision balance (Ohaus Discovery DV215CD, Pine Brook, New Jersey, USA) into each of three tin capsules (solids "light" 5×9 mm, Thermo Fisher Scientific, www.thermo.com) for isotope analysis. EA-IRMS analyses were performed as previously described⁷. The ¹³C and ¹⁵N abundance was first expressed as δ values (δ^{13} C and δ^{15} N, in ‰) as $\delta = R/R_{st} - 1$ where *R* is the heavy-to-light isotope ratio and R_{st} stands for the isotope ratio in the international reference (V-PDB for ¹³C and air N₂ for ¹⁵N).

The isotope abundance in sample fractions was expressed relative to that in total organic matter using the enrichment (in ‰) $\epsilon = \delta_f - \delta_{tom}$ where δ_f is the δ value in the fraction of interest (lipids, proteins, etc.) and δ_{tom} the (non-corrected) δ value in total organic matter. The enrichment ϵ is then positive if the sample fraction is enriched in the heavy isotope and negative if it is depleted. In this way, ϵ nearly equals the opposite of the isotope fractionation Δ (which is given by $[\delta_{tom} - \delta_f]/[\delta_f + 1] \approx \delta_{tom} - \delta_f$). For total organic matter in Figs 1 and 2 and Supplementary Figure S3, δ values were corrected to allow comparisons between patients (that may have had different nutritional preferences) and between cell lines (that had slightly different culture media), as follows:

In cultured cells, it was corrected for the δ value of the elemental C and N source using the classical relationships accounting for isotope fractionation: $\delta_{corr} = (\delta_{av} - \Delta)/(\Delta + 1)$ where $\Delta = (\delta_{srce} - \delta)/(\delta + 1)$, δ_{srce} is the δ value of the elemental source and δ_{av} is δ value used as the reference (here, the average of δ_{srce} values was used as the reference value; note that any change in this value would not change the pattern seen here, it would only change the absolute scale of δ_{corr} values). δ_{srce} was the $\delta^{15}N$ and $\delta^{13}C$ of source amino acids and sugars, respectively, in intact culture medium. When expressed as a fractionation (Fig. 2a), the isotope composition in total organic matter of cells is given as $\epsilon = \delta - \delta_{srce}$. That way, ϵ values are positive when it is enriched in the heavy isotope and negative when it is depleted, as for ϵ in individual cell fractions (see above).

For biopsies, δ_{srce} could not, quite obviously, be known because food composition of patients was not controlled. We therefore had to use a correction that could be calculated independently of food composition. The major influence of nutrition on elemental composition and isotopes is the protein/fat ratio of ingested food³⁹, with fat-rich and protein-poor food causing a general increase in %C (and decrease in %N) and a general decrease in both δ^{15} N and δ^{13} C in body tissues. Therefore, we renormalized the δ values in total organic matter by taking into account the %C and %N using $\delta_{\text{corr}} = \delta + (1 - f)\tau$ where *f* is the elemental composition relative to the average and τ is the apparent isotope fractionation between the whole population δ value (average isotope composition of human tissues in European western (French) population (δ_a), -22% and 9% for ¹³C and ¹⁵N, respectively) and the average of our samples (δ_m): $\tau = \delta_a - \delta_m$. For example, in this way, protein-rich samples (lean patients) that are naturally impoverished in elemental C and richer in ¹³C will have a negative correction for δ^{13} C. Of course, such a correction was not ideal because the knowledge of the average isotope composition of ingested food would have been better. In practice, this correction was minimal (on average, 0.02‰ for ¹³C and 0.5‰ for ¹⁵N) and did not cause a substantial bias in our data.

¹⁵N-abundance and amino acid content using gas chromatography - combustion - isotope ratio mass spectrometry (GC-C-IRMS). When cells were confluent, the culture medium was kept (10 mL from ~10⁸ cells), and cells (~10⁸) were washed with PBS and then lysed in 36 mL of 80% methanol. After homogenization, methanolic extracts were dried under a stream of N₂ gas at 40 °C. Amino acids were prepared as their *N*-pivaloyl, *O*-isopropyl esters for GC-C-IRMS analysis, and used for both quantification (µmol) and ¹⁵N values (‰) as previously described⁴⁰.

¹³C-abundance in metabolites using liquid chromatography coupled to chemical oxidation and isotope ratio mass spectrometry (LC-co-IRMS). For LC-co-IRMS analysis, ~10⁷ cells were used in 12 mL of 80% methanol. LC-co-IRMS analysis was performed as previously described, using water as a mobile phase and a RS Pak KC-811 column (Shodex)⁴¹. δ^{13} C values were corrected using pure standard solutions at different concentrations. In practice, such corrections due to non-linearity were very small, about 0.2‰.

GC-MS metabolomics. *Human biopsies.* If material remained after lyophilization and lipid extraction, an additional ~10–15 mg of tissue was ground and dissolved in a standard solution of DL-2-Aminobutyric acid (0.2 mM, 950 μ L) and methanol/water (950 μ L, 80:20 v/v) was added to extract the soluble metabolites. After centrifugation (14000 rpm, 15 min, 4 °C), the soluble fraction was mixed with standard solution of ribitol (2 mM, 95 μ L). The solution was centrifuged again and three aliquots of 200 μ L were dried using a speed-vac before derivatization and GC-MS analysis as described previously^{41,42}.

Cultured cells and extracellular medium. About 5×10^7 cells were dissolved in a solution of methanol/water (7 mL, 60:40 v/v) containing 500 µL of a standard solution D,L-2-aminobutyric acid (0.2 mM). After centrifugation (14000 rpm, 15 min, 4 °C), the soluble fraction was mixed with another standard solution of ribitol (2 mM, 50 µL). The solution was centrifuged again and three aliquots of 1.4 mL, i.e. 20% of ~ 5×10^7 cells, were dried using a speed-vac before derivatization and GC-MS analysis. For extracellular medium, the same standard solutions [D,L-2-aminobutyric acid (0.2 mM, 500 µL) and ribitol (2 mM, 50 µL) and methanol/water solution (500 µL, 60:40 v/v)] were added to 500 µL of media collected from the cultured cells. After a centrifugation, three aliquots of 300 µL were prepared for GC-MS analysis. All metabolomics data were normalized to the specific number of cells so that the exo- and endo-metabolome could be directly compared.

RT-qPCR. RNA was isolated from cell lines and 500 ng RNA was reverse transcribed as previously described⁴³. Quantitative PCR was done using the Maxima SYBR Green/ROX qPCR Master Mix (Life Technologies) and the MX4000 instrument (Stratagene, Basel, Switzerland), according to the manufacturer's instructions. The endogenous housekeeping genes *RPLPO*, *HPRT1* and *ACTB1* were used for normalization. Relative quantification was carried out by using the $\Delta\Delta$ Ct method. Primer sequences are listed in Supplementary Table S5.

Statistics. Pair-wise comparisons were carried out using Student-Welsh tests, with a p-threshold value of 0.01 unless otherwise stated in text. Metabolomics were examined using unsupervised analysis with principal component analysis (PCA) or supervised analysis with orthogonal partial least square discriminant analysis (OPLS-DA), both carried out with Simca (Umetrics). Heat maps and hierarchical clustering (HCL) were done using the open source software MeV⁴⁴. The HCL was carried out using the cosine correlation as a measure of proximity.

References

- 1. Graham, D. Y. et al. Campylobacter pylori detected noninvasively by the ¹³C-urea breath test. The Lancet 1, 1174–1177 (1987).
- Schmidt, H.-L., Robins, R. J. & Werner, R. A. Multi-factorial *in vivo* stable isotope fractionation: causes, correlations, consequences and applications. *Isotopes Environ Health Stud* 51, 155–199 (2015).
- Bartstow, T. J., Cooper, D. M., Epstein, S. & Wasserman, K. Changes in ¹³CO₂/¹²CO₂ consequent to exercise and hypoxia. J Appl Physiol 66, 936–942 (1989).
- Perkins, S. E. & Speakman, J. R. Measuring natural abundance of ¹³C in respired CO₂: variability and implications for non-invasive dietary analysis. *Funct Ecol* 15, 791–797 (2001).
- Balter, V. et al. Natural variations of copper and sulfur stable isotopes in blood of hepatocellular carcinoma patients. Proc Natl Acad Sci USA 4, 982–985 (2015).
- 6. De Berardinis, R. J., Sayed, N., Ditworth, D. & Thompson, C. B. Brick by brick: metabolism and tumor cell growth. Curr Op Gen Dev 18, 54–61 (2008).
- Tea, I., Martineau, E., Giraudeau, P., Akoka, S. & Barillé-Nion, S. Method for characterising the origin and/or condition of diseased or healthy cells and uses thereof in biology. Patent WO 2012123886-A1/US2014-0051116 (2012).
- 8. Colonese, A. C. et al. Archaeological bone lipids as palaeodietary markers. Rapid Commun Mass Spectrom 29, 611-618 (2015).
- Arneson, L. S. & MacAvoy, S. E. Carbon, nitrogen, and sulphur diet-tissue discrimination in mouse tissues. *Revue Can J Zool* 83, 989–995 (2005).
- Franck, C. J., McCreery, R. L. & Redd, D. C. B. Raman spectroscopy of normal and diseased human breast tissues. Anal Chem 67, 777–783 (1995).
- 11. Lazebnik, M. et al. A large-scale study of the ultrawideband microwave dielectric properties of normal, benign and malignant breast tissues obtained from cancer surgeries. Phys Med Biol 52, 6093–6115 (2007).
- 12. Petzke, K. J., Boeing, H., Klaus, Š. & Metges, C. C. Carbon and Nitrogen Stable Isotopic Composition of Hair Protein and Amino Acids Can Be Used as Biomarkers for Animal-Derived Dietary Protein Intake in Humans. *J Nutr* **135**, 1515–1520 (2005).
- 13. DeNiro, M. J. & Epstein, S. Influence of diet on the distribution of nitrogen isotopes in animals. *Geochim Cosmochim Acta* 45, 341–351 (1981).
- 14. DeNiro, M. J. & Epstein, S. Influence of diet on the distribution of carbon isotopes in animals. *Geochim Cosmochim Acta* 42, 495–506 (1978).
- O'Connell, T. C., Kneale, C. J., Tasevska, N. & Kuhnle, G. G. C. The Diet-Body Offset in Human Nitrogen Isotopic Values: A Controlled Dietary Study. Am J Phys Anthropol 149, 426–434 (2012).
- 16. Tang, X. et al. A joint analysis of metabolomics and genetics of breast cancer. Breast Cancer Res 16, 415 (2014).
- Budczies, J. et al. Remodeling of central metabolism in invasive breast cancer compared to normal breast tissue a GC-TOFMS based metabolomics study. BMC Genomics 13, 334 (2012).

- 18. Todor, I. N., Lukyanova, N. Y. & Chekhun, V. F. The lipid content of cisplatin- and doxorubicin-resistant MCF-7 human breast cancer cells. Exp Oncology 34, 97-100 (2012).
- 19. Schmidt, H. L. Fundamentals and systematics of the non-statistical distribution of isotopes in natural compounds. Naturwissenschaften 90, 537-552 (2003).
- 20. Wise, D. R. & Thompson, C. B. Glutamine addiction: a new therapeutic target in cancer. Trends Biochem Sci 35, 427-433 (2010).
- 21. Rishavy, M. A., Cleland, W. W. & Lusty, C. J. ¹⁵N isotope effects in glutamine hydrolysis by carbamyl phosphate synthetase: evidence for a tetrahedral intermediate in the mechanism. Biochemistry 39, 7309-7315 (2000).
- 22. Kim, S. C. & Raushel, F. M. Isotopic probes of the argininosuccinate lyase reaction. Biochemistry 25, 4744-4749 (1986).
- 23. Schmidt, H. L. & Medina, R. Possibilities and scope of the double isotope effect method in the elucidation of the mechanisms of enzyme catalyzed reactions. Isotopes Environ Health Studies 27, 1-4 (1991).
- 24. Marlier, J. F. & O'Leary, M. H. Carbon kinetic isotope effects on the hydration of carbon dioxide and the dehydration of bicarbonate ion. J Am Chem Soc 106, 5054-5057 (1984).
- 25. Tcherkez, G., Mahé, A. & Hodges, M. ¹²C/¹³C fractionations in plant primary metabolism. *Trends Plant Sci* 16, 499-506 (2011).
- 26. Marlier, J. F. & O'Leary, M. H. Carbon kinetic isotope effects on the hydration of carbon dioxide and the dehydration of bicarbonate ion. J Am Chem Soc 106, 5054-5057 (1984).
- 27. Lyon, R. C., Cohen, J. S., Faustino, P. J., Megnin, F. & Myers, C. E. Glucose metabolism in drug-sensitive and drug-resistant human breast cancer cells monitored by magnetic resonance spectroscopy. *Cancer Res* **48**, 870–877 (1988). Perkins, S. E. & Speakman, J. R. Measuring natural abundance of ¹³C in respired CO₂: variability and implications for non-invasive
- 28. dietary analysis. Funct Ecol 15, 791-797 (2001).
- 29. Hatch, K. A., Pinshow, B. & Speakman, J. R. The analysis of the 13C/12C ratio in exhaled CO₂: its advantages and potential application to field research to infer diet, changes in diet over time, and substrate metabolism in birds. Int Comp Biol 42, 21-33 (2002).
- 30. Cerussi, A. et al. In vivo absorption, scattering, and physiologic properties of 58 malignant breast tumors determined by broadband diffuse optical spectroscopy. J Biomed Optics 11, 044005 (2006).
- 31. Pavlova, N. N. & Thompson, C. B. The emerging hallmarks of cancer metabolism. Cell Metabolism 23, 27–47 (2016).
- 32. Delage, B. et al. Arginine deprivation and argininosuccinate synthetase expression in the treatment of cancer. Int. J. Cancer 126, 2762-2772 (2010).
- 33. Gerner, E. W. & Mevskens, F. L. Polvamines and cancer: old molecules, new understanding, Nat Rev 4, 781-792 (2004).
- 34. Slupsky, C. M. et al. Urine metabolite analysis offers potential early diagnosis of ovarian and breast cancer. Clin Cancer Res 16, 5835-5841 (2010).
- 35. Budczies, J. et al. Glutamate enrichment as new diagnostic opportunity in breast cancer. Int J Cancer 136, 1619–1628 (2015).
- 36. Singh, R., Pervin, S., Karimi, A., Cederbaum, S. & Chaudhuri, G. Arginase activity in human breast cancer cell lines: Nω-hydroxy-L-arginine selectively inhibits cell proliferation and induces apoptosis in MDA-MB-468 cells. Cancer Res 60, 3305–3312 (2000).
- 37. Straus, B., Cepelak, I. & Festa, G. Arginase, a new marker of mammary carcinoma. Clin Chim Acta 210, 5-12 (1992).
- 38. Llovd, S. M., Arnold, J. & Sreekumar, A. Metabolomic profiling of hormone-dependant cancers: a bird's eye view. Trends in Endocrinology and Metabolism 26, 477-485 (2015).
- Voigt, C. C., Rex, K., Michener, R. H. & Speakman, J. R. Nutrient routing in omnivorous animals tracked by stable carbon isotopes in tissue and exhaled breath. Oecologia. 157, 31-40 (2008).
- Tea, I. et al. Simultaneous determination of natural-abundance δ^{15} N values and quantity of individual amino acids in proteins from milk of lactating women and from infant hair using an isotope ratio mass spectrometry. Rapid Commun Mass Spectrom 27, 1345-1353 (2013)
- 41. Dellero, Y., Lamothe-Sibold, M., Jossier, M. & Hodges, M. Arabidopsis thaliana ggt1 photorespiratory mutants maintain leaf carbon:nitrogen balance by reducing RuBisCO content and plant growth. Plant J 83, 1005-1018 (2015).
- 42. Bathellier, C. et al. On the resilience of nitrogen assimilation by intact roots under starvation, as revealed by isotopic and metabolomic techniques. Rapid Commun Mass Spectrom 23, 2847-2856 (2009).
- 43. Bah, N. et al. Bcl-xL controls a switch between cell death modes during mitotic arrest. Cell Death Dis 5, e1291 (2014).
- 44. Saeed, A. I. et al. TM4: a free, open-source system for microarray data management and analysis. Biotechniques 34, 374–378 (2003).
- 45. Melzer, E. & Schmidt, H. L. Carbon isotope effects on the pyruvate dehydrogenase reaction and their importance for relative carbon-13 depletion in lipids. J Biol Chem 262, 8159-8164 (1987).
- Gilbert, A., Silvestre, V., Robins, R. J. & Remaud, G. S. Accurate quantitative isotopic ¹³C NMR spectroscopy for the determination of the intramolecular distribution of ¹³C in glucose at natural abundance. Anal Chem 81, 8978-8985 (2009)

Acknowledgements

The authors thanks Dr. P. Giraudeau, Pr. S. Akoka, and Dr. J.M. Chao de la Barca and Pr. P. Reynier for preliminary tests for metabolome analysis by NMR and discussion about the manuscript, respectively. The authors are grateful to M. Grand for ¹³C lipid analysis by EA-IRMS analyses. The authors thank the support of the Agence Nationale de la Recherche through a research grant Jeunes Chercheurs (under contract 12-000101), the metabolomics platform CORSAIRE from the Biogenouest network, the funding from SATT (Nantes University Technology Transfer Services), from PLAISIR Pays de la Loire Association for International Structure on "Isotopomic Research" French Regional International Strategy Grant and the Australian Research Council through a Future Fellowship (under contract FT14 0100645). The authors thank the company Elementar France for kindly providing equipment to carry out LC-co-IRMS analyses.

Author Contributions

I.T. and G.T. designed and supervised the project. E.M., I.A. and C.M. collected and generated the isotopic data and performed isotopic analyses; J.L. performed qPCR analyses; Metabolomics data were performed by F.G. and S.B.-N. provided tissue and cell samples. S.B.-N. and A.C.B. advised on breast cancer metabolism context. G.T. and I.T. performed statistical analyses, prepared the figures and tables and wrote the manuscript; S.B.-N. and A.C.B. revised the manuscript; all authors viewed, had the opportunity to comment on and approved the final draft.

Additional Information

Supplementary information accompanies this paper at http://www.nature.com/srep

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Tea, I. et al. ¹³C and ¹⁵N natural isotope abundance reflects breast cancer cell metabolism. Sci. Rep. 6, 34251; doi: 10.1038/srep34251 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/

© The Author(s) 2016