



Non-Coding RNAs in Cutaneous Melanoma Development, Progression and Dissemination

Virginie Vignard, Delphine Fradin

► To cite this version:

Virginie Vignard, Delphine Fradin. Non-Coding RNAs in Cutaneous Melanoma Development, Progression and Dissemination. OBM Genetics, 2018, 2 (2), pp.1 - 1. 10.21926/obm.genet.1802020 . inserm-01804848

HAL Id: inserm-01804848

<https://inserm.hal.science/inserm-01804848>

Submitted on 1 Jun 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Review Article

Non-Coding RNAs in Cutaneous Melanoma Development, Progression and Dissemination

Virginie Vignard ^{1,2}, Delphine Fradin ^{1,*}

1. CRCINA, Inserm, Université de Nantes, IRSUN 8 quai Moncousu, Nantes, France; E-Mail: delphine.fradin@inserm.fr

2. CHU Nantes, Nantes, France; E-Mail: virginie.vignard@univ-nantes.fr

* **Correspondence:** Delphine Fradin; E-Mail: delphine.fradin@inserm.fr

Academic Editors: Stéphane Viville and Marcel Mannens

Special Issue: [Epigenetic Mechanisms in Health and Disease](#)

OBM Genetics

2018, volume 2, issue 2

doi:10.21926/obm.genet.1802020

Received: March 13, 2018

Accepted: May 6, 2018

Published: May 16, 2018

Abstract:

Melanoma is a highly aggressive skin cancer with high incidence worldwide. There is growing evidence that aberrantly expressed non-coding RNAs (ncRNAs) play a role in the development, progression and dissemination of melanoma tumor cells. Among the many types of ncRNAs described in this review, the functions of micro- and long non-coding RNAs are described and related to the six hallmarks of cancer. Recently, ncRNAs discovered in body fluids have become known as one of the most promising groups of oncological biomarkers for use in non-invasive diagnosis, prognosis or therapeutic response prediction.

Keywords

Non-coding RNA; melanoma; cancer hallmarks



© 2018 by the author. This is an open access article distributed under the conditions of the [Creative Commons by Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is correctly cited.

1. Introduction

Melanoma is the deadliest form of skin cancer. The prognosis remains favorable for those diagnosed with early stage disease, where surgical therapy is the mainstay of treatment. However, treatment rarely succeeds if begun in the advanced stages of disease, although survival of late-stage disease has considerably improved in recent years [1, 2].

Interest in the identification of molecular pathways that contribute to melanoma development and progression has grown significantly over the last few years, revealing new insights for development of targeted melanoma therapies such as BRAF (B-Raf proto-oncogene, serine/threonine kinase) kinase inhibitor or MEK (mitogen-activated protein kinase kinase 1/2) inhibitor [3, 4]. Concurrently, the manipulation of the immune system [5] (e.g., through the blocking of immune checkpoints or through adoptive cell transfer) has become an important strategy for controlling disease progression during advanced melanoma stages. Although these therapies show promising clinical results, primary or acquired resistance to therapy is common and clinical outcomes for metastatic disease still remain poor.

Recently, epigenetic modifications have also been implicated in the pathogenesis of many human diseases, including melanoma. The most widely accepted definition of epigenetics includes stably inherited modulations in the expression of genes that do not involve changes in their DNA sequences. In this view, gene expression can vary due to effects exerted by RNA molecules. Specifically, increasing emphasis is being placed on the roles of non-coding RNAs (ncRNAs) to modulate gene expression through their functions as epigenetic modifiers [6]. Since ribosomal RNAs (rRNAs) and transfer RNAs (tRNAs) were discovered in the 1950s [7, 8], numerous other classes of RNAs located within the nucleus or cytoplasm have joined their ranks, including ncRNAs [9]. The two most described classes of ncRNAs are micro-RNAs (miRNAs) and the very large group of long ncRNAs (lncRNAs). Micro-RNAs, ncRNAs that are 19-22 nucleotides in length, modulate gene expression through mRNA silencing or degradation [10]. Notably, a single miRNA has the capacity to inhibit several different mRNA targets; this feature explains how miRNAs are such powerful regulators of gene expression. Moreover, they are variably expressed by cells and control many pathways that participate in cell metabolism, proliferation, differentiation and apoptosis [11]. Indeed, various studies have demonstrated that more than 60% of human genes are regulated by miRNAs [12] that may act as tumor suppressors (TS-miR) or oncogenes (oncomiR) depending on tissue, cellular context and target genes [13, 14]. The non-coding genome is also transcribed into many thousands of lncRNAs that are >200 nucleotides in length [15]. In contrast to miRNAs, which mainly negatively regulate gene expression, lncRNAs may either induce or repress gene expression and are more specific in terms of species, tissue and tumor type.

The objective of this review is to present information with regard to selected micro-RNAs and long non-coding RNAs in melanoma and relate their functions to the six hallmarks of cancer proposed by Hanahan and Weinberg in 2000 [16]. In that work, they suggested that cancer cells are characterized by six essential physiological alterations that are conducive to malignant growth: i) sustaining proliferative signaling; ii) evading growth suppressors; iii) enabling replicative immortality; iv) activating invasion and metastasis; v) inducing angiogenesis and vi) resisting cell death. This is the first review that relates short and long non-coding RNAs to current knowledge with regard to the six cancer hallmarks of cutaneous melanoma.

2. Non-Coding RNAs Maintain Proliferative Signaling in Melanoma Cells

To maintain a proliferative state, cancer cells often produce their own growth factors (GF) while concomitantly overexpressing GF receptors. Subsequently, GF bind to over-expressed GF receptors to induce autocrine proliferative stimulation, thus reducing their dependency on the external microenvironment for stimulation. Melanoma cells can also constitutively activate their own receptors via signals that are transduced within the tumor cells by activation of MAPK (mitogen-activated protein kinase) or NOTCH pathways (Figure 1). Such growth-signaling pathways in cancer cells have been linked to deregulation of the normal activities of non-coding RNA [17].




Figure 1 Differential non-coding RNA expression in normal and melanoma cells influences growth signals. Normal cells regulate their growth by modulating the activity of MAPK pathways and of the NOTCH signaling pathway. In tumor cells, aberrant expression of several micro-RNAs and long non-coding RNAs cause deregulation of these pathways and of cell proliferation. LFNG: LFNG O-fucosylpeptide 3-beta-N-acetylglucosaminyltransferase; NUMB: numb homolog; PTEN: phosphatase and tensin homolog; JNK: mitogen-activated protein kinase 8; ERK: elk-related tyrosine kinase; TGFβ: transforming growth factor beta; BANCN: BRAF-activated non-protein coding RNA; SAMMSON: survival associated mitochondrial melanoma-specific oncogenic non-coding RNA.

Non-coding RNAs can be categorized into two groups. The first group encompasses ncRNAs that normally function as tumor suppressors, but which are down-regulated and therefore not effectively present in cancer cells. For example, in non-tumor cells miR-31 targets kinases such as MET (proto-oncogene receptor tyrosine kinase), RAB27 (Ras-related protein in brain) and NIK (NFKB inducing kinase) to block signaling and inhibit cell proliferation [18]. Expression of each of these kinases is often enhanced in melanoma cells, due to the frequent deletion of the *miR-31* gene [19]. Another ncRNA, miR-204, also plays a central role in maintaining a non-proliferative state in noncancerous cells by down-regulating the NOTCH signaling pathway.

Conversely, other non-coding RNAs that act as oncogenes are over-expressed in melanoma cells. The miRNA miR-21 [20] and lncRNAs *RMEL3* [21] and *BANCN* (*BRAF-activated non-coding RNA*) [22, 23] activate the MAPK signaling pathway to induce cell proliferation even in the absence of GF. Indeed, up-regulation of miR-21 in melanoma is associated with worse prognosis [20, 24, 25]. This over-expression could be induced by environmental factors such as ultraviolet (UV)

exposure that contribute to melanomagenesis [26, 27]. Likewise, *RMEL3* has been described as an enriched lncRNA in melanoma tumors [21, 28], while *BANCR* stimulates MAPK pathways and the NOTCH cascade by sequestering miR-204 so that it is no longer able to repress NOTCH2 [22]. Meanwhile, miR-146a also participates in the regulation of this cascade by targeting LFNG (LFNG O-fucosylpeptide 3-beta-N-acetylglucosaminyltransferase) and the protein encoded by *NUMB* (numb homolog) [29, 30]. Finally, tumor cell proliferation is also enhanced in melanoma cells by the lncRNA *SAMMSON* (*survival associated mitochondrial melanoma-specific oncogenic non-coding RNA*) that confers a growth advantage to melanoma cells. *SAMMSON* mainly localizes to the cytoplasm and to a lesser degree to mitochondria, where it interacts with p32, a protein implicated in mitochondrial metabolism [31]. However, *SAMMSON* functions and downstream growth enhancement mechanisms are largely unknown.

3. Non-Coding RNAs Increase Melanoma Cell Insensitivity to Growth Suppressor Signals

Another hallmark of tumor cell development lies in cell evasion of control by growth inhibitors, resulting in sustained proliferative signaling. The principal antigrowth factor, or growth inhibitor, is TGF β (transforming growth factor beta). TGF β and other antigrowth factors bind to transmembrane cell surface receptors coupled to intracellular signaling pathways (Figure 1), leading to regulation of the cell cycle clock specifically *via* cyclins and kinases that govern cell cycle transit through the G1 phase (Figure 2).

Several ncRNAs that target proteins of the cell cycle are significantly down-regulated in melanoma cells compared to normal melanocytes, whereas in the latter they act as tumor suppressor ncRNAs. The miRNA most differentially expressed between normal melanocytes and primary melanomas, miR-211 [32- 36], targets TGF β [35] to stop the advancement of the cell cycle [32, 33, 37]. Another TS-miR that is under-expressed in melanomas and in a variety of other cancers, miR-206, inhibits translation of cyclin-dependent kinases such as CDK4 (cyclin-dependent kinase 4), cyclin D1 or cyclin C to induce G1 arrest in noncancerous cells [38]; similar effects are also achieved by miR-let-7b and miR-193b [39-41]. Likewise, tumor suppressor proteins such as those encoded by *CYLD* (cylindromatosis) [42-44] are down-regulated by at least two miRNAs that are over-expressed in melanoma cells, miR-767 [45] and miR-186 [46].

Conversely, two ncRNAs acting as a pair, miR-221 and miR-222, have been reported to drive oncogenesis of melanomas and many types of malignancies through their dynamic roles in removing inhibitory controls of proliferation and progression [47-49]. For example, both miR-221 and miR-222 have been shown to target p27, another important cell cycle regulator that induces cell cycle arrest when it binds to and blocks the function of cyclin D1 [48, 50]. Besides these miRNAs, the focally amplified lncRNA on chromosome 1 (*FALEC*) has been identified as an oncogenic lncRNA in melanoma [51] in addition to *PVT1* [52-54] and *UCA1* (urothelial cancer associated 1) [55]. *FALEC* affects p21 expression in melanoma by acting upon EZH2 (enhancer of zeste 2 polycomb repressive complex 2 subunit) [51], whereas *PVT1* enhances cell proliferation through the regulation of cyclin D1 [52] and sequestration of miR-26b [53]. The interplay between lncRNAs and miRNAs is also illustrated by the activity of the ncRNA pair *UCA1* and miR-507. The experimental depletion of *UCA1* in melanoma cell lines induces G0/G1 cell cycle arrest with increased expression of miR-507, suggesting that miR-507 directly binds *UCA1*. Moreover, the transcription factor forkhead box protein M1 (FOXO1), an essential effector of G2/M phase

transition, is also a direct target of miR-507. In melanoma, the elevated amount of *UCA1* sequesters miR-507, rendering the latter unable to repress FOXM1 [55].




Figure 2 Non-coding RNAs in cell cycle control of normal and melanoma cells. Cyclin-dependent kinases (CDK) trigger the transition from G1 to S phase and from G2 to M phase by phosphorylating distinct sets of substrates. Cyclin D family members and CDK4,6 regulate events in the G1 phase, cyclin E/CDK2 triggers the S phase, cyclin A/CDK2 regulates the completion of the S phase and cyclin B/CDK1 is essential for driving cells into mitosis. In melanoma cells, aberrant expression of micro- and long ncRNAs activates mainly the cyclin D/CDK4,6 pair of ncRNAs, leading to progression from G1 to S phase. FOXM1: forkhead box M1; *FALEC*: focally amplified long non-coding RNA in epithelial cancer; *PVT1*: Pvt1 oncogene; CYLD: cylindromatosis; *UCA1*: urothelial cancer associated 1.

4. Non-Coding RNAs Enable Replicative Immortality by Circumvention of Senescence in Melanoma Cells

Uncontrolled cellular proliferation is a defining feature of cancer. In contrast, cellular senescence is a state of irreversible growth arrest which can be triggered by various stimuli such as telomere shortening, DNA damage, oxidative stress or oncogene activation (Figure 3). E2F family members are critical barriers to senescence induction that allow a cell to avoid senescence either through activation or inhibition of cell replication.

Some ncRNAs promote senescence in normal cells under selective stimuli, such as miR-205 [56] or miR-377 [57]. In normal cells, these ncRNAs target E2F family members and induce a senescence phenotype with elevated expression of p16INK4A, while both are down-regulated in melanoma cells [56, 57]. *UCA1*, previously described in the regulation of the cell cycle, is also able to induce senescence by sequestering hnRNP A1 (heterogeneous nuclear ribonucleoprotein A1) and by increasing p16INK4A mRNA stability to prevent its degradation [58]. Conversely, Montes et

al. identified in melanoma patients a negative correlation between the lncRNA *MIR31HG* (MIR31 host gene) and p16INK4A [59]. Meanwhile, other studies suggest that p16INK4A could be also regulated by *ANRIL* (CDKN2B antisense RNA 1) through the recruitment of chromatin remodeling factors to its gene locus [60-62].




Figure 3 Regulation of cell senescence by non-coding RNAs in normal and melanoma cells. In normal cells responding to several signals, tumor suppressors p19ARF and p16INK4A are activated to prevent abnormal cell proliferation. In melanoma cells, p16INK4A is up-regulated by the expression of lncRNAs *MIR31HG* and *ANRIL* and upon down-regulation of miR-205, miR-377 and *UCA1*. p19ARF: cyclin-dependent kinase inhibitor 2A; Mdm2: MDM2 proto-oncogene, E3 ubiquitin protein ligase; *UCA1*: urothelial cancer associated 1; p16INK4A: cyclin-dependent kinase inhibitor 2A; hnRNPA1: heterogeneous nuclear ribonucleoprotein A1; CDK: cyclin-dependent kinase; Rb: retinoblastoma; *MIR31HG*: MIR31 host gene; *ANRIL*: CDKN2B antisense RNA 1.

5. Non-Coding RNAs Promote Melanoma Invasion and Metastasis

Melanoma cells are characterized by genetic instability and invasiveness through acquisition of migratory capacity. Several classes of proteins implicated as cell-cell adhesion molecules have been shown to be involved, such as immunoglobulin family members, calcium-dependent cadherin proteins or integrins, the latter of which link cells to the extracellular matrix. The metastatic capabilities of melanoma cancer cells also involve extracellular proteases such as MMP (matrix metalloproteinase) family members that are implicated in the breakdown of the extracellular matrix (Figure 4).

All of these pathways are regulated by ncRNAs. Indeed, miR-137 is down-regulated in melanoma cells [63-65] and represses transcription of the gene for E-cadherin by targeting the

transcription factor TBX3 (T-box transcription factor 3) [66]. Meanwhile, integrin-dependent pathways that are regulated by miR-let-7a are also down-regulated in malignant melanoma cells [67]. Moreover, MMP proteins are regulated by the lncRNA *GAS5* (growth arrest-specific 5) [68] and are also down-regulated in melanoma tissues [69], resulting in inhibition of their effects.

Conversely, numerous ncRNAs increase cell invasion by targeting the same pathways that function in cell migration. Thus, miR-21 (previously described in the first hallmark) targets TIMP3 (TIMP metalloproteinase inhibitor 3), a protein that inhibits MMP synthesis and activity [37, 70]. MMP family members are also regulated by the lncRNA *SLNCR1* (steroid receptor RNA activator 1-like non-coding RNA) through their interactions with Brn3a (brain-specific homeobox protein 3a) and AR (androgen receptor) transcription factors [71]. Metastasis-associated lung adenocarcinoma transcript 1 (*MALAT1*), first discovered in lung cancer, is also up-regulated in melanoma tumors [72]. *MALAT1* similarly regulates extracellular proteases, while also inducing migration by sequestering miR-22 [73] and miR183 [74] through respective targeting of MMP14 and IGTB1 (Integrin β 1). In addition to *MALAT1* regulation, integrin pathways are also controlled by the ncRNA pair miR-221/miR-222 through enhancement of cellular migration *via* activation of the Akt/PI3K signaling pathway [50].




Figure 4 Role of non-coding RNAs in cell invasion and migration. In a melanoma cancer cell, integrin binding to ligands in the extracellular matrix triggers multiple signals leading to changes in gene expression and cell migration. The loss of the epithelial cell-cell adhesion molecule E-cadherin also participates by inducing cell migration. PI3K: phosphatidylinositol-4, 5-bisphosphate 3-kinase; Akt: v-akt murine thymoma viral oncogene homolog; TBX3: T-box 3; MMP: matrix metalloproteinase; *GAS5*: growth arrest-specific 5; TIMP3: TIMP metalloproteinase inhibitor 3; *HOTAIR*: HOX transcript antisense RNA; *MALAT1*: metastasis-associated lung adenocarcinoma transcript 1.

Overall, *HOX transcript antisense RNA (HOTAIR)* promotes melanoma cell invasion and migration by regulating the epithelial-to-mesenchymal transition (EMT) through sequestration of miR-152-3p [75]. *HOTAIR* can also regulate target gene expression by acting as a guide for PRC2 (polycomb repressive complex 2), leading to the epigenetic silencing of metastasis suppressor genes [76]. Notably, *HOTAIR* levels are elevated in melanoma cells and are associated with poor prognosis [75, 77].

6. Non-Coding RNAs Induce Angiogenesis in Melanoma Tumors

Oxygen and nutrients supplied by the vasculature are crucial for tumor cell development and proliferation. The formation of new blood vessels is important for tumors to obtain an adequate supply of oxygen and nutrients and to evacuate metabolic waste and undergo metastatic processes (Figure 5). The principal regulator of this vascularization process is vascular endothelial growth factor (VEGF), a protein that is up-regulated by hypoxia and oncogenic signaling pathways.

Few non-coding RNAs have been described in melanoma angiogenesis; most of them target, either directly or indirectly, hypoxia inducible factor α (HIF1 α) [78, 79]. It has been suggested that because miR-21 is highly expressed in melanoma cells [25], it may target PTEN (phosphatase and tensin homolog) to activate MAPK signaling. In this way, PTEN may thereby enhance HIF1 α expression and VEGF transcription, as observed in prostate cancer cells [79]. Similarly, Mir-21 may also enhance angiogenesis *via* its target TIMP3 [80], while a previous study showed that miR-424 enhances VEGF expression by down-regulating cullin 2 (CUL2), a protein that stabilizes HIF α within a sequestered cytoplasmic complex [81].




Figure 5 Role of micro-RNAs in angiogenesis. In the melanoma cell under normal oxygen conditions, HIF1 α is degraded. Under hypoxia, HIF accumulates in the cytoplasm and moves to the nucleus to activate the production of proteins implicated in angiogenesis, including growth factor VEGF. Two micro-RNAs facilitate protein production by activating transcription of HIF1A and its accumulation in the cytoplasm. PTEN: phosphatase and tensin homolog; HIF1A: hypoxia inducible factor 1, alpha subunit; CUL2: cullin 2; VEGF: vascular endothelial growth factor A; ILK: integrin-linked kinase; PI3K: phosphatidylinositol-4, 5-bisphosphate 3-kinase.

7. Non-Coding RNAs Allow Evasion of Programmed Cell Death of Melanoma Cells

The ability of tumor cells to proliferate also stems from their resistance to programmed cell death. Apoptosis is dependent upon sensors which control the intra- and extracellular microenvironment, such as cell surface receptors and effectors which trigger cell death. Intracellular signals of apoptosis converge to act within mitochondria, where Bcl-2 family

members play their anti- or pro-apoptotic roles through preventing or promoting the release of cytochrome C, respectively, with the latter leading to the activation of effector caspases. At least three TS-miRNAs, miR-219 [82], miR-15b [83, 84] and miR-181a [85], enhance apoptosis by targeting Bcl-2. All of these TS-miRNAs are down-regulated in melanoma cells.

Overall, the lncRNA *GAS5* regulates apoptosis in melanoma *via* its interaction with the DNA binding domain of GRs (glucocorticoid receptors) by preventing binding of GRs to glucocorticoid response elements (GREs) in the genome. This suppresses the induction of several response genes implicated in apoptosis [86]. Finally, *SPRY4-IT1* (*SPRY4-IT1 intronic transcript 1*) is over-represented in melanoma cells [87, 88] and may block apoptosis by modulating lipid metabolism and lipotoxicity [89].




Figure 6 Role of non-coding RNAs in apoptosis. Apoptosis is induced by the activation of death receptors (extracellular signals) or intracellular signals, leading to the activation of Bcl-2 family members. Next, cytochrome c release activates effector caspases. In the nucleus, most Bcl-2 family genes are regulated by the binding of glucocorticoid receptor (GR) to their DNA sequences. BIM: BCL2-like 11; PUMA: BCL2 binding component 3; *GAS5*: growth arrest-specific 5; Bcl-2: B-cell CLL/lymphoma 2.

8. Conclusions

Among more than 3,000 miRNAs [90] and 24,793 annotated lncRNAs that have been identified within the human genome [91], only a few show deregulations in cutaneous melanoma (Table 1). Nevertheless, we are only at the beginning of our understanding and identification of ncRNAs; recent computational analyses using high-throughput expression data, including microarrays and RNA sequencing, have helped us identify deregulated mRNAs, lncRNAs and pseudogenes in melanoma [92, 93]. Consequently, the growing body of evidence strongly suggests a role of pseudogene transcription as a source of endogenous siRNAs that either bind to their cognate wild-type (WT) genes or act as sponges by binding miRNAs targeting their WT genes. The emerging picture underscores the functional interdependence of coding and non-coding RNAs [94, 95] whereby all perturbations in these interactions have widespread consequences that can lead to the malignant phenotypes of cancer cells.

Table 1 Non-coding RNAs deregulated in melanoma

Name	Functions in tumor	Targets	Up/down-regulated in melanoma
<i>lncRNA</i>			
<i>ANRIL</i>	Senescence	p16INK4A	Up
<i>BANCR</i>	Proliferation	mir-204	Up
<i>FALEC</i>	Cell cycle	p21	Up
<i>GAS5</i>	Invasion / Metastasis / Apoptosis	MMP, GR	Up
<i>HOTAIR</i>	Invasion / Metastasis	PRC2	Up
<i>MALAT1</i>	Invasion / Metastasis	miR-22, miR-183	Up
<i>MIR31HG</i>	Senescence	p16INK4A	Up
<i>PVT1</i>	Cell cycle	Cyclin D1, miR-26b	Up
<i>RMEL3</i>	Proliferation		Up
<i>SAMMSON</i>	Proliferation	p32	Up
<i>SLNCR1</i>	Invasion / Metastasis	Brn3a, AR	Up
<i>SPRY4-IT1</i>	Apoptosis		Up
<i>UCA1</i>	Cell cycle / Senescence	miR-507, hnRNPA1	Up
<i>miRNA</i>			
miR-137	Invasion / Metastasis	TBX3	Down
miR-146a	Proliferation	LFNG, NUMB	Up
miR-15b	Apoptosis	Bcl-2 family members	Down
miR-181a	Apoptosis	Bcl-2 family members	Down
miR-186	Cell cycle	CYLD	Up
miR-193b	Cell cycle	CDK4, Cyclin D1, Cyclin C	Down
miR-204	Proliferation		Down
miR-205	Senescence	E2F members	Down
miR-206	Cell cycle	CDK4, Cyclin D1, Cyclin C	Down
mir-21	Proliferation / Invasion / Metastasis / Angiogenesis	TIMP3, PTEN	Up
miR-211	Growth	TGFB	Down
miR-219	Apoptosis	Bcl-2 family members	Down
miR-221 / 222	Cell cycle / Invasion / Metastasis	p27, Akt/PI3K	Up
miR-31	Proliferation	MET, RAB27, NIK	Down
miR-377	Senescence	E2F members	Down
miR-424	Angiogenesis	CUL2	Up
miR-767	Cell cycle	CYLD	Up
miR-let-7a	Invasion / Metastasis	Integrin	Down
miR-let-7b	Cell cycle	CDK4, Cyclin D1, Cyclin C	Down

A more complete understanding of the cellular biology of ncRNAs, notably lncRNAs, is crucial to improving early diagnosis, prognosis and treatment. The latter, of course, first requires detailed knowledge about melanoma-specific lncRNAs functions. However, therapeutic modulation of cancer- and tissue-specific expression of lncRNAs could have major advantages over other therapeutics.

We also must consider the timing of action of ncRNAs, since many of the studies described in this review only investigated ncRNAs present at the time of diagnosis. Further studies need to be done to determine the variation in ncRNAs levels throughout the disease course to further evaluate their relevance to survival and treatment.

To conclude, the study of ncRNA effects on melanoma is a very attractive undertaking, since they participate in numerous cancer processes. Notably, because ncRNAs could be released into the bloodstream, they may serve as non-invasive biomarkers of melanoma activity.

Author Contributions

VV and DF wrote the manuscript and designed figures.

Competing Interests

The authors have declared that no competing interests exist.

References

1. Abdel-Rahman O. Evaluation of the eighth American Joint Committee on Cancer staging system for malignant melanoma of the skin. *Future Oncol.* 2018; 14: 471-481.
2. Miller KD, Siegel RL, Lin CC, Mariotto AB, Kramer JL, Rowland JH, et al. Cancer treatment and survivorship statistics, 2016. *CA Cancer J Clin.* 2016; 66: 271-289.
3. Kuske M, Rauschenberg R, Garzarolli M, Meredyth-Stewart M, Beissert S, Troost EGC, et al. Melanoma Brain Metastases: Local Therapies, Targeted Therapies, Immune Checkpoint Inhibitors and Their Combinations-Chances and Challenges. *Am J Clin Dermatol.* 2018.
4. Najem A, Krayem M, Perdrix A, Kerger J, Awada A, Journe F, et al. New Drug Combination Strategies in Melanoma: Current Status and Future Directions. *Anticancer Res.* 2017; 37: 5941-5953.
5. Marconcini R, Spagnolo F, Stucci LS, Ribero S, Marra E, Rosa F, et al. Current status and perspectives in immunotherapy for metastatic melanoma. *Oncotarget.* 2018; 9: 12452-12470.
6. Lee JT. Epigenetic regulation by long noncoding RNAs. *Science.* 2012; 338: 1435-1439.
7. Palade GE. A small particulate component of the cytoplasm. *J Biophys Biochem Cytol.* 1955; 1: 59-68.
8. Hoagland MB, Stephenson ML, Scott JF, Hecht LI, Zamecnik PC. A soluble ribonucleic acid intermediate in protein synthesis. *J Biol Chem.* 1958; 231: 241-257.
9. Morris KV, Mattick JS. The rise of regulatory RNA. *Nat Rev Genet.* 2014; 15: 423-437.
10. Djuranovic S, Nahvi A, Green R. A parsimonious model for gene regulation by miRNAs. *Science.* 2011; 331: 550-553.
11. Lee Y, Jeon K, Lee JT, Kim S, Kim VN. MicroRNA maturation: stepwise processing and subcellular localization. *Embo J.* 2002; 21: 4663-4670.

12. Friedman RC, Farh KK, Burge CB, Bartel DP. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res.* 2009; 19: 92-105.
13. Esquela-Kerscher A, Slack FJ. Oncomirs - microRNAs with a role in cancer. *Nat Rev Cancer.* 2006; 6: 259-269.
14. Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell.* 2005; 120 : 15-20.
15. Derrien T, Johnson R, Bussotti G, Tanzer A, Djebali S, Tilgner H, et al. The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. *Genome Res.* 2012; 22: 1775-1789.
16. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell.* 2000; 100: 57-70.
17. Zhang W, Liu HT. MAPK signal pathways in the regulation of cell proliferation in mammalian cells. *Cell Res.* 2002; 12: 9-18.
18. Asangani IA, Harms PW, Dodson L, Pandhi M, Kunju LP, Maher CA, et al. Genetic and epigenetic loss of microRNA-31 leads to feed-forward expression of EZH2 in melanoma. *Oncotarget.* 2012; 3: 1011-1025.
19. Poliseno L, Haimovic A, Segura MF, Hanniford D, Christos PJ, Darvishian F, et al. Histology-specific microRNA alterations in melanoma. *J Invest Dermatol.* 2012; 132: 1860-1868.
20. Jiang L, Lv X, Li J, Li J, Li X, Li W, et al. The status of microRNA-21 expression and its clinical significance in human cutaneous malignant melanoma. *Acta Histochem.* 2012; 114: 582-588.
21. Goedert L, Pereira CG, Roszik J, Placa JR, Cardoso C, Chen G, et al. RMEL3, a novel BRAFV600E-associated long noncoding RNA, is required for MAPK and PI3K signaling in melanoma. *Oncotarget.* 2016; 7: 36711-36718.
22. Cai B, Zheng Y, Ma S, Xing Q, Wang X, Yang B, et al. BANCER contributes to the growth and invasion of melanoma by functioning as a competing endogenous RNA to upregulate Notch2 expression by sponging miR204. *Int J Oncol.* 2017; 51: 1941-1951.
23. Li R, Zhang L, Jia L, Duan Y, Li Y, Bao L, et al. Long non-coding RNA BANCER promotes proliferation in malignant melanoma by regulating MAPK pathway activation. *PLoS One.* 2014; 9: e100893.
24. Grignol V, Fairchild ET, Zimmerer JM, Lesinski GB, Walker MJ, Magro CM, et al. miR-21 and miR-155 are associated with mitotic activity and lesion depth of borderline melanocytic lesions. *Br J Cancer.* 2011; 105: 1023-1029.
25. Satzger I, Mattern A, Kuettler U, Weinspach D, Niebuhr M, Kapp A, et al. microRNA-21 is upregulated in malignant melanoma and influences apoptosis of melanocytic cells. *Exp Dermatol.* 2012; 21: 509-514.
26. Yang CH, Yue J, Pfeffer SR, Handorf CR, Pfeffer LM. MicroRNA miR-21 regulates the metastatic behavior of B16 melanoma cells. *J Biol Chem.* 2011; 286: 39172-39178.
27. Melnik BC. MiR-21: an environmental driver of malignant melanoma? *J Transl Med.* 2015; 13: 202.
28. Sousa JF, Torrieri R, Silva RR, Pereira CG, Valente V, Torrieri E, et al. Novel primate-specific genes, RMEL 1, 2 and 3, with highly restricted expression in melanoma, assessed by new data mining tool. *PLoS One.* 2010; 5: e13510.
29. Forloni M, Dogra SK, Dong Y, Conte D, Jr., Ou J, Zhu LJ, et al. miR-146a promotes the initiation and progression of melanoma by activating Notch signaling. *Elife.* 2014; 3: e01460.
30. Raimo M, Orso F, Grassi E, Cimino D, Penna E, De Pitta C, et al. miR-146a Exerts Differential Effects on Melanoma Growth and Metastatization. *Mol Cancer Res.* 2016; 14: 548-562.

31. Leucci E, Vendramin R, Spinazzi M, Laurette P, Fiers M, Wouters J, et al. Melanoma addiction to the long non-coding RNA SAMMSON. *Nature*. 2016; 531: 518-522.
32. Aftab MN, Dinger ME, Perera RJ. The role of microRNAs and long non-coding RNAs in the pathology, diagnosis, and management of melanoma. *Arch Biochem Biophys*. 2014; 563: 60-70.
33. Weiss SA, Hanniford D, Hernando E, Osman I. Revisiting determinants of prognosis in cutaneous melanoma. *Cancer*. 2015; 121: 4108-4123.
34. Boyle GM, Woods SL, Bonazzi VF, Stark MS, Hacker E, Aoude LG, et al. Melanoma cell invasiveness is regulated by miR-211 suppression of the BRN2 transcription factor. *Pigment Cell Melanoma R*. 2011; 24: 525-537.
35. Levy C, Khaled M, Iliopoulos D, Janas MM, Schubert S, Pinner S, et al. Intronic miR-211 assumes the tumor suppressive function of its host gene in melanoma. *Mol Cell*. 2010; 40: 841-849.
36. Mazar J, DeYoung K, Khaitan D, Meister E, Almodovar A, Goydos J, et al. The regulation of miRNA-211 expression and its role in melanoma cell invasiveness. *PLoS One*. 2010; 5: e13779.
37. Latchana N, Ganju A, Howard JH, Carson WE, 3rd. MicroRNA dysregulation in melanoma. *Surg Oncol*. 2016; 25: 184-189.
38. Georgantas RW, 3rd, Streicher K, Luo X, Greenlees L, Zhu W, Liu Z, et al. MicroRNA-206 induces G1 arrest in melanoma by inhibition of CDK4 and Cyclin D. *Pigment Cell Melanoma R*. 2014; 27: 275-286.
39. Chen J, Feilotter HE, Pare GC, Zhang X, Pemberton JG, Garady C, et al. MicroRNA-193b represses cell proliferation and regulates cyclin D1 in melanoma. *Am J Pathol*. 2010; 176: 2520-2529.
40. Schultz J, Lorenz P, Gross G, Ibrahim S, Kunz M. MicroRNA let-7b targets important cell cycle molecules in malignant melanoma cells and interferes with anchorage-independent growth. *Cell Res*. 2008; 18: 549-557.
41. Xu D, Tan J, Zhou M, Jiang B, Xie H, Nie X, et al. Let-7b and microRNA-199a inhibit the proliferation of B16F10 melanoma cells. *Oncol Lett*. 2012; 4: 941-946.
42. Pannem RR, Dorn C, Ahlqvist K, Bosserhoff AK, Hellerbrand C, Massoumi R. CYLD controls c-MYC expression through the JNK-dependent signaling pathway in hepatocellular carcinoma. *Carcinogenesis*. 2014; 35: 461-468.
43. Xia JT, Chen LZ, Jian WH, Wang KB, Yang YZ, He WL, et al. MicroRNA-362 induces cell proliferation and apoptosis resistance in gastric cancer by activation of NF-kappaB signaling. *J Transl Med*. 2014; 12: 33.
44. Yang Y, Zhou J. CYLD - a deubiquitylase that acts to fine-tune microtubule properties and functions. *J Cell Sci*. 2016; 129: 2289-2295.
45. Zhang K, Guo L. MiR-767 promoted cell proliferation in human melanoma by suppressing CYLD expression. *Gene*. 2018; 641: 272-278.
46. Qiu H, Yuan S, Lu X. miR-186 suppressed CYLD expression and promoted cell proliferation in human melanoma. *Oncol Lett*. 2016; 12: 2301-2306.
47. Garofalo M, Quintavalle C, Romano G, Croce CM, Condorelli G. miR221/222 in cancer: their role in tumor progression and response to therapy. *Curr Mol Med*. 2012; 12: 27-33.
48. Felicetti F, Errico MC, Bottero L, Segnalini P, Stoppacciaro A, Biffoni M, et al. The promyelocytic leukemia zinc finger-microRNA-221/-222 pathway controls melanoma progression through multiple oncogenic mechanisms. *Cancer Res*. 2008; 68: 2745-2754.

49. Errico MC, Felicetti F, Bottero L, Mattia G, Boe A, Felli N, et al. The abrogation of the HOXB7/PBX2 complex induces apoptosis in melanoma through the miR-221&222-c-FOS pathway. *Int J Cancer*. 2013; 133: 879-892.
50. Felicetti F, De Feo A, Coscia C, Puglisi R, Pedini F, Pasquini L, et al. Exosome-mediated transfer of miR-222 is sufficient to increase tumor malignancy in melanoma. *J Transl Med*. 2016; 14: 56.
51. Ni N, Song H, Wang X, Xu X, Jiang Y, Sun J. Up-regulation of long noncoding RNA FALEC predicts poor prognosis and promotes melanoma cell proliferation through epigenetically silencing p21. *Biomed Pharmacother*. 2017; 96: 1371-1379.
52. Chen L, Ma D, Li Y, Li X, Zhao L, Zhang J, et al. Effect of long non-coding RNA PVT1 on cell proliferation and migration in melanoma. *Int J Mol Med*. 2018; 41: 1275-1282.
53. Wang BJ, Ding HW, Ma GA. Long Noncoding RNA PVT1 Promotes Melanoma Progression Via Endogenous Sponging MiR-26b. *Oncol Res*. 2017.
54. Chen X, Gao G, Liu S, Yu L, Yan D, Yao X, et al. Long Noncoding RNA PVT1 as a Novel Diagnostic Biomarker and Therapeutic Target for Melanoma. *Biomed Res Int*. 2017; 2017: 7038579.
55. Wei Y, Sun Q, Zhao L, Wu J, Chen X, Wang Y, et al. LncRNA UCA1-miR-507-FOXO1 axis is involved in cell proliferation, invasion and G0/G1 cell cycle arrest in melanoma. *Med Oncol*. 2016; 33: 88.
56. Dar AA, Majid S, de Semir D, Nosrati M, Bezrookove V, Kashani-Sabet M. miRNA-205 suppresses melanoma cell proliferation and induces senescence via regulation of E2F1 protein. *J Biol Chem*. 2011; 286: 16606-16614.
57. Zehavi L, Schayek H, Jacob-Hirsch J, Sidi Y, Leibowitz-Amit R, Avni D. MiR-377 targets E2F3 and alters the NF- κ B signaling pathway through MAP3K7 in malignant melanoma. *Mol Cancer*. 2015; 14: 68.
58. Kumar PP, Emechebe U, Smith R, Franklin S, Moore B, Yandell M, et al. Coordinated control of senescence by lncRNA and a novel T-box3 co-repressor complex. *Elife*. 2014; 3.
59. Montes M, Nielsen MM, Maglieri G, Jacobsen A, Hojfeldt J, Agrawal-Singh S, et al. The lncRNA MIR31HG regulates p16(INK4A) expression to modulate senescence. *Nat Commun*. 2015; 6: 6967.
60. Yap KL, Li S, Munoz-Cabello AM, Raguz S, Zeng L, Mujtaba S, et al. Molecular interplay of the noncoding RNA ANRIL and methylated histone H3 lysine 27 by polycomb CBX7 in transcriptional silencing of INK4a. *Mol Cell*. 2010; 38: 662-674.
61. Cunningham MS, Santibanez Koref M, Mayosi BM, Burn J, Keavney B. Chromosome 9p21 SNPs Associated with Multiple Disease Phenotypes Correlate with ANRIL Expression. *PLoS Genet*. 2010; 6: e1000899.
62. Xu S, Wang H, Pan H, Shi Y, Li T, Ge S, et al. ANRIL lncRNA triggers efficient therapeutic efficacy by reprogramming the aberrant INK4-hub in melanoma. *Cancer Lett*. 2016; 381: 41-48.
63. Luo C, Tetteh PW, Merz PR, Dickes E, Abukiwan A, Hotz-Wagenblatt A, et al. miR-137 inhibits the invasion of melanoma cells through downregulation of multiple oncogenic target genes. *J Invest Dermatol*. 2013; 133: 768-775.
64. Bennett PE, Bemis L, Norris DA, Shellman YG. miR in melanoma development: miRNAs and acquired hallmarks of cancer in melanoma. *Physiol Genomics*. 2013; 45: 1049-1059.
65. Li N. Low Expression of Mir-137 Predicts Poor Prognosis in Cutaneous Melanoma Patients. *Med Sci Monit*. 2016; 22: 140-144.
66. Rodriguez M, Aladowicz E, Lanfrancone L, Goding CR. Tbx3 represses E-cadherin expression and enhances melanoma invasiveness. *Cancer Res*. 2008; 68: 7872-7881.

67. Muller DW, Bosserhoff AK. Integrin beta 3 expression is regulated by let-7a miRNA in malignant melanoma. *Oncogene*. 2008; 27: 6698-6706.
68. Chen L, Yang H, Xiao Y, Tang X, Li Y, Han Q, et al. Lentiviral-mediated overexpression of long non-coding RNA GAS5 reduces invasion by mediating MMP2 expression and activity in human melanoma cells. *Int J Oncol*. 2016; 48: 1509-1518.
69. Bian D, Shi W, Shao Y, Li P, Song G. Long non-coding RNA GAS5 inhibits tumorigenesis via miR-137 in melanoma. *Am J Transl Res*. 2017; 9: 1509-1520.
70. Martin del Campo SE, Latchana N, Levine KM, Grignol VP, Fairchild ET, Jaime-Ramirez AC, et al. MiR-21 enhances melanoma invasiveness via inhibition of tissue inhibitor of metalloproteinases 3 expression: in vivo effects of MiR-21 inhibitor. *PLoS One*. 2015; 10: e0115919.
71. Schmidt K, Joyce CE, Buquicchio F, Brown A, Ritz J, Distel RJ, et al. The lncRNA SLNCR1 Mediates Melanoma Invasion through a Conserved SRA1-like Region. *Cell Rep*. 2016; 15: 2025-2037.
72. Tian Y, Zhang X, Hao Y, Fang Z, He Y. Potential roles of abnormally expressed long noncoding RNA UCA1 and Malat-1 in metastasis of melanoma. *Melanoma Res*. 2014; 24: 335-341.
73. Luan W, Li L, Shi Y, Bu X, Xia Y, Wang J, et al. Long non-coding RNA MALAT1 acts as a competing endogenous RNA to promote malignant melanoma growth and metastasis by sponging miR-22. *Oncotarget*. 2016; 7: 63901-63912.
74. Sun Y, Cheng H, Wang G, Yu G, Zhang D, Wang Y, et al. Deregulation of miR-183 promotes melanoma development via lncRNA MALAT1 regulation and ITGB1 signal activation. *Oncotarget*. 2017; 8: 3509-3518.
75. Luan W, Li R, Liu L, Ni X, Shi Y, Xia Y, et al. Long non-coding RNA HOTAIR acts as a competing endogenous RNA to promote malignant melanoma progression by sponging miR-152-3p. *Oncotarget*. 2017; 8: 85401-85414.
76. Wu L, Murat P, Matak-Vinkovic D, Murrell A, Balasubramanian S. Binding interactions between long noncoding RNA HOTAIR and PRC2 proteins. *Biochemistry*. 2013; 52: 9519-9527.
77. Cantile M, Scognamiglio G, Marra L, Aquino G, Botti C, Falcone MR, et al. HOTAIR role in melanoma progression and its identification in the blood of patients with advanced disease. *J Cell Physiol*. 2017; 232: 3422-3432.
78. Hwang HW, Baxter LL, Loftus SK, Cronin JC, Trivedi NS, Borate B, et al. Distinct microRNA expression signatures are associated with melanoma subtypes and are regulated by HIF1A. *Pigment Cell Melanoma Res*. 2014; 27: 777-787.
79. Liu LZ, Li C, Chen Q, Jing Y, Carpenter R, Jiang Y, et al. MiR-21 induced angiogenesis through AKT and ERK activation and HIF-1alpha expression. *PLoS One*. 2011; 6: e19139.
80. Qi JH, Ebrahim Q, Moore N, Murphy G, Claesson-Welsh L, Bond M, et al. A novel function for tissue inhibitor of metalloproteinases-3 (TIMP3): inhibition of angiogenesis by blockage of VEGF binding to VEGF receptor-2. *Nat Med*. 2003; 9: 407-415.
81. Ghosh G, Subramanian IV, Adhikari N, Zhang X, Joshi HP, Basi D, et al. Hypoxia-induced microRNA-424 expression in human endothelial cells regulates HIF-alpha isoforms and promotes angiogenesis. *J Clin Invest*. 2010; 120: 4141-4154.
82. Long J, Menggen Q, Wuren Q, Shi Q, Pi X. MiR-219-5p Inhibits the Growth and Metastasis of Malignant Melanoma by Targeting BCL-2. *Biomed Res Int*. 2017; 2017: 9032502.
83. Xia L, Zhang D, Du R, Pan Y, Zhao L, Sun S, et al. miR-15b and miR-16 modulate multidrug resistance by targeting BCL2 in human gastric cancer cells. *Int J Cancer*. 2008; 123: 372-379.

84. Friedman EB, Shang S, de Miera EV, Fog JU, Teilum MW, Ma MW, et al. Serum microRNAs as biomarkers for recurrence in melanoma. *J Transl Med.* 2012; 10: 155.
85. Du M, Zhang Z, Gao T. Piceatannol induced apoptosis through up-regulation of microRNA-181a in melanoma cells. *Biol Res.* 2017; 50: 36.
86. Kino T, Hurt DE, Ichijo T, Nader N, Chrousos GP. Noncoding RNA gas5 is a growth arrest- and starvation-associated repressor of the glucocorticoid receptor. *Sci Signal.* 2010; 3: ra8.
87. Khaitan D, Dinger ME, Mazar J, Crawford J, Smith MA, Mattick JS, et al. The melanoma-upregulated long noncoding RNA SPRY4-IT1 modulates apoptosis and invasion. *Cancer Res.* 2011; 71: 3852-3862.
88. Liu T, Shen SK, Xiong JG, Xu Y, Zhang HQ, Liu HJ, et al. Clinical significance of long noncoding RNA SPRY4-IT1 in melanoma patients. *FEBS Open Bio.* 2016; 6: 147-154.
89. Mazar J, Zhao W, Khalil AM, Lee B, Shelley J, Govindarajan SS, et al. The functional characterization of long noncoding RNA SPRY4-IT1 in human melanoma cells. *Oncotarget.* 2014; 5: 8959-8969.
90. Londin E, Loher P, Telonis AG, Quann K, Clark P, Jing Y, et al. Analysis of 13 cell types reveals evidence for the expression of numerous novel primate- and tissue-specific microRNAs. *Proc Natl Acad Sci USA.* 2015; 112: E1106-1115.
91. Ning Q, Li Y, Wang Z, Zhou S, Sun H, Yu G. The Evolution and Expression Pattern of Human Overlapping lncRNA and Protein-coding Gene Pairs. *Sci Rep.* 2017; 7: 42775.
92. Capobianco E, Valdes C, Sarti S, Jiang Z, Poliseno L, Tsinoremas NF. Ensemble Modeling Approach Targeting Heterogeneous RNA-Seq data: Application to Melanoma Pseudogenes. *Sci Rep.* 2017; 7: 17344.
93. Zhao H, Li Y, Wang S, Yang Y, Wang J, Ruan X, et al. Whole transcriptome RNA-seq analysis: tumorigenesis and metastasis of melanoma. *Gene.* 2014; 548: 234-243.
94. Kalyana-Sundaram S, Kumar-Sinha C, Shankar S, Robinson DR, Wu YM, Cao X, et al. Expressed pseudogenes in the transcriptional landscape of human cancers. *Cell.* 2012; 149: 1622-1634.
95. Milligan MJ, Harvey E, Yu A, Morgan AL, Smith DL, Zhang E, et al. Global Intersection of Long Non-Coding RNAs with Processed and Unprocessed Pseudogenes in the Human Genome. *Front Genet.* 2016; 7: 26.



Enjoy *OBM Genetics* by:

1. [Submitting a manuscript](#)
2. [Joining in volunteer reviewer bank](#)
3. [Joining Editorial Board](#)
4. [Guest editing a special issue](#)

For more details, please visit:

<http://www.lidsen.com/journals/genetics>