



HAL
open science

Efficient treatment of a metastatic melanoma patient with a combination of BRAF and MEK inhibitors based on circulating tumor DNA analysis: a case report

Gaëlle Quéreux, Guillaume Herbreteau, Anne-Chantal Knol, Audrey Vallée, Amir Khammari, Sandrine Théoleyre, Mélanie Saint-Jean, Brigitte Dréno, Marc G. Denis

► To cite this version:

Gaëlle Quéreux, Guillaume Herbreteau, Anne-Chantal Knol, Audrey Vallée, Amir Khammari, et al.. Efficient treatment of a metastatic melanoma patient with a combination of BRAF and MEK inhibitors based on circulating tumor DNA analysis: a case report. *BMC Biochemistry*, 2017, 10 (1), pp.320. 10.1186/s13104-017-2650-5 . inserm-01807889

HAL Id: inserm-01807889

<https://inserm.hal.science/inserm-01807889v1>

Submitted on 5 Jun 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CASE REPORT

Open Access



Efficient treatment of a metastatic melanoma patient with a combination of BRAF and MEK inhibitors based on circulating tumor DNA analysis: a case report

Gaëlle Quéreux^{1,2,3}, Guillaume Herbreteau^{2,4}, Anne-Chantal Knol^{2,3}, Audrey Vallée^{2,4}, Amir Khammari^{1,2,3}, Sandrine Théoleyre^{2,4}, Mélanie Saint-Jean^{1,2,3}, Brigitte Dréno^{1,2,3} and Marc G. Denis^{2,4*} 

Abstract

Background: Fixed tissues are the standard samples used in routine practice for molecular testing. But sometimes tissues are lacking or difficult to obtain. In these cases, circulating tumor DNA released from tumor cells can be used as an alternative source of tumor DNA.

Case presentation: We present the case of a 63-year-old Caucasian woman with a metastatic melanoma and a very poor performance status. A plasma sample was tested and the BRAF p.V600E mutation was detected. Based on this result, a treatment combining a BRAF inhibitor and a MEK inhibitor was immediately started. This patient achieved a complete response. In addition, by repeating the plasma test, we could obtain a precise kinetic of release of mutated BRAF DNA in plasma.

Conclusions: We report here for the first time the efficient treatment of a metastatic melanoma patient on the basis of circulating tumor DNA analysis. This urgent treatment provided a dramatic response in a patient with a very poor initial condition. The kinetic data most likely reflect treatment efficacy.

Keywords: BRAF, Melanoma, Circulating tumor DNA, Case report

Background

BRAF inhibitors have revolutionized the treatment of metastatic melanoma in patients presenting a BRAF V600 mutation in their tumor by showing highly significant clinical objective responses [1–5]. These drugs have been approved in many countries for the treatment of patients with unresectable or metastatic melanoma with a BRAF mutation. Therefore, daily practice requires BRAF mutation testing of patients' tumors.

Fixed tissues (formalin-fixed and paraffin-embedded) are the standard samples used in routine practice for molecular testing. But sometimes tissues are lacking or

difficult to obtain due to the metastases' location, requiring an invasive and potentially harmful procedure. Lastly, test can fail because of low cellularity or insufficient quality of the DNA. In these cases, circulating tumor DNA (ctDNA) released from tumor cells via mechanisms including necrosis and apoptosis [6–8] can be used as an alternative source of tumor DNA for noninvasive identification of biomarkers. In a recent report Tsao et al. concluded that BRAF mutant ctDNA could be used diagnostically where the tumour block was unavailable [9], but this has never been reported.

Case presentation

A 63-year-old Caucasian woman was referred to our unit with a history of rapidly increasing multiple metastases (duodenal, gastric and colon tumors, peritoneal and retroperitoneal carcinomatosis, many lung, bone and

*Correspondence: marc.denis@chu-nantes.fr

⁴ Department of Biochemistry, Nantes University Hospital, 9 quai Moncoussu, 44093 Nantes Cedex 01, France

Full list of author information is available at the end of the article

gallbladder metastases, mesenteric lymphadenopathies and one brain tumor) (Fig. 1). Biopsies of the colon and stomach concluded the existence of melanoma metastases. The patient had an excision of a primary melanoma of the thigh 15 years earlier but the BRAF status had not been determined. The patient was severely disabled with a very poor performance status (ECOG performance status of 4 and Karnofsky score of 20), ascites and anorexia. Lactate dehydrogenase was equal to 2 times the upper limit.

Because of the seriousness of the metastatic disease and its dramatically rapid progression, supportive care was initially discussed, but we finally decided to test the ctDNA for the presence of a BRAF mutation. Blood was collected, and plasma tested using a ctBRAF mutation test cartridge (Biocartis, Mechelen, Belgium) on an Idylla platform. This very rapid system of ctDNA analysis revealed the presence of the p.V600E mutation in less than 2 h (confirmed 2 weeks later by a test performed on the gastric metastasis using standard techniques [10]). The high concentration of BRAF V600E DNA copies (540 copies/mL plasma) is most likely related to the huge tumor burden [11].

Based on the ctDNA result, a treatment combining a BRAF inhibitor (vemurafenib) and a MEK inhibitor (cobimetinib) was immediately started. After 4 days of treatment, a clinical improvement was noted with a decrease in pain, a progressive recovery of daily living abilities and ascites regression.

In parallel, circulating cell-free DNA analysis was repeated to assess the kinetics of its evolution under treatment. DNA was extracted from plasma samples using the QIAamp circulating nucleic acid kit (Qiagen, Courtaboeuf, France), and analyzed by digital PCR using the QuantStudio 3D System and specific probes (Thermo Fischer, Courtaboeuf, France). The level of mutated BRAF DNA increased as early as 12 h after treatment initiation and reached a maximum after 3 days, followed by a significant decrease after day 4 (Fig. 2). These alterations were barely detectable after 4 weeks of treatment and no longer detectable after 8 weeks.

Tumor assessment according to Response Evaluation Criteria in Solid Tumors, version 1.1 [12], performed with a computed tomography scanner, showed a dramatic response after 8 weeks (86% reduction of the tumor burden). With a 20-week follow-up period, treatment was still ongoing and the response maintained [almost complete response observed at week 18: all target and non-target lesions became non-measurable and no new lesions were observed (Fig. 1)].

Discussion

We report here for the first time the efficient treatment of a metastatic melanoma patient on the basis of ctDNA analysis. We have recently shown that ctDNA samples may be less prone to heterogeneity and provide a better way of determining overall tumor mutation status than a small biopsy [11]. Another potential advantage of ctDNA





Fig. 1 Representative computed tomographic images demonstrating treatment efficacy. Lesions in lung and under liver are presented at baseline and after 8 and 18 weeks of treatment with a combination of BRAF and MEK inhibitors



is that it allows monitoring response [9]. Several reports have shown that, in melanoma patients responding to treatment, a decrease in ctDNA is seen after 1 month [13, 14]. We defined these kinetics more precisely, showing a very rapid increase in mutated ctDNA after treatment initiation, which probably indicates tumor lysis. It was followed by a rapid decrease, and this is most likely associated with response to treatment, as recently demonstrated for non-small cell lung carcinoma (NSCLC) treated with EGFR tyrosine kinase inhibitors [15, 16]. A similar kinetic has been previously reported [9]. Further studies are required to determine whether analysis of a plasma sample as early as 3 or 7 days after treatment initiation could be used to determine whether the patient will respond or not.

Another benefit of a “liquid biopsy” is that turnaround time for ctDNA analysis is expected to be shorter than for tissue genotyping. A prospective analysis of EGFR testing in the ctDNA of patients with newly diagnosed NSCLC recently demonstrated that the median turnaround time for plasma testing was 3 days versus 12 days for tissue genotyping [17].

Conclusions

In 2016, the European Medicines Agency and the US Food and Drug Administration approved the use of ctDNA for EGFR testing, authorizing treatment of

patients with metastatic NSCLC with osimertinib and erlotinib, respectively. Similarly, BRAF ctDNA testing of melanoma patients might also be used as pre-screening before attempting to obtain a result based on tissue samples. If the test is negative, tissue testing should be performed since BRAF mutations are not always detected in plasma of late-stage patients with BRAF mutation-positive tumors [11, 14, 18]. But if the ctDNA analysis is positive, then treatment could be started earlier. In our case, this urgent treatment provided a dramatic response in a patient with a very poor initial condition due to a huge and rapidly growing tumor burden.

Abbreviations

ctDNA: circulating tumor DNA; EGFR: epidermal growth factor receptor; NSCLC: non-small cell lung cancer.

Authors’ contributions

GQ carried out the clinical management of the patient, participated in the design of the study and in the writing of the manuscript. GH, AV, ST and ACK carried out the molecular studies and revised the manuscript. AK, MSJ and BD participated in the design of the study and the writing of the manuscript. MGD performed literature review and wrote the manuscript. All authors read and approved the final manuscript.

Author details

¹ Department of Dermatology, INSERM CIC1413, Nantes University Hospital, Nantes, France. ² CRCINA INSERM U1232, Nantes, France. ³ Immuno-Dermatology Laboratory, Nantes University Hospital, Nantes, France. ⁴ Department of Biochemistry, Nantes University Hospital, 9 quai Moncoussu, 44093 Nantes Cedex 01, France.

Acknowledgements

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article.

Consent for publication

Written informed consent was obtained from the patient for publication of this case report and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Ethics approval and consent to participate

Written informed consent was obtained from the patient and the study was approved by the Ethics Committee of Pays de La Loire (France).

Funding

This study was not funded by any outside source.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 24 October 2016 Accepted: 21 July 2017

Published online: 25 July 2017

References

1. Chapman PB, Hauschild A, Robert C, Haanen JB, Ascierto P, Larkin J, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med*. 2011;364:2507–16.
2. Flaherty KT, Puzanov I, Kim KB, Ribas A, McArthur GA, Sosman JA, et al. Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med*. 2010;363:809–19.
3. Robert C, Karaszewska B, Schachter J, Rutkowski P, Mackiewicz A, Stroiakovski D, et al. Improved overall survival in melanoma with combined dabrafenib and trametinib. *N Engl J Med*. 2015;372:30–9.
4. Long GV, Stroyakovskiy D, Gogas H, Levchenko E, de Braud F, Larkin J, et al. Dabrafenib and trametinib versus dabrafenib and placebo for Val600 BRAF-mutant melanoma: a multicentre, double-blind, phase 3 randomised controlled trial. *Lancet*. 2015;386:444–51.
5. Larkin J, Ascierto PA, Dreno B, Atkinson V, Liskay G, Maio M, et al. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med*. 2014;371:1867–76.

6. Fleischhacker M, Schmidt B. Circulating nucleic acids (CNAs) and cancer—a survey. *Biochim Biophys Acta*. 2007;1775:181–232.
7. Alix-Panabieres C, Schwarzenbach H, Pantel K. Circulating tumor cells and circulating tumor DNA. *Annu Rev Med*. 2012;63:199–215.
8. Schwarzenbach H, Hoon DS, Pantel K. Cell-free nucleic acids as biomarkers in cancer patients. *Nat Rev Cancer*. 2011;11:426–37.
9. Tsao SC, Weiss J, Hudson C, Christophi C, Cebon J, Behren A, Dobrovic A. Monitoring response to therapy in melanoma by quantifying circulating tumour DNA with droplet digital PCR for BRAF and NRAS mutations. *Sci Rep*. 2015;5:11198.
10. Saint-Jean M, Quereux G, Nguyen JM, Peuvrel L, Brocard A, Vallee A, et al. Younger age at the time of first metastasis in BRAF-mutated compared to BRAF wild-type melanoma patients. *Oncol Rep*. 2014;32:808–14.
11. Knol AC, Vallee A, Herbreteau G, Nguyen JM, Varey E, Gaultier A, et al. Clinical significance of BRAF mutation status in circulating tumor DNA of metastatic melanoma patients at baseline. *Exp Dermatol*. 2016;25:783–8.
12. Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer*. 2009;45:228–47.
13. Schreuer M, Meersseman G, Van Den Herrewegen S, Jansen Y, Chevolet I, Bott A, et al. Quantitative assessment of BRAF V600 mutant circulating cell-free tumor DNA as a tool for therapeutic monitoring in metastatic melanoma patients treated with BRAF/MEK inhibitors. *J Transl Med*. 2016;14:95.
14. Sanmamed MF, Fernandez-Landazuri S, Rodriguez C, Zarate R, Lozano MD, Zubiri L, et al. Quantitative cell-free circulating BRAFV600E mutation analysis by use of droplet digital PCR in the follow-up of patients with melanoma being treated with BRAF inhibitors. *Clin Chem*. 2015;61:297–304.
15. Marchetti A, Palma JF, Felicioni L, De Pas TM, Chiari R, Del Grammastrom M, et al. Early prediction of response to tyrosine kinase inhibitors by quantification of EGFR mutations in plasma of NSCLC patients. *J Thorac Oncol*. 2015;10:1437–43.
16. Vallee A, Audigier-Valette C, Herbreteau G, Merrien J, Tessonnier L, Theoleyre S, Denis MG. Rapid clearance of circulating tumor DNA during treatment with AZD9291 of a lung cancer patient presenting the resistance EGFR T790M mutation. *Lung Cancer*. 2016;91:73–4.
17. Sacher AG, Paweletz C, Dahlberg SE, Alden RS, O'Connell A, Feeney N, et al. Prospective validation of rapid plasma genotyping for the detection of EGFR and KRAS mutations in advanced lung cancer. *JAMA Oncol*. 2016;2:1014–22.
18. Santiago-Walker A, Gagnon R, Mazumdar J, Casey M, Long GV, Schadendorf D, et al. Correlation of BRAF mutation status in circulating-free DNA and tumor and association with clinical outcome across four BRAFi and MEKi clinical trials. *Clin Cancer Res*. 2016;22:567–74.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at www.biomedcentral.com/submit

