



**HAL**  
open science

# Draft Genome Sequence of Highly Rifampin-Resistant Propionibacterium namnetense NTS 31307302T Isolated from a Patient with a Bone Infection

Guillaume Ghislain Aubin, Stanimir Kambarev, Pascale Bémer, Paul A  
Lawson, Stéphane Corvec

► **To cite this version:**

Guillaume Ghislain Aubin, Stanimir Kambarev, Pascale Bémer, Paul A Lawson, Stéphane Corvec. Draft Genome Sequence of Highly Rifampin-Resistant Propionibacterium namnetense NTS 31307302T Isolated from a Patient with a Bone Infection. Genome Announcements, 2016, 4 (4), 10.1128/genomeA.00819-16 . inserm-01822375

**HAL Id: inserm-01822375**

**<https://inserm.hal.science/inserm-01822375>**

Submitted on 25 Jun 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Draft Genome Sequence of Highly Rifampin-Resistant *Propionibacterium namnetense* NTS 31307302<sup>T</sup> Isolated from a Patient with a Bone Infection

Guillaume Ghislain Aubin,<sup>a,b</sup> Stanimir Kambarev,<sup>c</sup> Pascale Bémer,<sup>a</sup> Paul A. Lawson,<sup>d</sup>  Stéphane Corvec<sup>a,e</sup>

Bacteriology and Hygiene Department, Nantes University Hospital, Nantes, France<sup>a</sup>; EA3826, Laboratory of Clinical and Experimental Therapeutics of Infections, Nantes Medicine School, Nantes, France<sup>b</sup>; Institut de Recherche en Santé de l'Université de Nantes INSERM U892—CNRS 6299 CRCNA Centre de Recherche en Cancérologie Nantes Angers, Université de Nantes, Team 13: Nuclear Oncology, Nantes, France<sup>c</sup>; Department of Microbiology and Plant Biology, University of Oklahoma, Norman, Oklahoma, USA<sup>d</sup>; Institut de Recherche en Santé de l'Université de Nantes INSERM U892—CNRS 6299 CRCNA Centre de Recherche en Cancérologie Nantes Angers, Université de Nantes, Team 2: Clinical and Translational Research in Skin Cancer, Nantes, France<sup>e</sup>

*Propionibacterium namnetense* was recently described as a potential bone pathogen, which is closely related to *Propionibacterium acnes*, a skin commensal microorganism. Here, we report the draft genome sequence of the highly rifampin-resistant strain NTS 31307302<sup>T</sup> isolated from a patient with a tibia infection.

Received 17 June 2016 Accepted 17 June 2016 Published 11 August 2016

**Citation** Aubin GG, Kambarev S, Bémer P, Lawson PA, Corvec S. 2016. Draft genome sequence of highly rifampin-resistant *Propionibacterium namnetense* NTS 31307302<sup>T</sup> isolated from a patient with a bone infection. *Genome Announc* 4(4):e00819-16. doi:10.1128/genomeA.00819-16.

**Copyright** © 2016 Aubin et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stéphane Corvec, [stephane.corvec@chu-nantes.fr](mailto:stephane.corvec@chu-nantes.fr).

*Propionibacterium acnes* is known as a Gram-positive bacterium constituting a significant part of the human skin microbiota (1). Its natural habitat is mainly the lipid-rich sebaceous glands (2, 3). *Acne vulgaris* is a common chronic skin disease and is usually associated with the detection of this organism. Moreover, *P. acnes* is more frequently identified in device-related infections (4, 5), producing biofilm in this context (6). Recently, this species has been subdivided into several phylogenetic types that were subsequently afforded subspecies status (7).

We present here the genome sequence of *Propionibacterium namnetense* NTS 31307302<sup>T</sup> isolated at Nantes University Hospital, France, during a bone infection (8). The isolate showed beta-hemolysis on a blood agar plate and was recently described as being related to *P. acnes* (9). Strain NTS 31307302<sup>T</sup> is resistant to rifampin, as it has been already reported for *in vitro*-selected mutants (10) or clinical strains of *P. acnes* involved in biofilm or device-related infections (11).

*P. namnetense* NTS 31307302<sup>T</sup> was grown overnight at 37°C on a Schaedler agar plate (Oxoid, United Kingdom) under an anaerobic atmosphere. Genomic DNA was extracted using a DNeasy blood and tissue kit (Qiagen GmbH, Germany), according to the provider's recommendation. A paired-end library was prepared with the NEBNext Ultra DNA library prep kit for Illumina (NEB) and sequenced (2 × 150 bp) on a MiSeq Sequencer (Illumina, USA). *De novo* assembly was performed with Velvet 1.2.10 and VelvetOptimiser 2.2.5 (optimal hash value, 127). A total of 2,846,458 reads were assembled into 24 contigs (15 of them >1 kb) with an average coverage of 127×. Contig reordering and annotation were performed with Mauve 2.3.1 (12) and the NCBI Prokaryotic Genome Automatic Annotation Pipeline (PGAAP) (13), respectively. Average nucleotide identities (ANI) were calculated using Oat 0.91 (14).

The final assembly has a total length of 2,369,664 bp, an  $N_{50}$  of

626 kb, and a G+C content of 60.5%. About 2,136 coding sequences (CDSs), 46 tRNAs, 84 pseudogenes, three rRNAs, and three noncoding RNAs were revealed by annotation.

To determine genomic differences between *P. namnetense* NTS 31307302<sup>T</sup> and the closely related *P. acnes* KPA171202, ATCC 11828, and ATCC 6919<sup>T</sup>, we performed a genomic comparison. The draft genome size of the newly sequenced strain is 2.37 Mb, which is 5.02 to 9.81% smaller than *P. acnes* reference strains. The ANI value was 88.5%. Interestingly, our strain showed an ANI value of 99.52% with *P. acnes* SK182B-JCVI (accession no. AFUN00000000.1), recovered during the Human Microbiome Project, which is significantly above the cutoff value of 95% for species delineation (14). Therefore, this strain is likely to be another isolate of this species. Comparing both sequences of the *rpoB* gene, we observed only one point mutation at nucleotide 1319 (G→A) in strain NTS 31307302<sup>T</sup> leading to an amino acid modification at position 440 (R440H), previously described to be involved in rifampin resistance in *P. acnes* (11).

This draft genome of *P. namnetense* NTS 31307302<sup>T</sup> will be used for studying virulence features associated with bone infection, especially hemolysin, lipase, or hyaluronidase (15).

**Accession number(s).** The draft sequence of *P. namnetense* NTS 31307302<sup>T</sup> studied in this project has been deposited at DDBJ/EMBL/GenBank under the accession no. [LWHO01000001](https://www.ncbi.nlm.nih.gov/nuclink/LWHO01000001). The version described in this paper is [LWHO01000001.1](https://www.ncbi.nlm.nih.gov/nuclink/LWHO01000001.1).

## ACKNOWLEDGMENTS

This work was supported by an internal grant.

We are grateful to the staff for the IRS-1 DNA sequencing platform for their advice and technical help.

## FUNDING INFORMATION

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## REFERENCES

- Findley K, Grice EA. 2014. The skin microbiome: a focus on pathogens and their association with skin disease. *PLoS Pathog* 10:e1004436. <http://dx.doi.org/10.1371/journal.ppat.1004436>.
- Aubin GG, Portillo ME, Trampuz A, Corvec S. 2014. *Propionibacterium acnes*, an emerging pathogen: from acne to implant-infections, from phylogeny to resistance. *Méd Mal Infect* 44:241–250. <http://dx.doi.org/10.1016/j.medmal.2014.02.004>.
- Leccia MT, Auffret N, Poli F, Claudel J-P, Corvec S, Dreno B. 2015. Topical acne treatments in Europe and the issue of antimicrobial resistance. *J Eur Acad Dermatol Venereol* 29:1485–1492. <http://dx.doi.org/10.1111/jdv.12989>.
- Portillo ME, Corvec S, Borens O, Trampuz A. 2013. *Propionibacterium acnes*: an underestimated pathogen in implant-associated infections. *BioMed Res Int* 2013:804391. <http://dx.doi.org/10.1155/2013/804391>.
- Bémer P, Corvec S, Tariel S, Asseray N, Boutoille D, Langlois C, Tequi B, Drugeon H, Passuti N, Touchais S. 2008. Significance of *Propionibacterium acnes*-positive samples in spinal instrumentation. *Spine* 33:E971–E976. <http://dx.doi.org/10.1097/BRS.0b013e31818e28dc>.
- Furustrand Tabin U, Corvec S, Betrisey B, Zimmerli W, Trampuz A. 2012. Role of rifampin against *Propionibacterium acnes* biofilm *in vitro* and in an experimental foreign-body infection model. *Antimicrob Agents Chemother* 56:1885–1891. <http://dx.doi.org/10.1128/AAC.05552-11>.
- Dekio I, Culak R, Misra R, Gaulton T, Fang M, Sakamoto M, Ohkuma M, Oshima K, Hattori M, Klenk H-P, Rajendram D, Gharbia SE, Shah HN. 2015. Dissecting the taxonomic heterogeneity within *Propionibacterium acnes*: proposal for *Propionibacterium acnes* subsp. *acnes* subsp. nov. and *Propionibacterium acnes* subsp. *elongatum* subsp. nov. *Int J Syst Evol Microbiol* 65:4776–4787. <http://dx.doi.org/10.1099/ijsem.0.000648>.
- Aubin GG, Bémer P, Kambarev S, Patel NB, Lemenand O, Caillon J, Lawson PA, Corvec S. 3 June 2016. *Propionibacterium namnetense* sp. nov., isolated from a human bone infection. *Int J Syst Evol Microbiol* [Epub ahead of print.] <http://dx.doi.org/10.1099/ijsem.0.001204>.
- Corvec S, Luchetta J, Aubin GG. 2015. Letter to the editor: is hemolysis a clinical marker of *Propionibacterium acnes* orthopedic infection or a phylogenetic marker? *Am J Orthop Belle Mead NJ* 44:E61–E62.
- Furustrand Tabin U, Trampuz A, Corvec S. 2013. *In vitro* emergence of rifampicin resistance in *Propionibacterium acnes* and molecular characterization of mutations in the *rpoB* gene. *J Antimicrob Chemother* 68:523–528. <http://dx.doi.org/10.1093/jac/dks428>.
- Furustrand Tabin U, Aubin GG, Eich G, Trampuz A, Corvec S. 2015. Occurrence and new mutations involved in rifampicin-resistant *Propionibacterium acnes* strains isolated from biofilm or device-related infections. *Anaerobe* 34:116–119. <http://dx.doi.org/10.1016/j.anaerobe.2015.05.003>.
- Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* 5:e11147. <http://dx.doi.org/10.1371/journal.pone.0011147>.
- Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity G, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of Standard Operating Procedures (SOPs) for (meta)genomic annotation. *Omic J Integr Biol* 12:137–141. <http://dx.doi.org/10.1089/omi.2008.0017>.
- Lee I, Kim YO, Park S-C, Chun J. 2016. OrthoANI: an improved algorithm and software for calculating average nucleotide identity. *Int J Syst Evol Microbiol* 66:1100–1103. <http://dx.doi.org/10.1099/ijsem.0.000760>.
- Scholz CF, Brüggemann H, Lomholt HB, Tettelin H, Kilian M. 2016. Genome stability of *Propionibacterium acnes*: a comprehensive study of indels and homopolymeric tracts. *Sci Rep* 6:20662. <http://dx.doi.org/10.1038/srep20662>.