

HIGHLIGHTS

- Peptide HLA Class I presentation by β cells is increased by inflammatory cytokines.
- Peptide sources feature several insulin granule proteins, e.g. SCG5, PCSK2, UCN3.
- SCG5-009 mRNA splice products and IAPP fusion peptides are also presented.
- In type 1 diabetes, peptide-reactive CD8⁺ T cells are enriched in the pancreas.

eTOC BLURB

The epitopes presented by β cells for recognition by autoimmune CD8⁺ T lymphocytes remain elusive. Gonzalez-Duque et al. identify them as derived from conventional processing, mRNA and peptide splicing. They are selectively recognized by CD8⁺ T lymphocytes of type 1 diabetic patients in the pancreas, but not in the blood.

Manuscript

Conventional and neo-antigenic peptides presented by β cells are preferentially targeted in the pancreas, but not in blood, of type 1 diabetic patients

Sergio Gonzalez-Duque¹⁻⁴, Marie Eliane Azoury¹⁻³, Maikel L. Colli⁵, Georgia Afonso¹⁻³, Jean-

Valery Turatsinze⁵, Laura Nigi⁶, Ana Ines Lalanne¹⁻³, Guido Sebastiani⁶, Alexia Carré¹⁻³,

Sheena Pinto⁷, Slobodan Culina¹⁻³, Noémie Corcos¹⁻³, Marco Bugliani⁸, Piero Marchetti⁸,

Mathieu Armanet9, Marc Diedisheim1-3,10, Bruno Kyewski7, Lars M. Steinmetz11,12, Søren

Buus¹³, Sylvaine You¹⁻³, Daniele Dubois-Laforgue^{1-3,10}, Etienne Larger^{1-3,10}, Jean-Paul

Beressi¹⁴, Graziella Bruno¹⁵, Francesco Dotta⁶, Raphael Scharfmann¹⁻³, Decio L. Eizirik^{5*},

Yann Verdier^{4*}, Joelle Vinh^{4*}, Roberto Mallone^{1-3,10**}.

¹INSERM, U1016, Cochin Institute, Paris, F-75014, France.

²CNRS, UMR8104, Cochin Institute, Paris, F-75014, France.

³Paris Descartes University, Sorbonne Paris Cité, Paris, F-75014, France.

⁴ESPCI Paris, PSL University, Spectrométrie de Masse Biologique et Protéomique, CNRS USR3149, Paris, F-75005, France.

⁵Université Libre de Bruxelles Center for Diabetes Research and Welbio, Medical Faculty, Université Libre de Bruxelles, Brussels, B-1070, Belgium.

⁶University of Siena, Department of Medicine, Surgery and Neuroscience, Diabetes Unit and Fondazione Umberto di Mario ONLUS, Toscana Life Sciences, Siena, I-53100, Italy.

⁷DKFZ, Division of Developmental Immunology, Heidelberg, D-69120, Germany.

⁸University of Pisa, Department of Clinical and Experimental Medicine, Pisa, I-56124, Italy. ⁹Assistance Publique Hôpitaux de Paris, Cell Therapy Unit, Saint Louis Hospital, Paris, F-75010, France.

¹⁰Assistance Publique Hôpitaux de Paris, Service de Diabétologie, Cochin Hospital, Paris, F-75014, France.

¹¹Stanford University, School of Medicine, Department of Genetics and Stanford Genome Technology Center, Stanford, 94305, USA.

¹²European Molecular Biology Laboratory, Genome Biology Unit, Heidelberg, D-69117 Germany.

¹³Panum Institute, Department of International Health, Immunology and Microbiology, Copenhagen, DK-2200, Denmark.

¹⁴Centre Hospitalier de Versailles André Mignot, Service de Diabétologie, Le Chesnay, F-78150, France.

¹⁵University of Turin, Department of Medical Sciences, Turin, I-10126, Italy.

*These authors contributed equally to this work

**Correspondence and Lead Contact: roberto.mallone@inserm.fr

SUMMARY

Although CD8⁺ T-cell-mediated autoimmune β -cell destruction occurs in type 1 diabetes (T1D), the target epitopes processed and presented by β cells are unknown. To identify them, we combined peptidomics and transcriptomics strategies. Inflammatory cytokines increased peptide presentation *in vitro*, paralleling upregulation of Human Leukocyte Antigen (HLA) Class I expression. Peptide sources featured several insulin granule proteins and all known β -cell antigens, barring islet-specific glucose-6-phosphatase catalytic subunit-related protein. Preproinsulin yielded HLA-A2-restricted epitopes previously described. Secretogranin V and its mRNA splice isoform SCG5-009, proconvertase-2, urocortin-3, the insulin gene enhancer protein ISL-1 and an islet amyloid polypeptide transpeptidation product emerged as antigens processed into HLA-A2-restricted epitopes which, as those already described, were recognized by circulating naïve CD8⁺ T cells in T1D and healthy donors, and by pancreas-infiltrating cells in T1D donors. This peptidome opens new avenues to understand antigen processing by β cells and for developing T-cell biomarkers and tolerogenic vaccination strategies.

Keywords: antigen, autoimmunity, β cell, epitope, Human Leukocyte Antigen (HLA), preproinsulin, T cell, type 1 diabetes.

INTRODUCTION

Autoimmune CD8⁺ T cells dominate the pancreatic immune infiltrates of human type 1 diabetes (T1D) (Coppieters et al., 2012), and lyse β cells by recognizing surface peptide-Human Leukocyte Antigen Class I (pHLA-I) complexes. Identifying these peptides is therefore critical for developing tolerogenic vaccination strategies and immune staging tools targeting islet-reactive CD8⁺ T cells.

Most islet antigens (Ags), namely insulin (INS) and its precursor preproinsulin (PPI), glutamic acid decarboxylase (GAD65/GAD2), islet Ag (IA)-2 (PTPRN) (Mallone et al., 2007; Martinuzzi et al., 2008), and zinc transporter 8 (ZnT8/SLC30A8) (Scotto et al., 2012), have been identified based on their targeting by auto-antibodies (aAbs). Other Ags such as islet-specific glucose-6-phosphatase catalytic subunit-related protein (IGRP) (Mallone et al., 2007), chromogranin A (CHGA) (Li et al., 2015) and islet amyloid polypeptide (IAPP) (Standifer et al., 2006) have been identified based on studies in the non-obese diabetic mouse and/or their islet-enriched expression. A systematic discovery effort is missing, and the available catalogue may be biased by the lack of information about the peptides that are naturally processed and presented by β cells.

Mutated sequences in tumor proteins become preferential CD8⁺ T-cell targets (Gubin et al., 2014), possibly because they are regarded as non-self and therefore not efficiently tolerized. Similarly, other processes in β cells may facilitate tolerance escape: post-translational modifications (PTMs) (McGinty et al., 2014; Rondas et al., 2015), transpeptidation, i.e. the splicing and fusion of non-contiguous peptide fragments from the same protein or from different ones (Babon et al., 2016; Delong et al., 2016), and the use of alternative transcription start sites (Kracht et al., 2017). These studies have mostly focused on CD4⁺ T cells, which are stimulated by pHLA Class II complexes presented by professional Ag-presenting cells that uptake β -cell material. These indirect Ag processing pathways do not reflect those that are

specific to β cells. Indeed, several arguments suggest an active role of β cells in their own demise (Eizirik et al., 2009).

First, we recently showed that some T1D susceptibility gene variants modulate islet inflammation (Marroqui et al., 2015; Marroqui et al., 2014; Moore et al., 2009), suggesting that the β -cell response to inflammation is genetically modulated. This response triggers cytokine/chemokine release, endoplasmic reticulum (ER) stress and HLA-I upregulation (Eizirik et al., 2009; Marroqui et al., 2017), which facilitate a productive autoimmune response. The alternative mRNA splicing signature induced by β -cell inflammation (Eizirik et al., 2012; Ortis et al., 2010) has received less attention, but may similarly generate neosequences not translated in the thymus and regarded as non-self.

Second, we recently reported a circulating islet-reactive CD8⁺ T-cell repertoire that is predominantly naïve and largely overlapping between T1D and healthy subjects (Culina et al., 2018). These findings reveal a general leakiness of central tolerance irrespective of T1D status, begging the question of what determines T1D progression versus the maintenance of a 'benign' state of autoimmunity. One hypothesis is that the target β cell and its response to inflammation may be critical in the progression toward T1D in the face of similar autoimmune T-cell repertoires across individuals.

In this context, it is crucial to understand the 'image' that human β cells deliver to CD8⁺ T cells through pHLA-I complexes. To this end, we implemented a strategy combining HLA-I peptidomics on β cells and RNAseq analysis of the splice isoforms transcribed by primary islets exposed or not to inflammatory cytokines and by thymic medullary epithelial cells (mTECs), focusing primarily on the most prevalent HLA-A2 variant.

4

RESULTS

The HLA-I peptidome of human β cells is enriched by cytokine exposure and displays the expected amino-acid length and motifs.

Our first epitope discovery pipeline employed HLA-I peptidomics experiments on the ECN90 β-cell line (Culina et al., 2018), which carries the HLA-I haplotype A*02:01/A*03:01/B*40:01/B*49:01/C*03:04/C*07:01 (A2/A3/B40/B49/C3/C7 from here on). ECN90 β cells were cultured overnight with or without interferon (IFN)- γ , alone or in combination with tumor necrosis factor (TNF)- α and interleukin (IL)-1 β , and lysed to immunopurify pHLA-I complexes. HLA-I-bound peptides were then analyzed by liquid chromatography-tandem mass spectrometry (LC-MS/MS). Although ECN90 β cells expressed surface HLA-I under basal conditions, this expression was significantly upregulated upon cytokine treatment (Fig. 1A-B), without significant cell death (Culina et al., 2018). The 2,997 eluted peptides were mostly (93%) 8-12-mers (Fig. 1C), i.e. the length required for HLA-I binding. The amino acid (aa) identity at pHLA-I anchor positions (p2 and p9) also revealed the preferences expected based on the aforementioned HLA-I haplotype (Fig. 1D-E). In line with the observed HLA-I upregulation, the number of eluted peptides was significantly higher in the presence of cytokines, and higher in β cells exposed to IFN- γ , TNF- α and IL-1 β compared with IFN-y alone (Fig. 1F).

These peptide datasets were subsequently analyzed using a stepwise bioinformatics pipeline (Fig. 1G). First, only peptides that were reproducibly detected in at least 2 of 5 biological replicates (85%; all percentages are given in relation to the number of peptides retained by the previous filter) and that displayed the expected 8-12-aa length (93%) were selected. β -cell-enriched peptides (both conventional and with PTMs, excluding those derived from peptide or mRNA splicing; red pipeline in Fig. 1G) were subsequently filtered based on non-ubiquitous (16%) and enriched β -cell expression (34%) of their source proteins. For other non-

conventional peptides (i.e. PTM or transcriptional variants), no expression filter was applied, as these species could be β -cell-specific in spite of a ubiquitous expression of the source protein or mRNA. PTM peptides (methionine, tryptophan, histidine and cysteine oxidation tryptophan conversion to kynurenine) derived from ubiquitous proteins accounted for 8% of the whole dataset (blue pipeline in Fig. 1G). Compounds potentially corresponding to peptide splice variants (0.5%; brown pipeline) were identified using an in-house script (Fig. S1 and Data S1) based on reported peptide splicing preferences (Berkers et al., 2015) applied to known and putative Ags.

For peptides derived from mRNA splice variants (dotted and green pipeline in Fig. 1G), the peptidomics dataset was interrogated against RNAseq datasets from primary human islets exposed or not to cytokines and from human mTECs (<u>Data S2</u>). First, higher gene expression can favor peptide processing and presentation. Hence, mRNA splice variants were selected based on a median Reads Per Kilobase per Million mapped reads (RPKM)>5 in islets (27%), based on the median RPKM of known islet Ags (Eizirik et al., 2012). Second, mRNA isoforms poorly expressed in mTECs might favor T-cell escape from clonal deletion. Thus, mRNA variants with a RPKM<0.1 in mTECs or with a fold-decrease>100 *vs.* islets were selected (6%). Third, we selected mRNA isoforms with >10-fold higher expression in islets *vs.* other tissues. We then analyzed the predicted aa neo-sequences encoded by these mRNA variants, yielding 88/166 mRNA variants (53%) and 336 peptide neo-sequences that were used to interrogate the HLA-I peptidomics dataset, with 2 hits found. Finally, each dataset was filtered to retain only those peptides found enriched in HLA-I- *vs.* mock-purified samples, leading to the overall exclusion of 48% hits. We focused on predicted HLA-A2-restricted peptides for subsequent HLA-A2 binding and CD8⁺ T-cell recognition studies.

Collectively, these results show that inflammatory cytokines increase pHLA-I presentation and that these peptides display the aa signatures required for HLA-I binding.

pHLA-I complexes from human β cells are enriched in peptides derived from secretory granule proteins, including known PPI epitopes.

The filtered HLA-I peptidomics dataset obtained is described in <u>Fig. 2</u> and detailed in <u>Table S1</u>. While 42/98 (43%) eluted peptides were shared among basal and cytokine-treated conditions, 45/98 (46%) peptides were only detected upon cytokine exposure (<u>Fig. 2A</u>). Also quantitatively, most peptides (62/98; 63%) were exclusively or more presented in cytokine-treated ECN90 β cells (Table S1).

Among the 40 source proteins of HLA-I-eluted peptides (Fig. 2B), the most represented ones were the known islet Ags CHGA (n=15 peptides) and PPI (n=12, plus one derived from an INS-006 mRNA splice variants). Besides the other known Ag IA-2 (PTPRN; n=3), the 5 top scoring proteins included two putative Ags: Kinesin Family Member 1A (KIF1A; n=9) and Secretogranin V (SCG5/7B2; n=3, plus one derived from a SCG5-009 mRNA splice variant). Other proteins included known Ags, i.e. GAD2 (GAD65), SLC30A8 (ZnT8) and IAPP (splice peptide), and several putative ones. Notably, all the HLA-A2-restricted PPI peptides identified, namely PPI₂₋₁₀, PPI₆₋₁₄, PPI₁₅₋₂₄, PPI₂₉₋₃₈ (INS_{B5-14}) and PPI₃₄₋₄₂ (INS_{B10-18}) (Fig. <u>2C</u>), are already described as major $CD8^+$ T-cell epitopes, thus validating our discovery strategy. Source proteins were enriched for insulin granule products (14/42, 33%; Fig. 2D), namely CHGA, INS, SCG5, PTPRN, ATP-binding cassette sub-family C member 8 (ABCC8), proprotein convertase 1 (PCSK1/PC1), urocortin-3 (UCN3), chromogranin B (CHGB), carboxypeptidase E (CPE), proprotein convertase 2 (PCSK2/PC2), secretogranin III (SCG3), SLC30A8; and IAPP and neuropeptide Y (NPY) generating splice peptides. The predicted HLA-I restrictions of these peptides (Fig. 2E) comprised all the alleles expressed by ECN90 β cells, while 10% of restrictions could not be assigned. For peptides derived from β cell-enriched proteins, 11/98 (11%) carried PTMs, with most of them (8/11; 73%) representing variants of unmodified peptides identified in this same dataset. Most of these modifications (7/11; 64%) were M(+15.99) methionine, C(+47.98) cysteine and W(+15.99) tryptophan oxidations, with W(+3.99) tryptophan to kynurenine transitions also detected. The 99 PTM peptides derived from ubiquitous source proteins are listed in <u>Table S2</u>.

To validate the results obtained using the ECN90 β -cell line, a HLA-A2⁺ primary human islet preparation that did not share other HLA-I alleles with ECN90 cells was analyzed similarly. The major source proteins of the HLA-I-bound peptides identified were largely overlapping with those found in ECN90 cells (Fig. 2F), with INS (n=12 peptides), CHGA (n=4), KIF1A (n=3) and SCG5 (n=3) ranking highest for both cells, and CHGB (n=3), PCSK2 (n=1) and an identical IAPP splice peptide also detected in both. When analyzing the identity of individual peptides (including length and PTM variants) (Table S3), 16/33 (48%) were shared between ECN90 and primary islet cells. This common repertoire increased to 12/13 peptides (92%) when considering only predicted HLA-A2 binders, supporting the validity of the ECN90 βcell model. Of note, shared peptides included all the PPI species already described as CD8⁺ Tcell epitopes, SCG5186-196 along with a shorter SCG5186-195 variant with higher HLA-A2 affinity, and the IAPP₁₅₋₁₇/IAPP₅₋₁₀ splice peptide VAL/KLQVFL. Although this product could also reflect PTPRN596-598/IAPP5-10 trans-splicing, IAPP15-17/IAPP5-10 is more likely because the intra-protein vicinity of these fragments is more favorable for transpeptidation. The new hits identified were mostly predicted to bind to the HLA-I molecules not shared with ECN90 cells, barring an HLA-A2-restricted CHGB440-448 peptide retained for further validation. Contrary to ECN90 cells, most peptides were detected at similar levels under basal and cytokine-treated conditions (Table S3). This mirrored a higher basal HLA-I expression in primary islets, possibly reflecting peri-mortem and tissue isolation stress conditions, which was less upregulated by cytokine treatment (Fig. 1G). Moreover, detection sensitivity may have been limited by the concomitant isolation of non- β -cell peptides, i.e. pancreatic polypeptide- and glucagon-derived sequences likely eluted from δ and α cells (n=4 and 5, respectively; not shown because excluded by the β -cell enrichment filter).

The fragmentation profile of the identified peptides was confirmed by comparing their MS/MS spectra with those of the corresponding synthetic peptides. Finally, the predicted HLA-A2 binding was experimentally verified (Fig. S2A-H), leading to the final selection of 18/19 (95%) HLA-I-eluted peptides for CD8⁺ T-cell studies (including CHGB₄₄₀₋₄₄₈ eluted from primary islets).

Collectively, these data show that β cells process and present several known HLA-A2restricted PPI epitopes and additional candidate ones, which are enriched for secretory granule products.

In silico analysis of mRNA splice variants yields additional predicted neo-Ag peptides.

The RNAseq dataset used for assigning m/z species was further mined *in silico*, independently of the HLA-I peptidomics pipeline, to identify other potential HLA-A2-restricted peptides (Fig. 1G, dotted pipeline). Selection was based on a predicted HLA-A2 binding, a 9-10-aa length and a neo-sequence stretch \geq 3 aa, with 39 candidates retained (Fig. S2I). These were splice variants of known β -cell Ags (GAD2-003, IAPP-002, IAPP-004, PTPRN-021, SLC30A8-002) and of candidate ones. Most of the source mRNA splice variants (35/39, 90%) were similarly expressed in untreated and cytokine-treated islets. HLA-A2 binding was experimentally confirmed for 34/39 candidates (87%; Fig. S2I), which were retained for further validation along with the 18 HLA-A2 binders identified by HLA-I peptidomics.

HLA-A2-restricted β -cell peptides are targeted by a circulating naïve CD8⁺ T-cell repertoire in healthy donors.

We previously documented that the vast majority of individuals, both type 1 diabetic and healthy, harbor similar frequencies of circulating, predominantly naïve HLA-A2-restricted CD8⁺ T cells reactive to known PPI, GAD65, IA-2, IGRP and ZnT8 epitopes (Culina et al., 2018). Similarly, we reasoned that the presence of a cognate naïve CD8⁺ T-cell repertoire is the preliminary requirement for the immunogenicity of the HLA-A2-restricted candidate epitopes identified in the in vitro HLA-I peptidomics and in silico transcriptomics pipeline (n=52; 18 and 34, respectively). We therefore started by screening these candidates for recognition by circulating CD8⁺ T cells in HLA-A2⁺ healthy donors (Table S4), using combinatorial double-coded HLA-A2 multimers (MMrs) loaded with the corresponding synthetic peptides (Culina et al., 2018). We retained those candidates that harbored a cognate naïve $CD8^+$ T-cell repertoire, based on *i*) the frequency of this repertoire, which is typically in the range of $1-50/10^6$ CD8⁺ T cells (Alanio et al., 2010; Culina et al., 2018); and *ii*) the pattern of HLA-A2 MMr staining, which is usually clustered rather than spread in the presence of a specific epitope-reactive population (James et al., 2018). The gating strategy is presented in Fig. 3 and representative dot plots in Fig. 4A-F. Using these two criteria, several candidate epitopes displayed a cognate naïve CD8⁺ T-cell repertoire in the expected range in a sizable fraction (\geq 50%) of the healthy individuals analyzed. The frequency of CD8⁺ T cells recognizing the known β -cell epitope PPI₆₋₁₄ previously analyzed (Culina et al., 2018) also fell in the same range, with some outliers noted. In total, 9/18 HLA-I-eluted peptides (50%; Fig. 4G) were validated, namely CHGA₃₄₄₋₃₅₂, insulin gene enhancer protein ISL1₂₇₆₋₂₈₄, potassium channel subfamily K member 16 (KCNK16)₁₂₉₋₁₃₇, KIF1A₁₃₄₇₋₁₃₅₅, PCSK2₃₀₋₃₈, SCG5₁₈₆₋₁₉₅, SCG5-009₁₈₆₋₁₉₄ and UCN3₁₋₉. Despite recognition in only 1 of 6 donors analyzed, the peptide splice product IAPP₁₅₋₁₇/IAPP₅₋₁₀ was also retained, since it was identified in both ECN90 and primary islet cells. Using the same criteria, 11/34 candidates selected in silico were validated (32%; Fig. 4H), namely cyclin I (CCNI)-00814-22, GAD2003₁₇₉₋₁₈₇, guanine nucleotide-binding protein G(s) subunit α isoforms short (GNAS)-036₆₇₋₇₅, GNAS-036₁₂₄₋₁₃₂, IAPP-002₃₃₋₄₂, PTPRN-021₃₉₂₋₄₀₂, PTPRN-021₃₉₈₋₄₀₇, phogrin/receptor-type tyrosine-protein phosphatase N2 (PTPRN2)-005₁₁₋₁₉, PTPRN2-005₁₉₋₂₇, mitochondrial oligoribonuclease (REXO2)-020₂₋₁₀, and SLC30A8-002₁₆₋₂₅. As previously observed for other known β-cell epitopes (Culina et al., 2018), including the PPI₆₋₁₄ here used as β-cell positive control, only a minority (median 16.4%, interquartile range 8.5-26.7%) of CD8⁺ T cells recognizing these candidate epitopes were Ag-experienced (CD45RA⁺CCR7⁻, CD45RA⁻ CCR7⁻ or CD45RA⁻CCR7⁺; Fig. 4I-J). Conversely, the Flu MP₅₈₋₆₆ peptide included as viral positive control displayed the expected predominantly Ag-experienced phenotype. The complete list of the 20 candidates validated for CD8⁺ T-cell recognition is presented in <u>Table</u> <u>1</u>. All the peptides validated came from source proteins whose gene expression was detected in islets, both under basal and cytokine-treated conditions. One notable exception was SCG5-009, whose expression was negligible under basal condition but strongly upregulated after cytokine treatment. Gene expression in mTECs was also negligible in all cases, with the exception of CHGA, ISL1 and SCG5.

Collectively, these results show that most of the β -cell peptides identified display a cognate naïve CD8⁺ T-cell repertoire in the blood of healthy individuals, making them potential targets of islet autoimmunity.

Circulating CD8⁺ T cells reactive to HLA-A2-restricted β -cell peptides display similar *ex-vivo* frequencies and a predominantly naïve phenotype in T1D and healthy subjects.

Thirteen of the 20 β -cell peptides validated for recognition by a naïve CD8⁺ T-cell repertoire were selected for further *ex-vivo* combinatorial double-coded MMr validation using blood samples from HLA-A2⁺ recent-onset T1D and healthy subjects (n=10/each; Table S4). For naturally processed and presented peptides identified by HLA-I peptidomics, we focused on 6 insulin granule putative Ags, namely IAPP₁₅₋₁₇₅₋₁₀, PCSK2₃₀₋₃₈, SCG5₁₈₆₋₁₉₅, SCG5-009₁₈₆₋₁₉₄ and UCN3₁₋₉; and the transcription factor ISL1₂₇₆₋₂₈₄. A less focused selection was made for 7 predicted mRNA splice peptides, as these may be derived from short-lived, unstable defective ribosomal products (DRiPs) (Anton and Yewdell, 2014). CCNI-008₁₄₋₂₂, GAD2-003₁₇₉₋₁₈₇, GNAS-036₆₇₋₇₅, GNAS-036₁₂₄₋₁₃₂, IAPP-002₃₃₋₄₂, PTPRN2-005₁₁₋₁₉ and SLC30A8-002₁₆₋₂₅ were thus selected. The frequency of circulating CD8⁺ T cells recognizing these peptides and the control PPI₆₋₁₄ epitope was similar in T1D and healthy subjects (Fig. 5A), and fell in the same range (1-50/10⁶ CD8⁺ T cells) detected in the preliminary screening performed on healthy subjects using different fluorochrome-labeled MMr combinations (Fig. 4G-H), with the exception of IAPP-002₃₃₋₄₂ for which virtually no MMr⁺ cells were detected. As in the screening round, frequencies were particularly high and clustered for 4 CD8⁺ T-cell specificities, namely SCG5-009₁₈₆₋₁₉₄, UCN3₁₋₉, CCNI-008₁₄₋₂₂ and GAD2-003₁₇₉₋₁₈₇. As reported for PPI₆₋₁₄ and other known β-cell epitopes (Culina et al., 2018), these MMr⁺ cells were predominantly naïve in both T1D and healthy subjects (Fig. 5B; median 8.3%, interquartile range 0-20.0%).

Collectively, these results show that the β -cell peptides identified are targeted by similar frequencies of predominantly naïve circulating CD8⁺ T cells in T1D and healthy subjects.

Pancreas-infiltrating cells reactive to the HLA-A2-restricted IAPP_{15-17/5-10}, ISL1₂₇₆₋₂₈₄ and UCN3₁₋₉ peptides are enriched in T1D patients.

Given the lack of difference in frequency or markers of prior Ag encounter observed for circulating islet-reactive CD8⁺ T cells between T1D and healthy donors, we verified whether these reactivities were present in the pancreas-infiltrating cells of HLA-A2⁺ T1D, aAb⁺, non-diabetic and type 2 diabetes donors by *in-situ* MMr staining of consecutive tissue sections from the Network for Pancreatic Organ Donors (nPOD) repository. We selected IAPP_{15-17/5-10},

ISL1₂₇₆₋₂₈₄ and UCN3₁₋₉, which are representative of the frequency range detected in peripheral blood and stained positive in a preliminary screening on pancreatic sections from one T1D donor (Fig. S3; with SCG-009₁₈₆₋₁₉₄ also staining positive).Fig. 6A-R displays representative images, with scattered MMr⁺ cells in the vicinity of the islet or exocrine tissue. The β -cell ZnT8₁₈₆₋₁₉₄ peptide and the melanocyte MelanA₂₆₋₃₅ peptide provided positive and negative controls, respectively (Culina et al., 2018). Results are summarized in Fig. 6S and Table S5. Whereas IAPP_{15-17/5-10}, ISL1₂₇₆₋₂₈₄, UCN3₁₋₉ (and ZnT8₁₈₆₋₁₉₄) MMr⁺ cells were significantly more abundant than MelanA₂₆₋₃₅ MMr⁺ cells in T1D, aAb⁺ (barring UCN3₁₋₉), and non-diabetic cases (barring IAPP_{15-17/5-10} and UCN3₁₋₉), all islet MMr⁺ cells were enriched in the pancreata of T1D vs. non-diabetic cases, and also in the pancreata of aAb⁺ vs. non-diabetic cases for IAPP_{15-17/5-10} and ISL1₂₇₆₋₂₈₄. In contrast, islet and MelanA₂₆₋₃₅ MMr⁺ cells were present at similar densities across all groups in pancreatic lymph node (PLN) sections from the same donors (Fig. 6T), as reported for ZnT8₁₈₆₋₁₉₄ (Culina et al., 2018).

Fluorescent confocal microscopy on pancreas sections from one T1D donor (Fig. 6U-W) detected 60 CD8⁺ cells (2.3 cells/mm²), among which 37 (61.7%) were CD45RO⁺ and 2 CD45RO⁺MMr⁺ using pooled IAPP_{15-17/5-10}/ISL1₂₇₆₋₂₈₄/UCN3₁₋₉ MMrs (5.4% of CD8⁺CD45RO⁺ and 3.3% of total CD8⁺ cells), suggesting that pancreas-infiltrating islet MMr⁺ cells are Ag-experienced CD8⁺ T cells.

Collectively, these results show that IAPP_{15-17/5-10}-, ISL1₂₇₆₋₂₈₄- and UCN3₁₋₉-reactive cells are enriched in the pancreas of T1D patients, similarly to $ZnT8_{186-194}$ -reactive cells (Culina et al., 2018), lending support to their relevance in T1D.

DISCUSSION

We here provide a first catalogue of the HLA-I peptidome of human β cells, using an immortalized β -cell line naturally expressing the most prevalent HLA-A2 variant. This cellular model proved informative, since several HLA-A2-restricted peptides identified were also naturally processed and presented by primary human islets. The technical strengths of our approach are the combined HLA-I peptidomics and transcriptomics pipelines; the use of small cell numbers (20x10⁶) for HLA-I purification, despite its low expression in β cells compared with professional Ag-presenting cells; and the use of a mock immunopurification to exclude peptides not bound to HLA-I. One limitation is the lower sensitivity of the LC-MS/MS data-dependent acquisition discovery mode used compared with targeted strategies. Indeed, previous studies on mouse NIT-1 β cells (Dudek et al., 2012) detected low amounts of the immunodominant IGRP₂₀₆₋₂₁₄ peptide only after using a targeted approach on IFN- γ -treated cells. Nonetheless, our sensitivity proved sufficient to detect several known β -cell Ags. Although this did not allow a precise quantitation of pHLA-I complexes, it allowed to detect HLA-I-bound peptides without *a priori* hypotheses.

Expectedly, only ~5% of the HLA-I peptidome originated from proteins preferentially expressed in β cells. Multiple PPI peptides described as major CD8⁺ T-cell epitopes were detected, validating our discovery approach and documenting their natural processing and presentation by β cells. Peptides derived from all the other known β -cell Ags were also identified, namely CHGA, PTPRN, GAD2, SLC30A8 and IAPP. The only known Ag missing was IGRP, which may reflect low amounts of IGRP pHLA-I complexes, as reported for murine NIT-1 β cells (Dudek et al., 2012). The same may be true for the ZnT8₁₈₆₋₁₉₄ epitope, for which a concordant *m*/*z* value was identified at the MS1 level in several HLA-I-purified ECN90 samples (1/5, 5/5 and 4/5 for basal, IFN- γ -treated and IFN- γ /TNF- α /IL-1 β -treated conditions, respectively; 1/15 mock-purified samples). More importantly, several as yet

undescribed peptides were identified, many of which were derived from secretory granule proteins, namely CHGA, INS, SCG5, PTPRN, ABCC8, PCSK1, UCN3, CHGB, CPE, PCSK2, SCG3, SCL30A8, NPY and IAPP. This is not surprising considering that granule proteins are abundantly synthesized by β cells, thus favoring HLA-I presentation (Bassani-Sternberg et al., 2015). Their fast turnover also increases the odds of producing misfolded proteins, which are rapidly routed toward proteasomal degradation and HLA-I presentation (Anton and Yewdell, 2014).

mRNA alternative splicing is another mechanism frequently leading to unstable DRiPs, which are rapidly degraded through different pathways (Anton and Yewdell, 2014). Moreover, these mRNA isoforms may translate aa neo-sequences when exons are either added or skipped compared to the canonical mRNA. We therefore performed a parallel *in silico* prediction of mRNA-translated peptide neo-sequences. Although no proof of natural processing and presentation could be provided for most of these theoretical peptide products, the finding of a naïve CD8⁺ T-cell repertoire capable of recognizing them supports their potential relevance as autoimmune T-cell targets. Of note, peptides derived from the alternative open reading frame INS mRNA (Kracht et al., 2017) were not detected.

Despite presentation by HLA-I molecules, peptides may still be ignored by CD8⁺ T cells, thus not triggering an autoimmune response. This primarily reflects the absence of a cognate naïve repertoire available for priming (Alanio et al., 2010). We therefore first screened healthy individuals for the presence of cognate naïve CD8⁺ T cells, which were found for several peptides. Although the poor expression of the genes encoding these proteins in mTECs may exert a facilitating effect, this is not an absolute requirement for peripheral CD8⁺ T-cell recognition. Indeed, *CHGA*, *ISL1* and *SCG5* were expressed in mTECs, and yet targeted by CD8⁺ T cells at frequencies comparable to those of T cells recognizing Ags not expressed in

mTECs, in line with the increasing appreciation that thymic clonal deletion is highly incomplete (Culina et al., 2018).

Based on our previous findings on known β -cell epitopes (Culina et al., 2018), we did not expect differences in circulating CD8⁺ T cells between T1D and healthy subjects, because the Ag-experienced fraction is rather limited, likely reflecting sequestration in the target tissue. This was also the case for the candidates studied herein, and for the well-described control PPI₆₋₁₄ epitope eluted from pHLA-I complexes. Together with the reactivity against IAPP₁₅-17/5-10, ISL1₂₇₆₋₂₈₄ and UCN3 1-9, which was preferentially detected in the pancreatic infiltrates of T1D (and aAb⁺) donors and localized in CD8⁺CD45RO⁺ T cells in one T1D case, these findings provide a first validation of their disease relevance. The degree of evidence for a relevance to T1D is higher for those peptides targeted by CD8⁺ T cells and naturally processed and presented by β cells, i.e. SCG5₁₈₆₋₁₉₅, PCSK2₃₀₋₃₈, UCN3₁₋₉ and ISL1₂₇₆₋₂₈₄. These also include the neo-antigenic peptides SCG-009₁₈₆₋₁₉₄ and IAPP_{15-17/5-10} generated by mRNA splicing and transpeptidation, respectively.

Complementary analyses of the current HLA-I peptidomics dataset will yield additional information. First, only few PTMs were searched and a dedicated analysis is required. This should include the distinction between biological and experimentally induced PTMs, since some of them, e.g. the tryptophan to kynurenine conversion of the PPI₁₅₋₂₄ peptide, were similarly detected in the corresponding synthetic peptides. Second, an unbiased analysis of transpeptidation beyond the described aa preference rules (Berkers et al., 2015) will likely yield additional fusion peptides, which may account for up to one third of the HLA-I peptidome (Liepe et al., 2016). Third, only HLA-A2-restricted peptides were analyzed for T-cell recognition, leaving several HLA-A3- and HLA-B39-restricted candidates available for follow-up studies. HLA-B39 was expressed by the primary islets analyzed and, although rare, is the HLA-I variant most strongly associated with T1D (Nejentsev et al., 2007).

Finally, the HLA-I peptidome obtained allows to formulate hypotheses about the Agprocessing pathways employed by β cells. Some peptides (UCN3₁₋₉, IAPP_{15-17/5-10}, PPI₂₋₁₀, PPI_{6-14} , PPI_{15-24}) are located in the leader sequence of proteins abundantly produced by β cells, which is cleaved in the ER at each protein synthesis. These byproducts may therefore provide a rich peptide source for HLA-I presentation and likely follow alternative Ag-processing pathways within the ER, independent of proteasome cleavage (El Hage et al., 2008; Skowera et al., 2008). It is also noteworthy that several source proteins of HLA-I-bound peptides, i.e. CHGA, INS, SCG5, PCSK1, UCN3, CHGB, CPE, PCSK2, SCG3, NPY and IAPP are synthesized as precursors and incorporated into β -cell granules, where they undergo intermediate processing by proconvertases to yield bioactive products. A notable example is SCG5, a PCSK2 chaperone that is gradually degraded along the secretory pathway to competitively prevent the premature activation of PCSK2 by autocatalytic cleavage (Mbikay et al., 2001). This continuous degradation may explain the abundance of HLA-I-bound SCG5 peptides. In this respect, the C-terminal SCG5₁₈₆₋₁₉₅ peptide is located between furin (RRKRR) and PCSK2 (KK) cleavage sites and, similar to leader sequence peptides, may behave as a byproduct of the intermediate SCG5 processing (Bartolomucci et al., 2011). The same is true for several CHGA peptides, e.g. CHGA344-352, which maps to the WE-14 neuropeptide produced by CHGA cleavage at dibasic KR motifs (Bartolomucci et al., 2011). These peptides may access the HLA-I pathway following crinophagy, i.e. the disposal of unused secretory granules through fusion with lysosomes (Goginashvili et al., 2015; Weckman et al., 2014). In this scenario, islet inflammation may provide a key switch for progression of the 'benign' autoimmunity of healthy individuals toward T1D at two levels: on T cells, by impairing peripheral immunoregulation; and on β cells, by increasing the overall number of pHLA-I complexes available for T-cell recognition.

LIMITATIONS OF THIS STUDY

One limitation of our work is that the pathogenic role of the CD8⁺ T cells recognizing the epitopes described herein remains to be definitely established. Islets from T1D patients were not available for HLA peptidomics studies, and the enrichment of these T cells in the pancreas of patients could reflect a cause or consequence of disease. It should be noted however that the same uncertainty applies to all the other islet-reactive T-cell specificities described to date. The capacity to kill β cells that has been described for some of them cannot be taken as conclusive evidence for their pathogenic role, since T-cell clones obtained from T1D and healthy donors kill β cells to a similar extent (Culina et al., 2018). Nonetheless, we here provide two key findings. First, that these epitopes are naturally processed and presented by β cells and are covered by the natural CD8⁺ T-cell repertoire, hence providing potential targets for β -cell killing. Second, that the CD8⁺ T cells recognizing them accumulate preferentially in the pancreas of T1D patients. This latter finding is unlikely to be serendipitous, as it did not apply to CD8⁺ T cells recognizing the melanocyte MelanA₂₆₋₃₅ epitope, despite the fact that their frequency in the blood is ~100-fold higher than for islet epitope-reactive CD8⁺ T cells (Culina et al., 2018).

In conclusion, the HLA-I peptidome of human β cells described herein provides information about the Ag processing features of β cells, the targets amenable to autoimmune recognition and a valuable tool for developing T-cell biomarkers and tolerogenic vaccines. Acknowledgements. We thank A. Jones, K. Zehrouni, and the Diabetology nurse and medical staff of the Cochin and A. Mignot Hospital for patient recruitment; C. Maillard for technical assistance; Univercell Biosolutions for the ECN90 β-cell line; the CyBio platform (Cochin Institute) for assistance with flow cytometry; and T. Loukanov (University of Heidelberg) for human thymi. This work was performed within the Département Hospitalo-Universitaire (DHU) AutHorS and supported by the *Ile-de-France* CORDDIM and by grants from the JDRF (1-PNF-2014-155-A-V, 2-SRA-2016-164-Q-R), the Agence Nationale de la Recherche (ANR-2015-CE17-0018-01), the Fondation pour la Recherche Médicale (DEQ20140329520), the Société Francophone du Diabète, the Association pour la Recherche sur le Diabète, and the Helmsley Charitable Trust Eisenbarth nPOD Award for Team Science (2015PG-T1D052), to RM; Lilly and Fondation Bettencourt-Schueller, to RS; European Research Council (ERC-2012-AdG), to BK; FRFSWelbio (CR-2015A-06) and NIH-NIDDK-HIRN Consortium (1UC4DK104166-01), to DLE; and Ile-de-France SESAME 2010 (10022268), the City of Paris and ESPCI Paris, to JV. This project received funding from the Innovative Medicines Initiative 2 Joint Undertaking (INNODIA, 115797), which receives support from the EU Horizon 2020 program, the European Federation of Pharmaceutical Industries and Associations, JDRF and the Helmsley Charitable Trust. This research was supported by nPOD, a collaborative T1D research project funded by JDRF. Organ Procurement Organizations partnering with nPOD are listed at www.jdrfnpod.org/ourpartners.php.

Author contributions. Conceptualization, D.L.E., R.M.; Methodology, S.G.D., M.E.A., M.L.C., G.A., J.V.T., A.C., A.I.L., F.D., Y.V., J.V., R.M.; Software, S.G.D.; Investigation, S.G.D., M.E.A., M.L.C., G.A., J.V.T., L.N., A.I.L., G.S., A.C., S.P., S.C., N.C., L.M.S., S.Y., F.D., D.L.E., Y.V., J.V., R.M.; Resources, S.P., M.B., P.M., M.A., M.D., B.K., L.M.S., S.B., D.D.L., E.L., J.P.B., G.B., R.S.; Data Curation, S.G.D., M.L.C., J.V.T., D.L.E., Y.V., J.V.,

R.M.; Writing – Original Draft, D.L.E., Y.V., R.M.; Writing – Review & Editing, S.G.D., S.Y., M.L.C., D.L.E., Y.V., J.V., R.M.; Visualization, S.G.D., M.E.A., M.L.C., G.A., J.V.T., L.N., G.S., Y.V., R.M.; Supervision, B.K., F.D., D.L.E., Y.V., J.V., R.M.; Funding Acquisition, S.P., B.K., S.Y., D.L.E., J.V., R.M.

Declaration of Interests. Some peptide sequences described herein are covered by pending patents filed by Inserm Transfert.

REFERENCES

Alanio, C., Lemaitre, F., Law, H.K., Hasan, M., and Albert, M.L. (2010). Enumeration of human antigen-specific naive CD8+ T cells reveals conserved precursor frequencies. Blood *115*, 3718-3725.

Anton, L.C., and Yewdell, J.W. (2014). Translating DRiPs: MHC class I immunosurveillance of pathogens and tumors. J. Leukoc. Biol. *95*, 551-562.

Babon, J.A., DeNicola, M.E., Blodgett, D.M., Crevecoeur, I., Buttrick, T.S., Maehr, R., Bottino, R., Naji, A., Kaddis, J., Elyaman, W., et al. (2016). Analysis of self-antigen specificity of islet-infiltrating T cells from human donors with type 1 diabetes. Nat. Med. 22, 1482-1487.

Bartolomucci, A., Possenti, R., Mahata, S.K., Fischer-Colbrie, R., Loh, Y.P., and Salton, S.R. (2011). The extended granin family: structure, function, and biomedical implications. Endocr. Rev. *32*, 755-797.

Bassani-Sternberg, M., Pletscher-Frankild, S., Jensen, L.J., and Mann, M. (2015). Mass spectrometry of human leukocyte antigen class I peptidomes reveals strong effects of protein abundance and turnover on antigen presentation. Mol. Cell. Proteomics *14*, 658-673.

Berkers, C.R., de Jong, A., Schuurman, K.G., Linnemann, C., Meiring, H.D., Janssen, L., Neefjes, J.J., Schumacher, T.N., Rodenko, B., and Ovaa, H. (2015). Definition of proteasomal peptide splicing rules for high-efficiency spliced peptide presentation by MHC Class I molecules. J. Immunol. *195*, 4085-4095.

Caron, E., Espona, L., Kowalewski, D.J., Schuster, H., Ternette, N., Alpizar, A., Schittenhelm, R.B., Ramarathinam, S.H., Lindestam Arlehamn, C.S., Chiek Koh, C., et al. (2015). An open-source computational and data resource to analyze digital maps of immunopeptidomes. eLife 4.

Coppieters, K.T., Dotta, F., Amirian, N., Campbell, P.D., Kay, T.W., Atkinson, M.A., Roep, B.O., and von Herrath, M.G. (2012). Demonstration of islet-autoreactive CD8 T cells in insulitic lesions from recent onset and long-term type 1 diabetes patients. J. Exp. Med. 209, 51-60.

Culina, S., Lalanne, A.I., Afonso, G., Cerosaletti, K., Pinto, S., Sebastiani, G., Kuranda, K., Nigi, L., Eugster, A., Osterbye, T., et al. (2018). Islet-reactive CD8+ T cell frequencies in the pancreas, but not in blood, distinguish type 1 diabetic patients from healthy donors. Sci Immunol *3*, eaao4013.

Delong, T., Wiles, T.A., Baker, R.L., Bradley, B., Barbour, G., Reisdorph, R., Armstrong, M., Powell, R.L., Reisdorph, N., Kumar, N., et al. (2016). Pathogenic CD4 T cells in type 1 diabetes recognize epitopes formed by peptide fusion. Science *351*, 711-714.

Dudek, N.L., Tan, C.T., Gorasia, D.G., Croft, N.P., Illing, P.T., and Purcell, A.W. (2012). Constitutive and inflammatory immunopeptidome of pancreatic beta-cells. Diabetes *61*, 3018-3025.

Eizirik, D.L., Colli, M.L., and Ortis, F. (2009). The role of inflammation in insulitis and betacell loss in type 1 diabetes. Nat. Rev. Endocrinol. *5*, 219-226.

Eizirik, D.L., Sammeth, M., Bouckenooghe, T., Bottu, G., Sisino, G., Igoillo-Esteve, M., Ortis, F., Santin, I., Colli, M.L., Barthson, J., et al. (2012). The human pancreatic islet transcriptome: expression of candidate genes for type 1 diabetes and the impact of pro-inflammatory cytokines. PLoS Genet. 8, e1002552.

El Hage, F., Stroobant, V., Vergnon, I., Baurain, J.F., Echchakir, H., Lazar, V., Chouaib, S., Coulie, P.G., and Mami-Chouaib, F. (2008). Preprocalcitonin signal peptide generates a cytotoxic T lymphocyte-defined tumor epitope processed by a proteasome-independent pathway. Proc. Natl. Acad. Sci. U.S.A. *105*, 10119-10124.

Goginashvili, A., Zhang, Z., Erbs, E., Spiegelhalter, C., Kessler, P., Mihlan, M., Pasquier, A., Krupina, K., Schieber, N., Cinque, L., et al. (2015). Insulin granules. Insulin secretory granules control autophagy in pancreatic beta cells. Science *347*, 878-882.

Gubin, M.M., Zhang, X., Schuster, H., Caron, E., Ward, J.P., Noguchi, T., Ivanova, Y., Hundal, J., Arthur, C.D., Krebber, W.J., et al. (2014). Checkpoint blockade cancer immunotherapy targets tumour-specific mutant antigens. Nature *515*, 577-581.

James, E.A., Abreu, J.R.F., McGinty, J.W., Odegard, J.M., Fillie, Y.E., Hocter, C.N., Culina, S., Ladell, K., Price, D.A., Alkanani, A., et al. (2018). Combinatorial detection of autoreactive CD8+ T cells with HLA-A2 multimers: a multi-centre study by the Immunology of Diabetes Society T Cell Workshop. Diabetologia *61*, 658-670.

Kracht, M.J., van Lummel, M., Nikolic, T., Joosten, A.M., Laban, S., van der Slik, A.R., van Veelen, P.A., Carlotti, F., de Koning, E.J., Hoeben, R.C., et al. (2017). Autoimmunity against a defective ribosomal insulin gene product in type 1 diabetes. Nat. Med. *23*, 501-507.

Li, Y., Zhou, L., Li, Y., Zhang, J., Guo, B., Meng, G., Chen, X., Zheng, Q., Zhang, L., Zhang, M., et al. (2015). Identification of autoreactive CD8+ T cell responses targeting chromogranin A in humanized NOD mice and type 1 diabetes patients. Clin. Immunol. *159*, 63-71.

Liepe, J., Marino, F., Sidney, J., Jeko, A., Bunting, D.E., Sette, A., Kloetzel, P.M., Stumpf, M.P., Heck, A.J., and Mishto, M. (2016). A large fraction of HLA class I ligands are proteasome-generated spliced peptides. Science *354*, 354-358.

Mallone, R., Martinuzzi, E., Blancou, P., Novelli, G., Afonso, G., Dolz, M., Bruno, G., Chaillous, L., Chatenoud, L., Bach, J.M., et al. (2007). CD8+ T-cell responses identify beta-cell autoimmunity in human type 1 diabetes. Diabetes *56*, 613-621.

Marroqui, L., Dos Santos, R.S., Floyel, T., Grieco, F.A., Santin, I., Op de Beeck, A., Marselli, L., Marchetti, P., Pociot, F., and Eizirik, D.L. (2015). TYK2, a candidate gene for type 1 diabetes, modulates apoptosis and the innate immune response in human pancreatic beta-cells. Diabetes *64*, 3808-3817.

Marroqui, L., Dos Santos, R.S., Op de Beeck, A., Coomans de Brachene, A., Marselli, L., Marchetti, P., and Eizirik, D.L. (2017). Interferon-alpha mediates human beta cell HLA class I overexpression, endoplasmic reticulum stress and apoptosis, three hallmarks of early human type 1 diabetes. Diabetologia *60*, 656-667.

Marroqui, L., Santin, I., Dos Santos, R.S., Marselli, L., Marchetti, P., and Eizirik, D.L. (2014). BACH2, a candidate risk gene for type 1 diabetes, regulates apoptosis in pancreatic beta-cells via JNK1 modulation and crosstalk with the candidate gene PTPN2. Diabetes *63*, 2516-2527.

Martinuzzi, E., Novelli, G., Scotto, M., Blancou, P., Bach, J.M., Chaillous, L., Bruno, G., Chatenoud, L., van, E.P., and Mallone, R. (2008). The frequency and immunodominance of islet-specific CD8+ T-cell responses change after type 1 diabetes diagnosis and treatment. Diabetes *57*, 1312-1320.

Mbikay, M., Seidah, N.G., and Chretien, M. (2001). Neuroendocrine secretory protein 7B2: structure, expression and functions. Biochem. J. *357*, 329-342.

McGinty, J.W., Chow, I.T., Greenbaum, C., Odegard, J., Kwok, W.W., and James, E.A. (2014). Recognition of post-translationally modified glutamic acid decarboxylase 65 epitopes in subjects with type 1 diabetes. Diabetes *63*, 3033-3040.

Montgomery, S.B., Sammeth, M., Gutierrez-Arcelus, M., Lach, R.P., Ingle, C., Nisbett, J., Guigo, R., and Dermitzakis, E.T. (2010). Transcriptome genetics using second generation sequencing in a Caucasian population. Nature *464*, 773-777.

Moore, F., Colli, M.L., Cnop, M., Esteve, M.I., Cardozo, A.K., Cunha, D.A., Bugliani, M., Marchetti, P., and Eizirik, D.L. (2009). PTPN2, a candidate gene for type 1 diabetes, modulates interferon-gamma-induced pancreatic beta-cell apoptosis. Diabetes *58*, 1283-1291.

Nejentsev, S., Howson, J.M., Walker, N.M., Szeszko, J., Field, S.F., Stevens, H.E., Reynolds, P., Hardy, M., King, E., Masters, J., et al. (2007). Localization of type 1 diabetes susceptibility to the MHC class I genes HLA-B and HLA-A. Nature *450*, 887-892.

Ortis, F., Naamane, N., Flamez, D., Ladriere, L., Moore, F., Cunha, D.A., Colli, M.L., Thykjaer, T., Thorsen, K., Orntoft, T.F., et al. (2010). Cytokines interleukin-1beta and tumor necrosis factor-alpha regulate different transcriptional and alternative splicing networks in primary beta-cells. Diabetes *59*, 358-374.

Pinto, S., Sommermeyer, D., Michel, C., Wilde, S., Schendel, D., Uckert, W., Blankenstein, T., and Kyewski, B. (2014). Misinitiation of intrathymic MART-1 transcription and biased TCR usage explain the high frequency of MART-1-specific T cells. Eur. J. Immunol. *44*, 2811-2821.

Ravassard, P., Hazhouz, Y., Pechberty, S., Bricout-Neveu, E., Armanet, M., Czernichow, P., and Scharfmann, R. (2011). A genetically engineered human pancreatic beta cell line exhibiting glucose-inducible insulin secretion. J. Clin. Invest. *121*, 3589-3597.

Rondas, D., Crevecoeur, I., D'Hertog, W., Bomfim Ferreira, G., Staes, A., Garg, A.D., Eizirik, D.L., Agostinis, P., Gevaert, K., Overbergh, L., et al. (2015). Citrullinated glucose-regulated protein 78 is an autoantigen in type 1 diabetes. Diabetes *64*, 573-586.

Scotto, M., Afonso, G., Larger, E., Raverdy, C., Lemonnier, F.A., Carel, J.C., Dubois-Laforgue, D., Baz, B., Levy, D., Gautier, J.F., et al. (2012). Zinc transporter (ZnT)8(186-194) is an immunodominant CD8+ T cell epitope in HLA-A2+ type 1 diabetic patients. Diabetologia *55*, 2026-2031.

Skowera, A., Ellis, R.J., Varela-Calvino, R., Arif, S., Huang, G.C., Van-Krinks, C., Zaremba, A., Rackham, C., Allen, J.S., Tree, T.I., et al. (2008). CTLs are targeted to kill beta cells in patients with type 1 diabetes through recognition of a glucose-regulated preproinsulin epitope. J. Clin. Invest. *118*, 3390-3402.

Standifer, N.E., Ouyang, Q., Panagiotopoulos, C., Verchere, C.B., Tan, R., Greenbaum, C.J., Pihoker, C., and Nepom, G.T. (2006). Identification of novel HLA-A*0201-restricted epitopes in recent-onset type 1 diabetic subjects and antibody-positive relatives. Diabetes *55*, 3061-3067.

Villate, O., Turatsinze, J.V., Mascali, L.G., Grieco, F.A., Nogueira, T.C., Cunha, D.A., Nardelli, T.R., Sammeth, M., Salunkhe, V.A., Esguerra, J.L., et al. (2014). Noval is a master regulator of alternative splicing in pancreatic beta cells. Nucleic Acids Res. *42*, 11818-11830. Weckman, A., Di Ieva, A., Rotondo, F., Syro, L.V., Ortiz, L.D., Kovacs, K., and Cusimano, M.D. (2014). Autophagy in the endocrine glands. J. Mol. Endocrinol. *52*, R151-163.

FIGURES

Figure 1. The HLA-I peptidome of human β cells is enriched by cytokine exposure and displays the expected amino-acid length and motifs. (A) HLA-I expression detected by flow cytometry on ECN90 β cells under basal (red), IFN- γ -stimulated (blue) and IFN- γ /TNF- $\alpha/IL-1\beta$ -stimulated conditions (green). The grey profile displays IFN- $\gamma/TNF-\alpha/IL-1\beta$ stimulated cells stained with an isotype control monoclonal antibody (mAb; the identical profiles of basal and IFN- γ -stimulated conditions are not shown). (**B**) HLA-I heavy chain expression detected by Western blot on the same cell aliquots, with α -tubulin bands shown as loading control. (C) as length of the 2,997 peptides detected in ≥ 2 out of 5 biological replicates. (D) Sequence logo plots displaying the HLA-A2 (top), -A3 (middle) and -B40/B49 (bottom) binding motifs for the 3 major Gibbs clusters. The X-axis shows the residue position within nonamer peptide sequences. The Y-axis shows the information content, with the size of each aa symbol proportional to its frequency. (E) Heat map of the predicted HLA-I binding affinities of nonamer peptides. Each column represents a unique peptide. Red, green and grey shades indicate high binders (K_D≤50 nM), low binders (K_D≤2,000 nM) and non-binders (K_D>2,000 nM), respectively. (F) Number of peptides identified in ECN90 β cells. **p=0.008 and p=0.02 by Mann-Whitney test. (G) The bioinformatics analysis pipeline used (see STAR methods for details). The 3,544 peptides identified were first filtered based on inter-sample reproducibility (n=2,997) and aa length (n=2,795). From this dataset, we identified 217 peptides derived from ubiquitous proteins carrying PTMs (blue), from which 99 (Table S2) were enriched in HLA-I-purified samples; and 15 peptide splice variants (brown; see Fig. S1 for the identification strategy), from which 10 were enriched in HLA-I-purified samples. The remaining β-cell-enriched peptides (both conventional and with PTMs; red) were filtered based on non-ubiquitous expression (n=411), β -cell-enriched expression (n=139) and enrichment in HLA-I-purified samples (n=86; Table S1). Finally, mRNA splice variants (green, n=2) were identified by searching the HLA-I peptidomics dataset against the predicted aa neo-sequences obtained from RNAseq analysis. For this RNAseq pipeline (dashed boxes), 53,280 mRNAs were filtered based on islet mRNA expression (n=14,504), low mTEC mRNA expression (n=908), islet enrichment compared to control tissues (n=166) and generation of a predicted aa neo-sequence (n=88). The 336 predicted aa neo-sequences obtained from these 88 mRNA variants were analyzed in parallel for the presence of predicted HLA-2 binders (n=66) with a 9-10-aa length (n=43) and a neo-sequence of \geq 3 aa (n=39). The indicated epitope candidates were further filtered on predicted and experimental HLA-A2 binding to focus subsequent CD8⁺ T-cell recognition studies on HLA-A2-restricted peptides (see text).

Figure 2. pHLA-I complexes of human β cells are enriched in peptides derived from secretory granule proteins, including known PPI epitopes. (**A**) Number of peptides (n=86 from β-cell-enriched proteins, n=10 transpeptidation products, n=2 mRNA splice products) identified in ECN90 β cells after bioinformatics filtering (Fig. 1G). (**B**) Source proteins of the 98 peptides (detailed in Table S1). (**C**) Mapping of HLA-A2-restricted PPI peptides identified in ECN90 β cells. PTM variants (see Table S1) and a PPI₁₅₋₂₆ length variant (in italics) were also identified. (**D**) Origin of the 42 proteins yielding the 98 peptides identified in ECN90 β cells, including splice peptides generated from IAPP and NPY. (**E**) Predicted HLA-I restrictions of the same peptides. (**F**) Source proteins of the 33 peptides (Table S3) identified in HLA-A2⁺ primary human islets. Asterisks indicate proteins for which some peptides identified were identical or length variants compared to those identified in ECN90 β cells, including the identical IAPP_{15-17/5-10} splice peptide. Exploded slices indicate source proteins not identified in ECN90 β cells. (**G**) HLA-I heavy chain expression detected by Western blot on primary islets and ECN90 β cells under basal and IFN-γ/TNF-α/IL-1β-stimulated conditions, with β -actin bands shown as loading control. A representative islet preparation out of 3 analyzed is shown.

Figure 3. Gating strategy for the combinatorial analysis of β-cell peptide MMr⁺CD8⁺ T cells in T1D and healthy subjects. (**A**) Frozen-thawed peripheral blood mononuclear cells (PBMCs) from T1D donor D314D were magnetically depleted of CD8⁻ cells before staining, acquisition and analysis (see STAR Methods). Cells were sequentially gated on small lymphocytes, singlets, live cells (Live/Dead Aqua⁻), CD3⁺CD8⁺ T cells and total PE⁺, PE-CF594⁺, APC⁺, BV650⁺, BV711⁺ and BV786⁺ MMr⁺ T cells. Using Boolean operators, these latter gates allowed to selectively visualize each double-MMr⁺ population by including only those events positive for the corresponding fluorochrome pair. (**B-C**) The final readout obtained for T1D donor D314D (B) and healthy donor H170S (C) is shown for the 15 peptides analyzed. Events corresponding to each epitope-reactive population are overlaid in different colors within each plot, with MMr⁻ events overlaid in light grey. The small dot plots on the right of each panel depict CD45RA (x-axis) and CCR7 (y-axis) expression in the corresponding MMr⁺ fraction. Numbers in each panel indicate the MMr⁺CD8⁺ T-cell frequency out of total CD8⁺ T cells and the percent naïve (CD45RA⁺CCR7⁺) fraction among MMr⁺ cells.

Figure 4. HLA-A2-restricted β-cell peptides are targeted by a circulating naïve CD8⁺ Tcell repertoire in healthy donors. MMr⁺CD8⁺ cells reactive to HLA-A2-binding β-cell peptides (Fig. S2) were stained *ex vivo* from PBMCs of 5-6 HLA-A2⁺ healthy donors (Table S4). (A-F) Representative dot plots of different MMr staining patterns: high frequency, clustered pattern (A; CCNI-008₁₄₋₂₂); intermediate frequency, clustered pattern (B; SCG5₁₈₆₋₁₉₅); low frequency, clustered pattern (C; CHGB₄₄₀₋₄₄₈); high frequency, spread pattern (D; LARP4-006₂₁₄₋₂₂₂); and the β -cell PPI₆₋₁₄ (E) and viral Flu MP₅₈₋₆₆ (F) positive controls. (G-H) MMr⁺CD8⁺ cells reactive to β -cell peptides identified in HLA-I peptidomics (G) and RNAseq (H) pipelines. PPI₆₋₁₄ and Flu MP₅₈₋₆₆ peptides were included as controls. Frequencies out of total CD8⁺ T cells are depicted. Dotted lines indicate the expected frequency of naïve MMr⁺CD8⁺ T cells, bars show median values. The 20 β -cell peptides displaying the expected CD8⁺ T-cell frequency and a clustered MMr staining pattern were retained and are marked with an asterisk (2 asterisks for the 13 peptides further analyzed). β cell peptides in italics were excluded despite their cognate MMr⁺CD8⁺ T-cell frequencies because of their spread MMr staining pattern. At least 0.6x10⁶ CD8⁺ T cells were counted for each donor (median 1.4x10⁶, range 0.6-4.9x10⁶). (I-J) Percent Ag-experienced cells (CD45RA⁻CCR7⁻, CD45RA⁺CCR7⁻ and CD45RA⁻CCR7⁺) out of total MMr⁺ cells for the β cell peptides depicted in panels G-H, respectively. Data points with <5 MMr⁺ cells were excluded from this analysis (median 19 MMr⁺ cells, range 5-808 for β -cell peptides). NA, not available.

Figure 5. Circulating CD8⁺ T cells reactive to HLA-A2-restricted β-cell peptides display similar *ex-vivo* frequencies and a predominantly naïve phenotype in T1D and healthy subjects. (A) Frequencies of MMr⁺CD8⁺ cells reactive to the indicated β-cell peptides in T1D (grey; n=10) and healthy subjects (white; n=10; donors listed in Table S4). PPI₆₋₁₄ and Flu MP₅₈₋₆₆ peptides were included as controls. At least $0.6x10^6$ CD8⁺ T cells were counted for each donor (median $1.0x10^6$, range $0.6-2.4x10^6$). (B) Percent Ag-experienced cells out of total MMr⁺ cells for the β-cell peptides depicted in panel A. Data points with <5 MMr⁺ cells were excluded (median 14 MMr⁺ cells, range 5-160 for β-cell peptides). Bars show median values. NA, not available. Fig. 6. Pancreas-infiltrating cells reactive to the HLA-A2-restricted IAPP_{15-17/5-10}, ISL1₂₇₆₋₂₈₄ and UCN3₁₋₉ peptides are enriched in T1D patients. Pancreas sections from nPOD cases (Table S5) were immunohistochemically stained *in situ* with MMrs loaded with IAPP_{15-17/5-10}, ISL1₂₇₆₋₂₈₄, UCN3₁₋₉ (selected from Fig. S3), positive control ZnT8₁₈₆₋₁₉₄ and negative control melanocyte MelanA₂₆₋₃₅ peptide. (**A-D**, **I-L**, **Q-R**) Representative pancreas images (20X magnification; scale bar 100 µm). (**E-H**, **M-P**) Higher magnification of the dotted areas highlighted on the left of each panel (scale bar 40 µm for panels E, M; 38 µm for F, N; 45 µm for G, O; 36 µm for H, P). (**S-T**) Number of MMr⁺ cells/mm² section area of pancreas (S) and PLNs (T). Each point represents an individual case, bars indicate median values. *P ≤ 0.05 and **P ≤ 0.009 by Mann-Whitney test. NA, not assessed. (**U-W**) Fluorescent confocal microscopy on pancreas sections from T1D EUnPOD case #060217 for CD8 (green), CD45RO (blue) and pooled IAPP_{15-17/5-10}/ISL1₂₇₆₋₂₈₄/UCN3₁₋₉ MMrs (red). Examples of CD8⁺CD45RO⁻MMr⁻, CD8⁺CD45RO⁺MMr⁻ and CD8⁺CD45RO⁺MMr⁺ cells are shown in panel U (scale bar 25 µm), V and W (scale bar 10 µm), respectively.

TABLES

Table 1. Summary of β-cell peptides tested for recognition by circulating naïve CD8⁺ T cells of healthy subjects. The peptides presented in Fig. 4 are alphabetically listed and classified according to type (conventional, peptide splice and mRNA splice). The corresponding median RPKM values detected in islets (control- and cytokine-treated) and mTECs (HLA Class II^{hi} and Class II^{lo}) are shown. For proteins not arising from alternative mRNA splicing, RPKM values refer to the most prevalent islet mRNA isoform coding for the corresponding peptide. Subsequent columns detail the number of T-cell⁺ donors (\geq 5/10⁷ MMr⁺ cells counted), the median MMr⁺ frequency, the median percent Agexperienced cells within the MMr⁺ fraction (for peptides/donors with \geq 5 MMr⁺ cells), the number of donors with \geq 5 MMr⁺ cells, the MMr staining pattern (clustered, spread or not detected, ND) and the final validation outcome. The 20 validated β-cell peptides (in bold) displayed the expected CD8⁺ T-cell frequency and a clustered MMr staining pattern and are marked with an asterisk (2 asterisks for the 13 peptides further analyzed). NA, not available.

Protein or mRNA	Peptide	Sequence	Туре	Islet RPKM, control	lslet RPKM, cytokines	mTEC RPKM, HLA-II ^{hi}	mTEC RPKM, HLA-III⁰	T-cell⁺ donors	Frequency among CD8⁺ cells	% Ag experienced	Donors with ≥5 MMr⁺ cells	MMr staining pattern	Validation
ACLY-004	70-78	GLVGVNLTL	mRNA splice	22.5	5.7	0.0	0.0	0/6	0.00E+00	NA	0/6	ND	No
C15orf48-003	73-81	FLLQNPCPL	mRNA splice	9.8	18.1	0.0	0.0	0/6	8.30E-08	29	1/6	Spread	No
C16orf70-010	44-53	VLYSEQVIEV	mRNA splice	0.7	12.3	0.0	0.0	0/6	2.54E-07	14	1/6	Spread	No
CCNI-008	11-19	ILDKLNWDL	mRNA splice	41.0	25.9	0.4	0.3	1/6	1.05E-07	20	1/6	Spread	No
**CCNI-008	14-22	KLNWDLHTA	mRNA splice	41.0	25.9	0.4	0.3	4/6	5.05E-06	10	6/6	Clustered	Yes
CCNI-008	53-61	SLPLNSVYV	mRNA splice	41.0	25.9	0.4	0.3	1/6	5.20E-08	NA	0/6	Spread	No
*CHGA	344-352	KMDQLAKEL	Conventional	587.2	1881.9	7.9	13.2	3/6	9.75E-07	18	5/6	Clustered	Yes
CHGA	402-411	SLEAGLPLQV	Conventional	587.2	1881.9	7.9	13.2	0/6	1.66E-07	NA	0/6	Spread	No
CHGB	440-448	FLGEGHHRV	Conventional	1212.4	935.2	9.6	18.4	1/6	6.67E-07	28	4/6	Clustered	No
CLDN7-008	60-68	GMMSCKIGL	mRNA splice	20.8	17.5	0.0	0.1	0/6	0.00E+00	NA	0/6	ND	No
FAM171B	19-28	VLLKARLVPA	Conventional	4.3	4.0	6.5	5.0	3/6	1.70E-06	7	5/6	Spread	No
**GAD2-003	179-187	KIIKLFFRL	mRNA splice	22.2	21.2	0.0	0.0	5/6	5.14E-06	19	6/6	Clustered	Yes
GNAS-002	31-39	ALLWLSCSI	mRNA splice	256.1	123.2	0.6	0.0	0/6	4.92E-07	30	2/6	Clustered	No
GNAS-002	32-41	LLWLSCSIAL	mRNA splice	256.1	123.2	0.6	0.0	0/6	0.00E+00	NA	0/6	ND	No
GNAS-002	34-42	WLSCSIALL	mRNA splice	256.1	123.2	0.6	0.0	1/6	3.16E-07	23	2/6	Clustered	No
**GNAS-036	67-75	YMCTHRLLL	mRNA splice	48.9	45.1	0.0	0.0	4/6	1.25E-06	9	5/6	Clustered	Yes
GNAS-036	67-76	YMCTHRLLLL	mRNA splice	48.9	45.1	0.0	0.0	0/6	0.00E+00	NA	0/6	ND	No
**GNAS-036	124-132	AMSNLVPPV	mRNA splice	48.9	45.1	0.0	0.0	5/6	1.30E-06	14	5/6	Clustered	Yes
GNAS-036	185-194	QLIDCAQYFL	mRNA splice	48.9	45.1	0.0	0.0	0/6	4.76E-08	NA	0/6	ND	No
GPR119	48-56	AVADTLIGV	Conventional	9.4	7.4	0.0	0.0	0/6	1.26E-07	NA	0/6	Spread	No
**IAPP/IAPP	15-17/5-10	VALKLQVFL	Peptide splice	1257.1	392.4	0.3	0.0	1/6	2.45E-07	9	1/6	Clustered	Yes
**IAPP-002	33-42	VLSRNILLEL	mRNA splice	74.0	24.3	0.0	0.0	3/6	9.63E-07	35	4/6	Clustered	Yes
IAPP-004	9-18	CLDQIPIFTV	mRNA splice	609.8	257.1	0.0	0.0	1/6	2.59E-07	37	2/6	Clustered	No
IGF2BP3	552-560	KIQEILTQV	Conventional	0.2	0.4	5.3	0.9	0/5	2.66E-07	24	4/5	Spread	No

**ISL1	276-284	GLQANPVEV	Conventional	26.2	27.9	17.0	26.9	3/6	1.83E-06	26	3/6	Clustered	Yes
*KCNK16	129-137	ALLGIPLNV	Conventional	33.3	22.5	0.0	0.0	5/6	4.44E-06	13	6/6	Clustered	Yes
KCNK16-002	234-242	SLAAIWILL	mRNA splice	33.3	22.5	0.0	0.0	0/6	4.37E-08	NA	0/6	ND	No
KCNK16-002	240-248	ILLGLAWLA	mRNA splice	33.3	22.5	0.0	0.0	0/6	1.30E-07	NA	0/6	Spread	No
*KIF1A	1347-1355	VLDTSVAYV	Conventional	19.5	17.7	1.0	1.9	3/6	2.25E-06	33	5/6	Clustered	Yes
KIF1A	1480-1488	KLSEMSVTL	Conventional	19.5	17.7	1.0	1.9	0/5	5.2E-07	18.2	5/5	Spread	No
LARP4-006	214-222	RLMDSSIYS	mRNA splice	6.8	10.5	0.0	0.0	2/6	8.95E-07	20	4/6	Spread	No
LARP4-006	355-363	YLQKETSTL	mRNA splice	6.8	10.5	0.0	0.0	2/6	7.94E-07	21	5/6	Spread	No
LGMN-012	68-76	VMINPTPGI	mRNA splice	0.0	10.9	0.0	0.0	0/6	0.00E+00	NA	0/6	ND	No
PCSK1-002	8-17	FLFFSQIGSL	mRNA splice	16.2	13.8	0.0	0.0	0/6	0.00E+00	NA	0/6	ND	No
**PCSK2	30-38	FTNHFLVEL	Conventional	73.2	43.4	0.7	0.0	4/6	1.25E-06	9	6/6	Clustered	Yes
PCSK2-001	11-19	AAAGFLFCV	mRNA splice	36.5	17.8	0.0	0.1	2/6	5.40E-07	23	2/6	Spread	No
PCSK2-001	15-23	FLFCVMVFA	mRNA splice	36.5	17.8	0.0	0.1	2/6	3.79E-07	54	2/6	Spread	No
PDXDC1-013	174-183	YLCNQDVAFL	mRNA splice	13.5	0.8	0.0	0.0	2/6	6.21E-07	17	3/6	Spread	No
PIK3R3	144-152	SLAQYNPKL	Conventional	2.2	2.4	18.7	9.2	3/6	8.50E-07	12	4/6	Spread	No
PRPH	171-179	GLAEDLAAL	Conventional	4.0	3.8	1.1	0.5	3/6	1.08E-06	19	5/6	Spread	No
*PTPRN-021	392-402	SLAAGVKLLEI	mRNA splice	81.5	98.3	0.0	0.0	5/5	4.49E-06	17	5/5	Clustered	Yes
*PTPRN-021	398-407	KLLEILAEHV	mRNA splice	81.5	98.3	0.0	0.0	5/5	4.30E-06	13	5/5	Clustered	Yes
PTPRN-021	402-409	ILAEHVHM	mRNA splice	81.5	98.3	0.0	0.0	1/5	5.41E-07	9	4/5	Clustered	No
**PTPRN2-005	11-19	LLLLLPPRV	mRNA splice	33.3	10.8	0.0	0.0	4/6	1.67E-06		3/6	Clustered	Yes
*PTPRN2-005	19-27	VLPAAPSSV	mRNA splice	33.3	10.8	0.0	0.0	6/6	5.53E-06	8	6/6	Clustered	Yes
*REXO2-020	2-10	SVANALWIV	mRNA splice	4.2	10.5	0.0	0.0	5/6	1.94E-06	28	5/6	Clustered	Yes
**SCG5	186-195	YLQGQRLDNV	Conventional	252.5	163.2	3.2	2.9	5/6	1.16E-06	20	6/6	Clustered	Yes
**SCG5-009	186-194	FLSGAVNRL	mRNA splice	0.3	41.4	0.0	0.0	5/5	6.62E-06	7	5/5	Clustered	Yes
**SLC30A8-002	16-25	KMYAFTLESV	mRNA splice	43.1	38.1	0.3	0.0	4/6	1.33E-06	7	2/6	Clustered	Yes
ST18	304-312	SLLEQAIAL	Conventional	8.2	4.8	0.2	0.5	3/6	6.70E-07	33	3/6	Spread	No
**UCN3	1-9	MLMPVHFLL	Conventional	61.9	26.7	0.2	0.1	5/5	1.30E-05	11	5/5	Clustered	Yes
WARS-035	66-74	GLDEIDSAV	mRNA splice	0.3	31.8	0.0	0.2	0/6	3.36E-07	0	2/6	Spread	No

STAR METHODS

Contact for reagent and resource sharing

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Roberto Mallone (roberto.mallone@inserm.fr).

Experimental model and subject details

Cell lines

The ECN90 cell line (HLA-A*02:01/03:01, -B*40:01/49:01, -C*03:04/07:01) was derived from a human neonatal pancreas using described protocols (Ravassard et al., 2011). Cells were seeded in 15-cm diameter tissue culture dishes (Techno Plastic Products AG) coated with 0.1% fibronectin solution from human plasma (Sigma; 400 ng/cm²) and extracellular matrix from Engelbreth-Holm-Swarm murine sarcoma (Sigma; 1-2.4 mg/cm²). They were maintained in DMEM/F12 medium supplemented with 2% bovine serum albumin, 6.7 ng/ml sodium selenite, 10 mM nicotinamide, 50 μ M β -mercaptoethanol and penicillin/streptomycin. IFN- γ (R&D) was added to the cell culture at 80-90% confluence at a final concentration of 500 U/ml for 16-18 h. IFN- γ , TNF- α and IL-1 β (all from R&D) were added at a final concentration of 2,000 U/ml, 1,100 U/ml, and 1,000 U/ml, respectively.

Primary human tissues and PBMCs

For HLA-I peptidomics experiments, transplantation-grade, undispersed primary human islets (75% purity; HLA-A*02:01/25:01, -B*39:01/51:01, -C*12:03/14:02) were obtained from a brain-dead non-diabetic organ donor (age 49 years, male, BMI 37 kg/m²; protocol approved by the *Agence de la Biomédecine*) with standard procedures and maintained in CMRL 1066 medium (Sigma) supplemented with 10% fetal bovine serum. For RNAseq analyses, primary human islets from 5 brain-dead non-diabetic organ donors (mean age 50.6±10.2 years, 3 females, 2 males, BMI 25±2 kg/m²; 57±5% β cells; protocol approved by the Ethics

Committee of the University of Pisa, Italy) were exposed or not to IFN- γ (1,000 U/ml) and IL-1 β (50 U/ml) for 48 h. For HLA-I expression analyses (Fig. 2G), primary human islets were from 3 non-diabetic organ donors (mean age 54.7±14.3 years, 1 female, 2 males, BMI 29±5 kg/m²; protocol approved by the Ethics Committee of the University of Pisa, Italy). Primary human HLA Class II¹⁰ and Class II^{hi} mTECs were purified as described (Pinto et al., 2014) from the thymi of 3 children (male gender, age 6 days, 4 months and 9 months) undergoing corrective cardiac surgery at the University of Heidelberg, Germany (Ethics approval 367/2002). Blood was drawn into 9 ml sodium heparin tubes from T1D and healthy donors (Table S4) under the Ethics approval DC-2015-2536 Ile-de-France I. Informed consent was obtained from all subjects, or next-of-kin for islet donors. For *in-situ* MMr staining, pancreas and PLN sections were provided by nPOD.

Method details

Purification of pHLA-I complexes

W6/32 and HC10 anti-HLA-I mAbs were purified on a protein A Prosep Ultraplus column (Millipore) from hybridoma supernatants. The W6/32 mAb recognizes a conformational epitope formed by the interaction of the HLA-I heavy chain and β 2-microglobulin and was used for purifying pHLA-I complexes and for flow cytometry in conjunction with an FITC-conjugated goat anti-mouse Ig Ab (BD). The HC10 mAb recognizes a linear epitope on the HLA-I heavy chain and was used for Western blotting along with an anti- α -tubulin mAb for loading control (Fig. 1B). Since α -tubulin expression is higher in ECN90 β cells than in primary islets, an anti- β -actin mAb was used for loading control to compare HLA-I expression and upregulation between the two cell types (Fig. 2G). To this end, the membrane previously probed for HLA-I (45 kD) was stripped and reprobed for β -actin (42 kD).

The HLA-I peptidome of the ECN90 β -cell line was obtained from 5 biological replicates. A single biological replicate was available for primary human islets. Frozen cell pellets (~20x10⁶/condition for ECN90 cells; ~25,000 islet equivalents/condition for primary islets, corresponding to ~19x10⁶ β cells) were resuspended in a buffer containing 10 mM Tris-HCl pH 8.0, 150 mM NaCl, 5 mM EDTA, 0.1% (v/v) Complete Protease Inhibitor Cocktail (Roche), and 1% (w/v) octyl- β -D glucopyranoside (Sigma). Lysis was carried out at 4°C for 1 h under rotation, with two sonication steps at 30 and 60 min. Lysates were cleared by centrifugation and pHLA-I complexes immunoaffinity-purified with the W6/32 mAb covalently bound to Protein A Sepharose CL-4B beads (GE Healthcare) by dimethyl pimelimidate cross-linking. Beads were subsequently loaded on GELoader Tips (20 µl; ThermoFisher) and washed before elution of pHLA-I complexes with 10% acetic acid. Aliquots were collected at each washing and elution step for analysis by 12% SDS-PAGE and Western blot using the HC10 mAb to verify the yield and purity of the eluted HLA-I.

Eluted peptides and the associated HLA-I heavy chain and β 2-microglobulin obtained from $20x10^6$ cells were concentrated to 20 µl by vacuum centrifugation, acidified with 10 µl of 1% formic acid (Normapur) and loaded on C18 stage tips (ThermoFisher) prewashed with 100% methanol and equilibrated with 2% acetonitrile (ACN) in 0.1% formic acid in LC-MS grade water. After loading, the C18 stage tips were washed with 2% ACN/0.1% formic acid and peptides separated from the HLA-I heavy chain and β 2-microglobulin species by eluting them with 50% ACN/0.1% formic acid. The ACN was evaporated by vacuum centrifugation and the peptides resuspended up to 6 µl of volume in a solution of 2% ACN/0.1% formic acid and spiked with 10 fmol/µl of a cytomegalovirus pp65 495-503 peptide (NLVPMVATV) as internal control. For LC-MS analysis, 5 µl of this peptide solution were used.

LC-MS/MS

Peptides were loaded and separated by a nanoflow HPLC (RSLC Ultimate 3000, Dionex) on a C18 Acclaim PepMap nanocolumn (50 cm length, 75 µm internal diameter; Dionex) coupled on-line to a nano-electrospray ionization Q Exactive HF mass spectrometer (ThermoFisher) with a glass emitter (New Objective). Peptides were eluted with a linear gradient of 2-50% buffer B (80% ACN, 0.05% formic acid) at a flow rate of 220 nl/min over 60 min at 35°C. Data was acquired using a data-dependent acquisition "Top 20" method. We acquired one full-scan MS spectrum at a resolution of 60,000 at 200 m/z with an automatic gain control (AGC) target value of 3x10⁶ ions, followed by 10 MS/MS spectra in higher energy collisional dissociation mode on the 10 most intense ions at a resolution of 15,000 at 200 m/z with an AGC target value of 1×10^5 with a maximum injection time of 120 ms and a dynamic exclusion of 20 s. Unassigned precursor ion charge states and charge states >4 were excluded. The peptide match option was set to 'preferred'. The resulting spectra were analyzed by MaxQuant (www.coxdocs.org) using a custom database comprising: a) the reference human proteome (Swiss-Prot/UniProt, up000005640, release December 2012); b) an in-house database containing 119,305 predicted peptide splice products (Berkers et al., 2015) from major known and candidate β -cell protein Ags (Fig. S1 and Data S1); and c) the predicted aa neo-sequences encoded by mRNA splice variants identified by RNASeq. The following parameters were set: enzyme specificity: unspecific; variable modifications: methionine, tryptophan and histidine oxidation (+15.99 Da), cysteine oxidation to cysteic acid (+47.98 Da) and tryptophan conversion to kynurenine (+3.99 Da); maximum false discovery rate 5%. Since the MS identification was targeted on HLA-I-eluted peptides rather than on proteins, the protein false discovery rate parameter was set to 100%. The initial allowed mass deviation of the precursor ion was set to 10 ppm and the maximum fragment mass deviation was set to 20 mDa. The "match between runs" option was enabled to match identifications across

different replicates in a time window of 0.5 min and an initial alignment time window of 20 min.

RNAseq

RNAs from 5 individual preparations of primary human islets exposed or not to IFN- γ and IL-1 β for 48 h and from HLA Class II¹⁰ and Class II^{hi} human mTECs were sequenced on an Illumina HiSeq 2000 at high depth (coverage >150x10⁶ reads, which is sufficient to detect >80% of slice variants). RNA sequencing reads were mapped to the human reference genome hg19 using TopHat 2 and the Gencode release 18 annotation dataset. Mapped reads were used to quantify abundance and analyse the differential expression of genes and transcripts using Flux Capacitor (Montgomery et al., 2010).

HLA-I peptidomics and transcriptomics bio-informatics analysis

Predicted HLA-I binding affinities for each nonamer peptide were visualized as sequence logo plots using Gibbs clustering (www.cbs.dtu.dk/services/GibbsCluster; Fig. 1D), and as heat maps (Fig. 1E) using the open-source script of Caron et al. (Caron et al., 2015).

For conventional peptides, source proteins were selected with Perseus (www.coxdocs.org) based on: *a*) a non-ubiquitous expression pattern, based on the Human Protein Atlas (www.proteinatlas.org); *b*) a pancreas- and β -cell-enriched expression pattern, based on the Human Protein Atlas, the Human Protein Reference Database (www.hprd.org) and the Single-Cell Gene Expression Atlas of Human Pancreatic Islets (http://sandberg.cmb.ki.se/pancreas). The bioinformatics analysis pipeline is detailed in Fig. 1G. The 3,544 peptides identified were first filtered based on inter-sample reproducibility (≥ 2 of 5 biological replicates; n=2,997, 85%) and aa length (8-12 aa; n=2,795, 93%). For all peptides, the final filter was based on an enrichment in HLA-I-purified samples compared with mock-purified ones based on *m*/*z* peak intensity, which verified the specific association of the identified peptides with pHLA-I complexes. From this dataset, we identified:

1) 217 (8%) peptides derived from ubiquitous proteins carrying PTMs (blue pipeline in Fig.1G), from which 99 (46%; Table S2) were enriched in HLA-I-purified samples.

2) 15 (0.5%) peptide splice variants (brown pipeline in Fig. 1G; see Fig. S1 for the identification strategy), from which 10 (67%) were enriched in HLA-I-purified samples (listed in Table S1).

3) The remaining 2,561 peptides (both conventional and PTM species derived from β -cellenriched proteins; red pipeline in Fig. 1G) were filtered by Perseus based on non-ubiquitous expression of their source proteins (n=411; 16%), using the Human Protein Atlas; and pancreas- and β -cell-enriched expression (n=139, 34%), using the Human Protein Atlas, the Human Protein Reference Database and the Single-Cell Gene Expression Atlas of Human Pancreatic Islets. Finally, the 86 (62%) remaining peptides were retained as enriched in HLA-I-purified samples (listed in Table S1).

4) mRNA splice variants (green pipeline in Fig. 1G; n=2, 0.1%; both enriched in HLApurified samples; listed in Table S1) were identified by analyzing the HLA-I peptidomics dataset against the predicted aa neo-sequences obtained from RNAseq analysis. For this RNAseq pipeline (dashed boxes in Fig. 1G), 53,280 mRNAs were filtered based on:

a) A median RPKM>5 in islets, either under basal or inflammatory conditions, a cut-off selected based on the median RPKM of known islet Ags (islet expression filter; n=14,504, 27%).

b) A median RPKM<0.1 in mTECs (either HLA Class II^{lo} or Class II^{hi}), or a median RPKM fold-decrease>100 *vs.* islet (mTEC expression filter; n=908, 6%).

c) A median RPKM fold-increase>10 in islets compared to 12 control tissues (adipose tissue, breast, colon, heart, kidney, liver, lung, lymph node, ovary, prostate, skeletal muscle, white blood cells), using the Illumina BodyMap 2.0 dataset (islet enrichment filter; n=166, 18%).

Tissues of neuroendocrine origin (brain, testis, adrenal gland and thyroid) were excluded for this filtering.

d) We subsequently focused our analysis on mRNA isoforms, as described (Eizirik et al., 2012; Villate et al., 2014). The predicted translation products were aligned using MUSCLE 3.8 (www.ebi.ac.uk/Tools/msa/muscle), and aa neo-sequences were defined by comparing the predicted aa sequence of each mRNA isoform with that of the reference mRNA, taking as reference the longest and/or most prevalent mRNA isoform in islets (neo-sequence generation filter; n=88, 53%).

The 336 predicted aa neo-sequences obtained from these 88 mRNA variants were used to interrogate HLA-I peptidomics datasets and searched in parallel for:

e) Potential HLA-A2 binders, based on their predicted HLA-A2 binding affinity (K_D <100 nM by NetMHC 4.0; www.cbs.dtu.dk/services/NetMHC) and stability (half-life \geq 1.5 h by NetMHCstab 1.0; www.cbs.dtu.dk/services/NetMHCstab-1.0) (predicted HLA-A2 binding filter; n=66, 20%).

f) A 9-10-aa length (peptide length filter; n=43, 65%).

g) A neo-sequence ≥ 3 aa (neo-sequence filter; n=39, 92%).

Overall, this combined HLA-I peptidomics and transcriptomics analysis pipeline led to the identification of 99 *in vitro* PTM candidates from ubiquitous proteins (Table S2); 10 *in vitro* peptide splice candidates, 86 *in vitro* candidates from β -cell proteins (including PTM variants) and 2 *in vitro* mRNA splice candidates (Table S1); and 39 *in silico* mRNA splice candidates (Fig. S2I). PTM candidates were not further analyzed. The other candidates underwent additional filtering steps to focus subsequent studies on HLA-A2-restricted peptides. Predicted HLA-A2 binders were first selected *in silico* using NetMHC 4.0 and NetMHCstab 1.0 (except for *in silico* mRNA splice candidates, which had already been filtered for

predicted HLA-A2 binding in the same way), then *in vitro* for experimental HLA-A2 binding and, finally, for CD8⁺ T-cell recognition.

HLA-A2 binding assays

Peptide binding to HLA-A*02:01 was measured using the transporter associated with Ag processing (TAP)-deficient T2 cell line. TAP deficiency leads to defective translocation of endogenous peptides from the cytosol into the ER, leading to the expression of unstable, empty HLA-A*02:01 molecules on the cell surface. This expression is stabilized in the presence of HLA-A*02:01-binding peptides, resulting in a higher surface expression that can be monitored by flow cytometry. T2 cells were washed in RPMI medium and plated in roundbottom 96-well plates at 0.2×10^6 cells/200 µl in the presence of 5 µg/ml β2-microglobulin. Peptides prepared in DMSO were sequentially diluted 4-fold in RPMI and added to final concentrations of 102.4 to 0.1 µM for 24 h at 37°C, 5% CO₂. The HLA-A*02:01-binding peptide Flu MP₅₈₋₆₆ (GILGFVFTL), and a non-binding peptide CHGA₃₈₂₋₃₉₀ (HPVGEADYF) were included as positive and negative controls, respectively. After incubation, the cells were washed twice with ice-cold phosphate-buffered saline (PBS) and stained with the viability marker Live/Dead Red and mouse anti-HLA-A2 mAb BB7.2, followed by an Alexa Fluor 488-labeled goat-anti-mouse IgG Ab (Interchim). Following acquisition on a BD Fortessa cytometer, results were analyzed by gating on viable cells and expressed as the median fluorescence intensity fold increase of the test peptide compared with the negative control peptide at the same concentration.

HLA-A2 MMr assays

All peptides were synthesized at >90% purity (Synpeptides). HLA-A2 MMrs were produced and used as described (Culina et al., 2018). Each pHLA-A2 complex was used at a final concentration of 8-27 nM and conjugated with fluorochrome-labeled streptavidin at a 1:4 ratio. The combinatorial MMr panel was first set up by staining HLA-A2⁺ PBMCs with the same set of fluorescent streptavidin-labeled MMrs, all loaded with the Flu MP₅₈₋₆₆ epitope. Compensations were set using fluorescence-minus-one samples (i.e. omitting one streptavidin at a time). The concentration of each fluorescent MMr was corrected for the variable staining index of each streptavidin, in order to obtain a distinct double-MMr⁺ population for each fluorochrome pair. The identification of the same MMr⁺ population by each pair validated the panel. PBMCs were isolated by density gradient centrifugation using 50 ml Leucosep® tubes (Greiner/Dominique Dutscher), washed twice in RPMI medium supplemented with AB human serum (Sigma), counted on a ThermoFisher Countess II automated counter and frozen in pre-chilled 10% DMSO solution in AIM-V medium (ThermoFisher) using CoolCell containers (Biocision) stored overnight at -80°C prior to transfer into liquid nitrogen. At thawing, PBMCs were immediately diluted in pre-warmed AIM-V medium. Following centrifugation and one additional wash in AIM-V, PBMCs were counted and rested in the presence of 50 nM dasatinib for 30 min at 37°C before magnetic depletion of CD8⁻ cells (StemCell Technologies). Staining was performed for 20 min at 20°C in 20 µl PBS-dasatinib for 10⁷ cells with the combinatorial double-coded MMr panels (Culina et al., 2018) detailed in Fig. 3, followed, without washing, by mAb and Live/Dead Aqua staining at 4°C for 20 min. After one wash, cells were acquired using a FACSAria III cytometer configured as detailed in Table S6. Candidate epitopes binding to HLA-A2 (Fig. S2) that did not yield any appreciable MMr staining provided negative controls for each panel. Data was analyzed with FlowJo software as described in Fig. 3. Cells were sequentially gated on small lymphocytes, singlets, live cells (Live/Dead Aqua⁻), CD3⁺CD8⁺ T cells and total PE⁺, PE-CF594⁺, APC⁺, BV650⁺, BV711⁺ and BV786⁺ MMr⁺ T cells. Using Boolean operators, these latter gates allowed to selectively visualize each double-MMr⁺ population by including only those events positive for the corresponding fluorochrome pair. For example, UCN3₁₋₉ MMr⁺ cells (PE⁺PE-CF594⁺) were visualized by gating on events that were PE+PE-CF594+APC-BV650-BV711-BV786-.

Events negative for all MMr fluorochromes (PE⁻PE-CF594⁻APC⁻BV650⁻BV711⁻BV786⁻) were represented in the same PE/PE-CF-594 dot plot to set the double-MMr⁺ gate, as shown in Fig. 4A-F. CD45RA and CCR7 staining was subsequently visualized by gating on these MMr⁺ cells. Each dot plot of Fig. 3B-C displays a color-coded overlay of each double-MMr⁺ fraction and of the MMr⁻ population to visualize the separation of each epitope-reactive CD8⁺ T-cell fraction relative to the others.

In-situ HLA-A2 MMr staining

In-situ immunohistochemistry staining was performed as described (Culina et al., 2018). Unfixed, frozen sections were dried for 2 h, loaded with 1 μ g of PE-labeled MMrs overnight at 4°C, washed gently with PBS and fixed in 2% paraformaldehyde for 10 min. After a further wash, endogenous peroxidase activity was blocked with 0.3% H₂O₂. Sections were then incubated serially with a rabbit anti-PE Ab (Abcam), horseradish peroxidase-conjugated swine anti-rabbit Ig (Dako) and 3,3'-diaminobenzidine tetrahydrochloride substrate (ThermoFisher). After a final wash, sections were counterstained with hematoxylin, dehydrated via sequential passages in 95-100% ethanol and xylene, mounted and analyzed using a Nikon Eclipse Ni microscope with NIS-Elements D software v4.40.

In-situ immunofluorescence staining was performed similarly, but non-specific reactions were blocked with 5% goat serum for 2 h at room temperature before serial incubations with rabbit anti-PE Ab (1:250, 1.5 h at room temperature) and Alexa Fluor 594-conjugated goat anti-rabbit IgG (ThermoFisher; 1:500, 1 h at room temperature). After a further wash, sections were incubated for 1 h at room temperature with rat anti-CD8 mAb (Abcam; 1:100) together with mouse anti-CD45RO mAb (BioLegend; 1:200) followed, after one wash, by one final incubation for 1 h at room temperature with Alexa Fluor 488-conjugated goat anti-rat IgG together with Alexa Fluor 647-conjugated goat anti-mouse IgG (1:500/each; both from

ThermoFisher). After DNA counterstaining with DAPI, sections were mounted and analyzed using a Leica TCS SP5 confocal laser scanning microscope with LAS software v2.6.0.7266.

Quantification and statistical analysis

Statistical details of experiments can be found in the legends of each figure. A two-tailed p<0.05 cut-off was used to define statistical significance.

Data and software availability

The custom script used to predict peptide splice products is provided in Data S1. The islet RNAseq datasets have been deposited under GEO accession number GSE108413. The mTEC RNAseq dataset is provided in Data S2.

SUPPLEMENTAL DATA

Data S1. Related to Fig. 1G-S1. In-house script used to predict peptide splice products.

Data S2. Related to Fig. 1G. mTEC RNAseq dataset of mRNA isoforms expressed in human islets.













KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse monoclonal anti-HLA Class I, clone W6/32	ATCC	ATCC HB-95; RRID:
		CVCL_7872
Mouse monoclonal anti-HLA Class I, clone HC10	Laboratory of	RRID:
	Hidde L. Ploegh	AB_2728622
Mouse monoclonal anti-α-tubulin, clone DM1A	Thermo Fisher	Cat#14-4502-80;
	eBioscience	RRID: AB_1210457
Mouse monoclonal anti-β-actin, clone AC-15	Sigma	Cat#A5441; RRID: AB_476744
Goat polyclonal Anti-mouse Ig (H+L) HRP	SouthernBiotech	Cat#1010-05; RRID: AB_2728714
Mouse monoclonal anti-HLA-A2, clone BB7.2	ATCC	ATCC HB-82; RRID: CVCL 7246
Goat polyclonal anti-mouse Ig FITC	BD Pharmingen	Cat#554001; RRID:
Goat polyclopal anti-mouse InG (H+L) Alexa Fluor 488	Interchim	Cat#EP-SA/010
Guar polycional anti-mouse igo (II+L) Alexa i idoi 400	Interchini	RRID: ΔR 2728715
EasySen™ Human CD8+ T Cell Enrichment Kit	StemCell	Cat#19053: RRID:
	Technologies	AB 2728716
Mouse monoclonal anti-human CD3 APC-H7, clone SK7	BD Pharmingen	Cat#560176: RRID:
		AB 1645475
Mouse monoclonal anti-human CD8 PE-Cy7, clone	BD Pharmingen	Cat#557746; RRID:
RPA-T8		AB_396852
Mouse monoclonal anti-human CD45RA FITC, clone	BD Pharmingen	Cat#555488; RRID:
HI100		AB_395879
Mouse monoclonal anti-human CCR7 BV421, clone 150503	BD Horizon	Cat#562555; RRID: AB_2728119
LIVE/DEAD™ Fixable Aqua Dead Cell Stain Kit	Thermo Fisher	Cat#L34965
LIVE/DEAD™ Fixable Red Dead Cell Stain Kit	Thermo Fisher	Cat# L23102
Streptavidin PE	BD Pharmingen	Cat#554061; RRID:
		AB_10053328
Streptavidin PE-CF594	BD Horizon	Cat#562284; RRID:
		AB_11154598
Streptavidin APC	BD Pharmingen	Cat#554067; RRID:
		AB_10050396
Streptavidin BV650	BD Horizon	Cat#563855
Streptavidin BV711	BD Horizon	Cat#563262
Streptavidin BV786	BD Horizon	Cat#563858
Rabbit polyclonal anti-B phycoerythrin	Abcam	Cat#ab7011; RRID:
		AB_305700
Swine polyclonal anti-rabbit Ig HRP	Dako	Cat#P0217; RRID:
		AB_2728719
Rat monoclonal anti-human CD8, clone YTC182.20	Abcam	Cat#ab60076; RRID: AB 940921
Mouse monoclonal anti-human CD45RO, clone UCHL1	BioLegend	Cat#304202; RRID:
	-	AB_314418
Goat polyclonal anti-rabbit IgG (H+L) Alexa Fluor 594	Thermo Fisher	Cat#A-11037; RRID:
		AB_2534095
Goat polyclonal anti-rat IgG (H+L) Alexa Fluor 488	Thermo Fisher	Cat# A-11006;
		RRID: AB 2534074

Goat polyclonal anti-mouse IgG (H+L) Alexa Fluor 647	Thermo Fisher	Cat# A-21236; RRID: AB_2535805
Bacterial and Virus Strains		
Biological Samples		
Human pancreas unfixed frozen sections	nPOD	www.jdrfnpod.org
Human pancreatic lymph node unfixed frozen sections	nPOD	www.jdrfnpod.org
Chemicals, Peptides, and Recombinant Proteins		
HLA-A2 heavy chain	ImmunAware	N/A
β2-microglobulin	Lee Biosolutions	Cat#126-11
Dasatinib	LC Laboratories	Cat#D-3307
Critical Commercial Assays		
Deposited Data		
Human reference proteome Swiss-Prot/UniProt,	Uniprot	http://www.uniprot.or
up000005640, release December 2012		g/proteomes/UP000
Liver on Drotein Atlance 40.0	Liveran Dratain Atlan	005640
Human Protein Atlas V.13.0	Human Protein Atlas	www.proteinatias.org
Human Protein Reference Database, release 9	Reference Database	www.npra.org
Single-Cell Gene Expression Atlas of Human Pancreatic	Sandberg Lab	http://sandberg.cmb.
Islets		ki.se/pancreas
Gencode release 18	The GENCODE	https://www.gencode
	Project	genes.org
liiumina Bodymap 2.0	http://www.ncbi.nim.ni	GEO #GSE30611
	MTAB-513	
Human islet RNAseq	This paper	GEO # GSE108413
Human mTEC RNAseq	This paper	Data S2
Experimental Models: Cell Lines		
Human: ECN90 β-cell line	Culina et al., 2018;	N/A
	Univercell Biosolutions	
Human: 12 (174 x CEM.12) cell line	AICC	ATCC CRL-1992;
Mouse: W6/32 hybridoma cell line	ATCC	ATCC HB-95' RRID'
	/1100	CVCL 7872
Mouse: HC10 hybridoma cell line	Laboratory of	RRID:
	Hidde L. Ploegh	AB_2728622
Mouse: BB7.2 hybridoma cell line	ATCC	ATCC HB-82; RRID: CVCL 7246
Experimental Models: Organisms/Strains		—
Oligonucleotides		
Recombinant DNA		
Software and Algorithms	1	1
FlowJo v10	FlowJo, LLC	www.flowjo.com

Flux Capacitor v1.6.1 Montgomery et al., 2010 http://sammeth.net/c onfluence/display/FL UX/Home Gibbs clustering v1.1 DTU Bioinformatics www.cbs.dtu.dk/serv icces/GibbsCluster LAS v2.6.0.7266 Leica Microsystems N/A MaxQuant v1.5.3.8 Max Planck Institute of Biochemistry http://www.coxdocs. org MUSCLE v3.8 EMBL-EBI www.ebi.ac.uk/Tools /msa/muscle NetMHC v4.0 DTU Bioinformatics www.cbs.dtu.dk/serv icces/NetMHC NetMHCStab v1.0 DTU Bioinformatics www.cbs.dtu.dk/serv icces/NetMHC NetMHCStab v1.0 DTU Bioinformatics www.cbs.dtu.dk/serv icces/NetMHCStab-1.0 NIS-Elements D v4.40 Nikon N/A Perseus v1.5.2.4 Max Planck Institute of Biochemistry http://www.coxdocs. org/articles/07661/fig ures#SD9-data Python and R Scripts Caron et al., 2015 http://esciences. org/articles/07661/fig ures#SD9-data TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html			
2010Ontildence/display/FL UX/HomeGibbs clustering v1.1DTU BioinformaticsWww.cbs.dtu.dk/serv ices/GibbsClusterLAS v2.6.0.7266Leica MicrosystemsN/AMaxQuant v1.5.3.8Max Planck Institute of Biochemistryhttp://www.coxdocs. orgMUSCLE v3.8EMBL-EBIwww.ebi.ac.uk/Tools /msa/muscleNetMHC v4.0DTU Bioinformaticswww.ebs.dtu.dk/serv ices/NetMHCNetMHC stab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCNetMHCStab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstab- 1.0NIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataTopHat v2.0.10J. Hopkins University Center for Computational BiologyData S1OtherMaxNikonhttp://ccb.jhu.edu/sof tware/tophat/index.s html	Flux Capacitor v1.6.1	Montgomery et al.,	http://sammeth.net/c
Gibbs clustering v1.1DTU BioinformaticsDXHomeGibbs clustering v1.1DTU Bioinformaticswww.cbs.dtu.dk/serv ices/GibbsClusterLAS v2.6.0.7266Leica MicrosystemsN/AMaxQuant v1.5.3.8Max Planck Institute of Biochemistryhttp://www.coxdocs. orgMUSCLE v3.8MaxPlanck Institute of Biochemistryhttp://www.cbs.dtu.dk/serv ices/NetMHCNetMHC v4.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCNetMHCStab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstabpan nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataTopHat v2.0.10J. Hopkins University Center for Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html		2010	onfluence/display/FL
Gibbs clustering v1.1DTU Bioinformaticswww.cbs.dtu.dk/serv ices/GibbsClusterLAS v2.6.0.7266Leica MicrosystemsN/AMaxQuant v1.5.3.8Max Planck Institute of Biochemistryhttp://www.coxdocs. orgMUSCLE v3.8EMBL-EBIwww.ebi.ac.uk/Tools /msa/muscleNetMHC v4.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCNetMHCStab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCStab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCStabpan nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataTopHat v2.0.10J. Hopkins University Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html			UX/Home
LAS v2.6.0.7266Leica MicrosystemsN/AMaxQuant v1.5.3.8Max Planck Institute of Biochemistryhttp://www.coxdocs. orgMUSCLE v3.8EMBL-EBIwww.ebi.ac.uk/Tools /msa/muscleNetMHC v4.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCNetMHCstab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCStab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCStabpan nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015http://cb.jhu.edu/sof tware/tophat/index.s computational BiologyOtherJ. Hopkins University Center for Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html	Gibbs clustering v1.1	DIU Bioinformatics	www.cbs.dtu.dk/serv
LAS v2.6.0.7266 Leica Microsystems N/A MaxQuant v1.5.3.8 Max Planck Institute of Biochemistry http://www.coxdocs. org MUSCLE v3.8 EMBL-EBI www.ebi.ac.uk/Tools /msa/muscle NetMHC v4.0 DTU Bioinformatics www.cbs.dtu.dk/serv ices/NetMHC NetMHCStab v1.0 DTU Bioinformatics www.cbs.dtu.dk/serv ices/NetMHCstab- 1.0 NetMHCStabPan v1.0 DTU Bioinformatics www.cbs.dtu.dk/serv ices/NetMHCstabpa n NIS-Elements D v4.40 Nikon N/A Perseus v1.5.2.4 Max Planck Institute of Biochemistry http://www.coxdocs. org Python and R Scripts Caron et al., 2015 http://elifesciences. org/articles/07661/fig ures#SD9-data TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html Other Uter Iter/org http://ccb.jhu.edu/sof			ices/GibbsCluster
MaxQuant v1.5.3.8 Max Planck Institute of Biochemistry http://www.coxdocs. org MUSCLE v3.8 EMBL-EBI www.ebi.ac.uk/Tools /msa/muscle NetMHC v4.0 DTU Bioinformatics www.cbs.dtu.dk/serv ices/NetMHC NetMHCstab v1.0 DTU Bioinformatics www.cbs.dtu.dk/serv ices/NetMHCstab- 1.0 NetMHCStabPan v1.0 DTU Bioinformatics www.cbs.dtu.dk/serv ices/NetMHCstabpan 1.0 NIS-Elements D v4.40 Nikon N/A Perseus v1.5.2.4 Max Planck Institute of Biochemistry http://www.coxdocs. org Python and R Scripts Caron et al., 2015 https://elifesciences. org/articles/07661/fig ures#SD9-data TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s httml	LAS v2.6.0.7266	Leica Microsystems	N/A
BiochemistryorgMUSCLE v3.8EMBL-EBIwww.ebi.ac.uk/Tools /msa/muscleNetMHC v4.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCNetMHCstab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCStab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCStabpan nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataTopHat v2.0.10J. Hopkins University Center for Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s httml	MaxQuant v1.5.3.8	Max Planck Institute of	http://www.coxdocs.
MUSCLE v3.8EMBL-EBIwww.ebi.ac.uk/Tools /msa/muscleNetMHC v4.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCNetMHCstab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCStab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCStabpan nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015http://elifesciences. org/articles/07661/fig ures#SD9-dataTopHat v2.0.10J. Hopkins University Center for computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html		Biochemistry	org
NetMHC v4.0DTU Bioinformatics/msa/muscleNetMHC stab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCNetMHCStab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstabpa nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataTopHat v2.0.10J. Hopkins University Center for Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s 	MUSCLE v3.8	EMBL-EBI	www.ebi.ac.uk/Tools
NetMHC v4.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCNetMHCstab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstabpa nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015http://elifesciences. org/articles/07661/fig ures#SD9-dataTopHat v2.0.10J. Hopkins University Center for Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html			/msa/muscle
NetMHCstab v1.0DTU Bioinformaticsices/NetMHCNetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstabpa nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataTopHat v2.0.10J. Hopkins University Center for Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html	NetMHC v4.0	DTU Bioinformatics	www.cbs.dtu.dk/serv
NetMHCstab v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstab- 1.0NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstabpa nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015http://clifesciences. org/articles/07661/fig ures#SD9-dataTopHat v2.0.10J. Hopkins University Center for Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html			ices/NetMHC
NetMHCStabPan v1.0DTU Bioinformaticsices/NetMHCstab- 1.0NIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataScript for the prediction of peptide splice productsThis paperData S1TopHat v2.0.10J. Hopkins University Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html	NetMHCstab v1.0	DTU Bioinformatics	www.cbs.dtu.dk/serv
NetMHCStabPan v1.0DTU Bioinformatics1.0NIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataScript for the prediction of peptide splice productsThis paperData S1TopHat v2.0.10J. Hopkins University Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html			ices/NetMHCstab-
NetMHCStabPan v1.0DTU Bioinformaticswww.cbs.dtu.dk/serv ices/NetMHCstabpa nNIS-Elements D v4.40NikonN/APerseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataScript for the prediction of peptide splice productsThis paperData S1TopHat v2.0.10J. Hopkins University Center for Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html			1.0
NIS-Elements D v4.40 Nikon N/A Perseus v1.5.2.4 Max Planck Institute of Biochemistry http://www.coxdocs. org Python and R Scripts Caron et al., 2015 https://elifesciences. org/articles/07661/fig ures#SD9-data Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html	NetMHCStabPan v1 0	DTU Bioinformatics	www.cbs.dtu.dk/serv
NIS-Elements D v4.40 Nikon N/A Perseus v1.5.2.4 Max Planck Institute of Biochemistry http://www.coxdocs.org Python and R Scripts Caron et al., 2015 https://elifesciences.org/articles/07661/figures#SD9-data Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html Other Image: Script Scr		Die Diemeinatie	ices/NetMHCstabpa
NIS-Elements D v4.40 Nikon N/A Perseus v1.5.2.4 Max Planck Institute of Biochemistry http://www.coxdocs.org Python and R Scripts Caron et al., 2015 https://elifesciences.org/articles/07661/fig ures#SD9-data Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html Other Image: Script Sc			n
Nite Lientering D V1.46 Niterin Niterin Niterin Perseus v1.5.2.4 Max Planck Institute of Biochemistry http://www.coxdocs.org Python and R Scripts Caron et al., 2015 https://elifesciences.org/articles/07661/figures#SD9-data Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html Other Image: Script	NIS-Elements D v4 40	Nikon	N/A
Perseus v1.5.2.4Max Planck Institute of Biochemistryhttp://www.coxdocs. orgPython and R ScriptsCaron et al., 2015https://elifesciences. org/articles/07661/fig ures#SD9-dataScript for the prediction of peptide splice productsThis paperData S1TopHat v2.0.10J. Hopkins University Center for Computational Biologyhttp://ccb.jhu.edu/sof tware/tophat/index.s html		T T T T T T T T T T T T T T T T T T T	
Perseus v1.5.2.4 Max Planck Institute of Biochemistry http://www.coxdocs. org Python and R Scripts Caron et al., 2015 https://elifesciences. org/articles/07661/fig ures#SD9-data Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html			
Biochemistry org Python and R Scripts Caron et al., 2015 https://elifesciences. org/articles/07661/fig ures#SD9-data Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html Other Image: Computational Biology Image: Computational Biology	Perseus v1.5.2.4	Max Planck Institute of	http://www.coxdocs.
Python and R Scripts Caron et al., 2015 https://elifesciences.org/articles/07661/fig ures#SD9-data Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.shtml Other Image: Computational Biology Image: Computational Biology		Biochemistry	org
org/articles/07661/fig ures#SD9-data Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html Other Image: Computational Biology html	Python and R Scripts	Caron et al., 2015	https://elifesciences.
Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s Other Image: Computational Biology http://ccb.jhu.edu/sof			org/articles/07661/fig
Script for the prediction of peptide splice products This paper Data S1 TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html Other Image: Computational Biology http://ccb.jhu.edu/sof			ures#SD9-data
TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s Other J. Hopkins University Center for http://ccb.jhu.edu/sof	Script for the prediction of peptide splice products	This paper	Data S1
TopHat v2.0.10 J. Hopkins University Center for Computational Biology http://ccb.jhu.edu/sof tware/tophat/index.s html Other Image: Computational Biology http://ccb.jhu.edu/sof			
Center for tware/tophat/index.s Computational Biology html	TopHat v2.0.10	J. Hopkins University	http://ccb.jhu.edu/sof
Other Computational Biology html		Center for	tware/tophat/index.s
Other		Computational Biology	html
	Other		

SUPPLEMENTAL ITEMS

Figure S1. Related to Fig. 1G. Schematic of the strategy used for predicting peptide splice variants.

Figure S2. Related to Fig. 4. HLA-A2 binding measurements for the β -cell peptides selected for CD8⁺ T-cell validation.

Fig. S3. Related to Fig. 6. Screening of HLA-A2-restricted peptide reactivities in pancreasinfiltrating cells from a T1D case.

Table S1. Related to Fig 1G, 2A-E. List of the 98 HLA-I-bound peptides identified in ECN90 β cells.

Table S2. Related to Fig 1G. List of the 99 HLA-I-bound PTM peptides originating from ubiquitous proteins identified in ECN90 β cells.

Table S3. Related to Fig 2F. List of the 33 HLA-I-bound peptides identified in primary human islets.

Table S4. Related to Fig. 4-5. Characteristics of HLA-A2⁺ study subjects for ex-vivo MMr studies on PBMCs.

Table S5. Related to Fig. 6. nPOD cases analyzed by *in-situ* tissue MMr staining.

Table S6. Related to STAR Methods. Configuration of the flow cytometer used for HLA-A2MMr assays.

Data S1. Related to Fig. 1G-S1. In-house script used to predict peptide splice products.

Data S2. Related to Fig. 1G. mTEC RNAseq dataset of mRNA isoforms expressed in human islets.



Figure S1. Related to Fig. 1G. Schematic of the strategy used for predicting peptide splice variants. The indicated known or putative β -cell Ags (first column; PPI, IGRP, CHGA, SCG5, PTPRN/IA-2, NPY, SLC30A8/ZnT8, IAPP, PCSK2, GAD2/GAD65) were scanned with an in-house Python script (Data S1) based on reported peptide splicing preferences (Berkers et al., 2015) for the presence of the indicated N-and C-terminal preferred precursor sequences (second and third column, respectively, where Φ indicates any hydrophobic aa residue). Each N-terminal precursor sequence identified was then combined with each C-terminal precursor sequence, derived either from the same protein (*cis*-splicing) or from a different protein (*trans*-splicing). An example is shown for PPI and IGRP. The 119,305 predicted peptide splice sequences thus obtained (26,396 *cis*-spliced and 92,909 *trans*-spliced) were saved in FASTA format and appended to the Swiss-Prot human proteome database used for assigning aa sequences to the identified MS spectra with the MaxQuant algorithm.



Figure S2. Related to Fig. 4. HLA-A2 binding measurements for the β-cell peptides selected for CD8⁺ T-cell validation. (A-F) Flow cytometry measurement of peptide binding to HLA-A2 using TAP-deficient T2 cells. T2 cells were pulsed with the indicated peptides at 4-fold sequential dilutions (102.4 to 0.1 μ M) at 37°C and stained for HLA-A2. Representative stains at 0.4, 6.4 and 102.4 µM are shown for peptides that are weak binders (IAPP15-17/5-10; A), intermediate binders (SCG5-009186-194; B), strong binders (PCSK230-38; C) and non-binders (GAD2257-265; D), along with the non-binding (ChgA382-390; E) and binding (Flu MP58-66; F) control peptides included in each assay. G. Plotting of relative median fluorescence intensity values (arbitrary units, AU) for all peptide dilutions normalized to the non-binding ChgA₃₈₂₋₃₉₀ control peptide. Representative examples of two independent experiments are shown. (H-I) Summary of the HLA-A2 binding values measured for the β -cell peptides selected from the *in vitro* HLA-I peptidomics (H) and *in* silico RNAseq pipeline (I) using the T2 assays depicted in panels A-G. The relative median fluorescence intensity measured at 25.6 and 102.4 µM peptide concentrations is shown along with their relative ranking: - (<1.50 AU); + (1.50-1.99 AU); ++ (2-2.49 AU); +++ (\geq 2.50 AU). The *in silico* predicted HLA-A2 binding affinity (NetMHC 4.0) and stability values (NetMHCstab 1.0) are also shown. Peptides in bold red fonts bound HLA-A2 and were retained for CD8⁺ T-cell validation, with those eventually validated (see Fig. 4) highlighted in grey. In panel H, the GAD2₂₅₇₋₂₆₅ peptide eluted from ECN90 cells with a predicted HLA-A3 restriction is included as negative control. For β-cell peptides selected from the *in silico* RNAseq pipeline (panel I), RPKM values of the source mRNA splice variant in control- and cytokine-treated islets are shown in the last two columns. The color codes indicate no change in mRNA expression (<2 log₂ foldchange, either positive or negative; green), an increase for cytokine-treated islets (\geq +2 log₂ fold-change; red) or a decrease for cytokine-treated islets ($\leq -2 \log_2$ fold-change; blue).



Fig. S3. Related to Fig. 6. Screening of HLA-A2-restricted peptide reactivities in pancreasinfiltrating cells from a T1D case. Pancreas sections from T1D EUnPOD case #060217 (39-year-old female, T1D duration 21 years, positive for anti-GAD aAbs) were immunohistochemically stained *in situ* with the following MMrs: IAPP15-17/5-10 (A-B), ISL1276-284 (C-D), SCG5-009186-194 (E-F), UCN31-9 (G-H) and the positive control PPI6-14 (I-J). For each MMr, pancreas images at 20X magnification are shown on the left (scale bar 100 μ m), and higher magnifications of the dotted areas are shown on the right (scale bar 36 μ m).

	Accession number(s)	Amino acid positions	Sequence	HLA restriction	Predicted affinity (nM)	Predicted stability (h)	IFN-v (%)	IFN-v/TNF-a/IL-1B (%)	Basal (%)	IFN-v log2 FC	IFN-v/TNF-a/IL-18 log2 FC
ABCC8	Q09428	611-619	SEFLSSAEI	B*40:01	46.5	1.6	20	80	0	IFN-y only	IFN-y/TNF-a/IL-1B only
ABCC8	Q09428	787-795	IIFESPFNK	A*03:01	15.6	2.2	80	60	0	IFN-y only	IFN-v/TNF-a/IL-1B only
AMPH	P49418	459-467	EEPVEEAVI	B*40:01	3182.1	0.3	80	40	40	2.64	4.25
AP3B2	013367	403-411	NETNIPTVI	B*40:01	44.7	1.3	40	40	0	IEN-y only	IEN-y/TNE-g/II-18 only
C1011:C1014	086723:075973	196-204	ASNSVILHI	C*03:04	837.2	0.2	100	80	20	4.85	4.19
CADPS:CADPS2	Q86UW7:Q9ULU8	780-788	SLLERVLMK	A*03:01	27.6	5.1	20	60	20	-2.34	3.38
CADPS:CADPS2	086UW7:09UUU8	780-788	SLLERVIM(+15.99)K	A*03:01	27.6	5.1	60	0	20	3 34	Basal only
CHEA	010645	3 10	PEANIALI	C*02:04	568.0	0.2	40	80	40	2.07	6.16
CHGA	P10045	120 120	AVEEDSSKDVA(+1E 00)	C*02:04	2110 5	0.3	40	100	40	2.25	5.27
CIGA	P10045	120-130	AVECP33KDVM(+15.95)	03.04	5115.5	0.2	40	100	40	0.59	3.27
CHGA	P10845	120-131	AVEEPSSKDVIVI(+15.99)E	0840-01	701.2	NA O.F.	40	60	60	U.56	-1.17
CHGA	P10045	121-130		B-40:01	/61.2 NA	0.5	60	60	20	A 25	2.29
CHGA	P10845	121-131	VEEPS3KDVM(+15.99)E	NA	INA 510.0	NA	60	40	20	4.35	2.28
CHGA	P10645	344-352	KMDQLAKEL	A*02:01	543.8	0.6	0	60	80	Basal only	-2.08
CHGA	P10645	344-354	KMDQLAKELTA	A*02:01	823.2	0.6	100	100	40	-1.08	2.45
CHGA	P10645	344-354	KM(+15.99)DQLAKELTA	A*02:01	823.2	0.6	0	0	40	Basal only	Basal only
CHGA	P10645	344-355	KMDQLAKELTAE	NA	NA	NA	20	100	80	-3.38	2.06
CHGA	P10645	380-389	AYGFRGPGPQ	NA	NA	NA	0	40	60	Basal only	-5.55
CHGA	P10645	381-390	YGFRGPGPQL	C*03:04	38.0	0.2	100	80	40	2.91	2.15
CHGA	P10645	383-390	FRGPGPQL	C*07:01	305.3	0.1	60	100	0) IFN-y only	IFN-γ/TNF-α/IL-1β only
CHGA	P10645	402-409	SLEAGLPL	A*02:01	300.4	0.8	0	40	20	Basal only	1.34
CHGA	P10645	402-411	SLEAGLPLQV	A*02:01	115.1	1.3	100	100	0	IFN-γ only	IFN-γ/TNF-α/IL-1β only
CHGA	P10645	403-411	LEAGLPLQV	B*40:01, B*49:01	402.0, 343.9	0.8, 0.7	100	80	0	IFN-γ only	IFN-γ/TNF-α/IL-1β only
CHGB	P05060	603-613	RVAQLDQLLHY	A*03:01	641.8	0.9	40	80	40	-1.04	0.32
CPE	P16870	83-91	FEGRELLVI	B*40:01, B*49:01	223.1, 264.4	0.6, 0.4	100	100	0	IFN-γ only	IFN-γ/TNF-α/IL-1β only
CPE/PTPRN;CPE/PTPRN;CPE/PTPRN	P16870/Q16849;P16870/Q16849;P16870/Q16849	61-66/133-135;61-66/28-30;61-66/949-951	EALVSV/RPG	NA	NA	NA	80	20	0	IFN-y only	IFN-y/TNF-a/IL-1B only
DLK1	P80370	4-12	TEALLRVLL	B*40:01	20.4	1.3	60	80	0	IFN-y only	IFN-y/TNF-a/IL-1B only
ESRRG	P62508	361-369	LEKEEFVTL	B*40:01	45.5	0.9	80	80	C	IFN-y only	IFN-v/TNF-a/IL-1B only
FAM171B	Q6P995	19-28	VLLKARLVPA	A*02:01	148.8	2.4	60	100	40	-0.24	0.69
GAD2	005329	257-265	KM(+15.99)FPEVKEK	A*03:01	48.0	4.7	40	0	0	IFN-y only	NA
GAD2/GAD2	005329/005329	97-104/460-462	LLPAC(+47.98)DGERAK	A*03:01	3984.3	0.3	60	20	20	2.00	-1.64
GPR119	O8TDV5	48-56	AVADTLIGV	A*02:01	93	3.2	20	80		IEN-y only	IEN-y/TNE-g/II-18 only
HSEY1	096116	143-151	ESKIDONEO	NA	5.5 NA	5.2 NA	20	00		IFN-y only	IEN-v/TNF-g/II-18 only
USDAE /USDAE	011021/011021	215 220/74 76	EGEVNIDI	NA	NA	NA	00	100		Not detector	IEN w/TNE g/IL 19 grite
IADD/IADD/DTDDAt/IADD	P11021/F11021	15 17/5 10:505 509/5 10		C*02-04 **03-C*	612 E 2500 4	0.2.0.5	100	100		1.00	1.00
ICCODDO	P10557/F10557;Q10045/P10357	12-11/2-10/220-220/2-10		C 03:04, A 02:01	30.5	0.2, 0.5	100	80	40	1.00	1.96
INTZEPS	0100425	552-56U		A-02:01	20.5	9.3	60	100	0	IFN-y only	IFN-γ/TNF-α/IL-1β only
INA	Q16352	1/0-1/8	GLAEEVQRL	A=02:01	30.7	2.3	80	20	20	3.04	1.68
INS	P01308	2-10	ALWMRLLPL	A*02:01	26.7	5.6	0	40	0	Not detected	IFN-γ/TNF-α/IL-1β only
INS	P01308	2-10	ALW(+3.99)M(+15.99)RLLPL	A*02:01	26.7	5.6	100	80	40	3.75	1.60
INS	P01308	6-14	RLLPLLALL	A*02:01	11.6	8.0	80	60	0	IFN-y only	IFN-γ/TNF-α/IL-1β only
INS	P01308	15-24	ALWGPDPAAA	A*02:01	95.5	1.8	80	100	20	-0.91	3.80
INS	P01308	15-24	ALW(+3.99)GPDPAAA	A*02:01	95.5	1.8	60	100	80	-1.83	2.02
INS	P01308	15-24	ALW(+15.99)GPDPAAA	A*02:01	95.5	1.8	20	60	60	-4.27	-0.57
INS	P01308	15-26	ALWGPDPAAAFV	A*02:01	8.2	6.6	0	60	0	Not detected	IFN-y/TNF-a/IL-1B only
INS	P01308	29-38	HLC(+47.98)GSHLVEA	A*02:01	521.1	1.5	60	0	40	1.40	Basal only
INS	P01308	34-42	HIVEALVIV	A*02:01	3.3	17.8	100	100	0	IEN-y only	IEN-y/TNE-g/II-18 only
INS	P01308	56-66	REAEDLOVGOV	B*40:01	127.8	1.0	0	60	0	Not detected	IEN-y/TNE-g/II-18 only
INIS	001208	E8 66	AEDLOVGOV	R#40-01	064.9	0.6	100	100	40	2.25	2.54
INIS	001208	58 68	AEDLOVGOVEL	R#40:01	26.1	1.5	200	100	40	4.60	4.94
ING DOC	101308	58-08		0 40.01	30.1	1.5	30	00	40	4.05	4.54
INS-UUD	103-008	15 21/427 420	REAEDLOGSL	B-40.01	9.9	1.4	20	100	20		2.27
INS/PIPRN	P01308/Q18849	15-21/437-439	ALWGPDP/KVN	N/A	INA	NA	40	100	20	-1.22	2.27
INS/SLC30A8	P01308/Q8IW04	15-20/30-33	ALWGPD/KPVN	NA	NA	NA	40	40	80	0.18	-2.22
ISLI	P61371	276-284	GLQANPVEV	A*02:01	82.6	2.4	100	U	u	IFN-y only	NA
JAKMIP2;JAKMIP3	Q5VZ66;Q96AA8	31-39	KLTDIQIEL	A*02:01	8.3	6.2	0	80	0	Not detected	IFN-y/TNF-a/IL-1B only
KCNK16	Q96T55	129-137	ALLGIPLNV	A*02:01	11.5	3.8	40	40	0	IFN-γ only	IFN-γ/TNF-α/IL-1β only
KCNK16	Q96T55	13-21	RVLPLLLAY	A*03:01	150.3	0.5	80	100	0) IFN-y only	IFN-γ/TNF-α/IL-1β only
KIF1A;KIF1B;KIF1C	Q12756;O43896;O60333	153-161	RVRDLLNPK	A*03:01	138.5	6.6	100	80	20	3.72	4.53
KIF1A	012756	219-228	AVFNIIFTQK	A*03:01		5.0	80	80	40		5.01
			10 T A A 1 A 10 1 10 1		24.9	5.0	00			4.40	5.01
KIF1A;KIF1B	Q12756;O60333	734-743	KEANAISVEL	B*40:01	24.9 10.6	2.6	0	60	U	0 4.40 Not detected	IFN-γ/TNF-α/IL-1β only
KIF1A;KIF1B KIF1A	Q12756;O60333 Q12756	734-743 796-805	ATHYWTLEKL	B*40:01 C*03:04	24.9 10.6 1925.8	2.6	0	60 100	0	0 4.40 Not detected 0 IFN-γ only	IFN-γ/TNF-α/IL-1β only IFN-γ/TNF-α/IL-1β only
KIF1A;KIF1B KIF1A KIF1A	Q12756;060333 Q12756 Q12756	734-743 796-805 860-868	ATHYWTLEKL LLYPVPLVH	B*40:01 C*03:04 A*03:01	24.9 10.6 1925.8 119.8	2.6 0.2 0.6	0 20 60	60 100 100	0 60	4.40 Not detected IFN-y only 2.37	IFN-γ/TNF-α/IL-1β only IFN-γ/TNF-α/IL-1β only 3.46
KIF1A;KIF1B KIF1A KIF1A KIF1A	012756;060333 012756 012756 012756 012756	734-743 796-805 860-868 877-885	REANAISVEL ATHYWTLEKL LLYPVPLVH GEVKGFLRV	B*40:01 C*03:04 A*03:01 B*40:01	24.9 10.6 1925.8 119.8 106.6	2.6 0.2 0.6 0.7	0 20 60 40	60 100 100 100	60	4.40 Not detected IFN-y only 2.37 2.87	IFN-γ/TNF-α/IL-1β only IFN-γ/TNF-α/IL-1β only 3.46 5.02
KIF1A,KIF1B KIF1A KIF1A KIF1A KIF1A	Q12756;O60333 Q12756 Q12756 Q12756 Q12756 Q12756	734-743 796-805 860-868 877-885 1347-1355	ATHYWTLEKL LLYPVPLVH GEVKGFLRV VLDTSVAYV	B*40:01 C*03:04 A*03:01 B*40:01 A*02:01	24.9 10.6 1925.8 119.8 106.6 6.3	2.6 0.2 0.6 0.7 8.3	0 20 60 40	60 100 100 100 100	0 60 20	4.40 Not detected IFN-y only 2.37 2.87 3.01	IFN-γ/TNF-α/IL-1β only IFN-γ/TNF-α/IL-1β only 3.46 5.02 4.39
KIF1A;KIF1B KIF1A KIF1A KIF1A KIF1A KIF1A	012756,060333 012756 012756 012756 012756 012756 012756 012756 012756	734-743 796-805 860-868 877-885 1347-1355 1480-1488	KEANAISVEL ATHYWTLEKL LLYPVPLVH GEVKGFLRV VLDTSVAYV KLSEMSVTL	B*40:01 C*03:04 A*03:01 B*40:01 A*02:01 A*02:01	24.9 10.6 1925.8 119.8 106.6 6.3 6.9	2.6 0.2 0.6 0.7 8.3 6.2	0 20 60 40 60 40	60 100 100 100 100 20	60 60 20	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only	1.01 IFN-y/TNF-α/IL-1β only IFN-y/TNF-α/IL-1β only 3.46 5.02 4.39 IFN-y/TNF-α/IL-1β only
KIF1A;KIF18 KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A	012756.060333 012756 012756 012756 012756 012756 012756 012756 012756 012756	734-743 796-805 860-868 877-885 1347-1355 1480-1488 1480-1488	KEANAISVEL ATHYWTLEKL LLYPVPLVH GEVKGFLRV VLDTSVAAV KLSEMSVTL KLSEMSVTL KLSEM(+15.99)SVTL	B*40:01 C*03:04 A*03:01 B*40:01 A*02:01 A*02:01 A*02:01	24.9 10.6 1925.8 119.8 106.6 6.3 6.9 6.9 6.9	2.6 0.2 0.6 0.7 8.3 6.2 6.2	0 20 60 40 60 40 40	60 100 100 100 100 20 40	60 60 20 0 40	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71	IFN-yTNF-α/IL-1β only IFN-yTNF-α/IL-1β only 3.46 5.02 4.39 IFN-yTNF-α/IL-1β only -0.54
KIF1A,KIF18 KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A MYO3A	012756;060333 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756	734-743 796-805 860-868 877-885 1347-1355 1480-1488 1480-1488 1480-1488 1517-1525	KEANADVEL ATHYWTLEKL LLYPVPLVH GEVKGFLRV VLDTSVAVY KLSEM(+15.99)SVTL SIQEEKRP	8*40:01 C*03:04 A*03:01 8*40:01 A*02:01 A*02:01 A*02:01 NA	24.9 10.6 1925.8 119.8 106.6 6.3 6.9 6.9 NA	2.6 0.2 0.6 0.7 8.3 6.2 6.2 NA	0 20 60 40 60 40 40 0	60 100 100 100 20 40 40	0 60 60 20 0 40 20	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 Basal only	IFN-γ/TNF-α/IL-1β only IFN-γ/TNF-α/IL-1β only 3.46 5.02 4.39 IFN-γ/TNF-α/IL-1β only -0.54 3.00
KIF1A,KIF18 KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A MY03A NEUROD1	012756,06033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 031562 031562	734-743 796-805 860-868 877-885 1347-1355 1480-1488 1480-1488 1517-1525 291-299	KEANAISVEL ATHYWTLEKL LLYPVPLVH GEVKGFLRV VLDTSVAVV KLSEMSVTL KLSEMSVTL SIQEEKRRP AEFEKNYAF	B*40:01 C*03:04 A*03:01 B*40:01 A*02:01 A*02:01 A*02:01 NA B*40:01	24.9 10.6 1925.8 119.8 106.6 6.3 6.9 6.9 NA 19.9	5.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 6.2 NA 1.7	0 0 20 60 40 60 40 40 40 60 60 60 60 60 60 60 60 60 6	60 100 100 100 20 40 40 100	0 60 60 20 0 40 20 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4.40 Not detected IFN-y only 2.37 3.01 IFN-y only 0.71 Basal only IFN-y only	IFN-γ/TNF-σ/IL-1β only IFN-γ/TNF-σ/IL-1β only 3.66 5.02 4.39 IFN-γ/TNF-σ/IL-1β only -0.54 3.00 IFN-γ/TNF-σ/IL-1β only
KIF1A,KIF1B KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A MY03A NEUR0D1 NPCILI	0.12756,06033 0.12756	734-743 796-805 860-868 877-885 1347-1355 1480-1488 1480-1488 1480-1488 1480-1488 1157-1525 291-299 1158-1167	REAMAISVEL ATHYWTLEKL LLYPVPLVH GEWGFLRV VLDTSVAVV KLSEMSVTL KLSEM(+15.99)SVTL SIQEEKRP AEFEKNYAF LGLDIRSGLL	B*40:01 C*03:04 A*03:01 B*40:01 A*02:01 A*02:01 A*02:01 NA B*40:01 C*03:04	24.9 10.6 1925.8 109.8 6.3 6.9 6.9 NA 19.9 4828.2	5.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 6.2 NA 1.7 0.1	0 0 20 60 40 60 40 40 40 60 40 60 20 60 20 60 20 60 20 60 60 60 60 60 60 60 60 60 6	60 100 100 200 40 40 100 80	0 60 60 20 0 40 20 0 0 0 0 0	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 Basal only IFN-y only IFN-y only IFN-y only	IFN-γTNF-σ/L-1β only IFN-γTNF-σ/L-1β only 3.46 5.02 4.39 IFN-γTNF-σ/L-1β only -0.54 3.00 IFN-γTNF-σ/L-1β only IFN-γTNF-σ/L-1β only
KIF1A,KIF1B KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A MY03A NFUROD1 NPCILI NP//PTPRN,NP//SCG5	012756/06033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 021756 021756 021756 021756 021756 021756 021756 021756 021757 020	734.743 796.805 860.868 1397.1355 1480.1488 1480.1488 1517.1525 291.299 1158.1167 1158.1167 13.821,65.458,13.13,14,488.490	REARAISVEL ATHYVTLERL LLYPVPLVH GEVKGFLRV VLDTSVAVV KLSEMSVTL KLSEMSVTL SIQEEKRP AEFEKNYAF LGLDLRSGLL TALSL/KLL	B*40:01 C*03:04 A*03:01 B*40:01 A*02:01 A*02:01 A*02:01 NA B*40:01 C*03:04 A*02:01	24.9 10.6 1925.8 106.6 6.3 6.9 6.9 6.9 NA 19.9 4828.2 754.6	3.0 2.6 0.2 0.6 0.7 8.3 6.2 0.7 8.3 6.2 NA 1.7 0.1 0.7	0 0 20 60 40 60 40 40 0 0 60 20 80	660 100 100 100 20 40 40 40 40 40 40 40 40 40 40		4.40 Not detected 1FN-y only 2.37 2.87 3.01 1FN-y only 0.71 8asal only 1FN-y only 1FN-y only 1FN-y only 1FN-y only	IFN-yTNF-σ/IL-1β only IFN-yTNF-σ/IL-1β only IFN-yTNF-σ/IL-1β only 3.66 5.02 4.39 IFN-yTNF-σ/IL-1β only -0.54 3.00 IFN-yTNF-σ/IL-1β only IFN-yTNF-σ/IL-1β only IFN-yTNF-σ/IL-1β only IFN-yTNF-σ/IL-1β only IFN-yTNF-σ/IL-1β only IFN-yTNF-σ/IL-1β only
KEFLAKEFLB KEFLA KEFLA KEFLA KEFLA KEFLA KEFLA KEFLA KEFLA MYOJA NEURODI NPC/LIT NPC/LIT NPC/SCS PCSK1	0.12756,06033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.2084V/4 0.13562 0.2094K0 0.13562 0.2094K0 0.13624 0.2094K0	734.743 795.805 860.868 1347.1355 1480.1488 1480.1488 1137.1325 291.299 1138.1167 1138.1167 1138.1167 1138.1167 1138.118	REARAISVEL ATHYWTLEKL LLYPVPLVH GEWGFLRV VLDTSVAVV KLSEMSVTL KLSEM(+15.99)SVTL SIQEEKRP AEFEKNYAF GLODR/SGLL TLALSL/KLL RRGDLHYTL	B*40:01 C*03:04 A*03:01 B*40:01 A*02:01 A*02:01 A*02:01 NA B*40:01 C*03:04 A*02:01 C*03:04 A*02:01 C*07:01	24.9 10.6 1925.8 119.8 6.9 6.9 NA 19.9 4828.2 754.6 204.5	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 NA 1.7 0.1 0.1	0 0 20 60 40 60 40 60 60 60 20 80 100	60 100 100 100 20 40 40 40 30 40 20 20 20 20 20 20 20		4.40 Not detected IFN-y only 2.37 3.01 IFN-y only 0.71 Basal only IFN-y only IFN-y only IFN-y only IFN-y only IFN-y only	IFN-γTNF-67/L-1β only IFN-γTNF-67/L-1β only IFN-γTNF-67/L-1β only 5.02 4.39 IFN-γTNF-67/L-1β only 0.54 3.00 IFN-γTNF-67/L-1β only IFN-γTNF-67/L-1β only IFN-γTNF-67/L-1β only IFN-γTNF-67/L-1β only IFN-γTNF-67/L-1β only
KETA,KETAB KETAA KE	012756/06033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 021757 021757 020	734-743 796-805 860-868 1397-1355 1480-1488 1480-1488 1517-1525 291-299 1158-1167 13.82/165-853 13.82/164-848-490 517-525 517-	REARAISVEL ATHYVITLEKL LLYPPULVH GEVKGFLRV VUDTSVAVV KLSEM(+15.99)SVTL SIGEEKRRP AEFEKNYAF GEJDLRSGLL TLALSL/KLL RRGDLHVTL KINRPSOLK	B*40:01 C*03:04 A*03:01 B*40:01 A*02:01 A*02:01 A*02:01 NA B*40:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01	24.9 10.6 1925.8 119.8 106.6 6.9 6.9 8.9 19.9 4828.2 754.6 204.5 16.9	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 6.2 NA 1.7 0.1 0.1 0.7 0.1 13.1	0 0 20 60 40 40 40 40 60 20 80 100 40	60 100 100 20 40 40 100 80 40 100 80 100 100 100 100 100 100		4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 8.8331 only IFN-y only IFN-y only IFN-y only IFN-y only IFN-y only IFN-y only	IFN-yTNF-67/L-1β only IFN-yTNF-67/L-1β only IFN-yTNF-67/L-1β only 5.02 4.39 IFN-yTNF-67/L-1β only -0.54 3.00 IFN-yTNF-67/L-1β only
KGF1A, KGF1A KGF1A KGF1A KGF1A KGF1A KGF1A KGF1A MC03A NCURCO1 NCUL1 NCV/DFNRyAPY/SCG5 PCSK1 PCSK1 PCSK2	0.12756.06033 0.12756	734.743 778.753 860.868 1347.1355 1460.1488 1480.1488 1480.1488 1189.1487 118.9167 11.18/167 11.18/164-563.13.18/488-490 517.255 705.714 30.38 51.255 5	REAMARSVEL ATHWATTERK LLYPPPUM GEVKGERRV VLDTSVATV KLSEMSVTL SIGEEKRRP AEFENTVAF LGIDLRSGLL TLALS/KLL RRGDH/VTL KLINEPSQLK ETMERVFI	B*40:01 C*03:04 A*03:01 B*40:01 A*02:01 A*02:01 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01 C*07:01 C*07:01 C*03:04 A*02:01 C*07:01 C*03:04 C*03:04 C*03:04 C*03:04 C*03:04 C*03:04 C*03:05 C*03:04 C*03:05 C*0	24.9 10.6 1925.8 109.6 6.9 6.9 6.9 NA 19.9 4828.2 754.6 204.5 16.9 34.4.00	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 6.2 0.1 1.7 0.1 1.7 0.7 0.1 13.1 2.6 0.4	00 00 60 40 60 40 40 40 60 20 80 100 60	660 100 100 200 400 400 100 800 400 200 100 800 800 800 800 800 800 800 800 8	0 60 60 20 40 40 40 40 40 40 40 40 40 4	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 1.712	IFN+γTNF-σ(1)-1β only IFN+γTNF-σ(1)-1β only IFN+γTNF-σ(1)-1β only 0.54 3.00 IFN+γTNF-σ(1)-1β only IFN+γTNF-σ(1)-1β only IFN+γTNF-
KF1A,KF18 GF1A GF1A GF1A GF1A GF1A GF1A GF1A GF1A KF1A MF0A M	012756/06033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 021562 021556 021556 021556 021556 021556 021556 021556 021556 020	734-743 778-705 660-868 1347-1355 1480-1488 1480-1488 1517-1525 151-1	REAMANSVEL ATHWATTERK LLYPP/PU/H GEVKGFLRV VUDTSVANV KLSEM/011 SLGEKRRP ALSEM/011 GLDLRSGLL TLASL/KL RRGDH/VTL KLNPSQLK FTNHFVEL PFTG/RPI1	8*40:01 C*03:04 A*02:01 A*02:01 A*02:01 A*02:01 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 C*0	24.9 10.6 1925.8 119.8 106.6 6.9 6.9 8.9 19.9 4828.2 754.6 204.5 16.9 43.4,40.0 2916.3	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 NA 1.7 0.1 0.7 0.1 13.1 2.6,0.4 0.2	0 20 60 40 40 40 0 60 20 80 100 60 60 60	60 100 100 100 200 40 40 100 80 40 200 80 80 40 200 80 100 80 100 80 100 100 100	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 Basal only IFN-y only IFN-y only IFN-y only IFN-y only IFN-y only 1FN-y	IFN-r/TN-e7/L-15 only IFN-r/TN-e7/L-15 only
KRF1A,KRF1B KRF1A K	0.12756.06033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.13562 0.01960 P01301/(10549/p01303/P05408 P29120 P91202 P91210 P95131 P55131/016849/P15519/016849 P55345	734.743 778.753 860.868 1347.1355 1460.1488 1480.1488 1480.1488 1189.1487 118.9167 11.18/1467 11.18/1465 11.525 706.714 30.38 479.483/085-888 10.0.17 10.1	REAMARSVEL ATHWYTLER. LLYPPVUH GEVKGFLRV VUDTSVAVV KLSEM/15.99JSVTL KLSEM/15.99JSVTL KLSMSGLL TLALS/LKL REGDIH/VTL KLNKPSQLK TTMFLVEL PPTG/RPLL EERDRV/I	8*40:01 C*03:04 A*03:01 8*40:01 A*02:01 A*02:01 A*02:01 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 B*40:01 C*03:04 B*40:01 C*03:04 B*40:01 C*03:04 B*40:01 C*03:04 B*40:01 C*03:04 D*02:01 C*03:04 D*02:01 C*03:04 D*02:01 C*03:04 D*02:01 C*03:04 D*02:01 D*0	24.9 10.6 1925.8 119.8 6.9 6.9 6.9 19.9 4828.2 754.6 204.5 16.9 43.4,40.0 2316.3 6.2 16.9 16.9 16.9 16.9 16.9 16.9 19.9 19.9 10.6 19.9 10.6 19.9 10.6 19.9 10.6 19.9 10.6 10.9 10.9 10.9 10.6 10.9 10.	5.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 NA 1.7 0.1 0.7 0.1 13.1 2.6,0.4 0.2 0.9	0 0 20 60 40 40 40 40 60 20 80 100 40 60 60 20 20 20 20 20 20 20 20 20 2	660 100 100 200 40 40 40 40 20 30 40 100 30 40 100 30 30 100 30 30 30 30 30 30 30 30 30 30 30 30 3	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 Basal only IFN-y only	IFN-y(TNI-sc/L-1) only IFN-y(TNI-sc/L-1) only IFN-y(TNI-sc/L-1) only 1.40 4.30 IFN-y(TNI-sc/L-1) only
KF1A,KF18 KF1A K	012756/06033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 013562 000405 013562 000405 013562 000405 013562 000405 013562 000405 0005000000	734-743 778-805 660-868 1347-1355 1480-1488 1480-1488 1480-1488 1517-1535 134-1185 1153-1187 13-141/64-164,13-18/488-490 157-535 706-714 15-535 706-714 17-535 706-714 17-535 706-714 17-535 706-714 17-535 706-714 17-535 706-714 17-535 706-714 17-535 706-714 17-535 706-714 70-545 70-555 70-555 70-555 70-555 70-555 70-555 70-555 70-555 70-555 70-555 70-555 70-555 70-555 70-555 70-555	REAMANSVEL ATHWATTERK LLYPP/PU/H GEVKGFLRV VUDTSVATV KLSEM/015.90JSVTL SIGEEKRRP AEEENVAF GJDLRSGUL TLALSJ/KUL RRGDH/VTL KLNOPSQUK FTNHFLVEL EFPNRVQL EEPPRVQL EEPPRVQL	8*40.01 C*03.04 A*03.01 8*40.01 A*02.01 A*02.01 A*02.01 A*02.01 C*03.04 A*02.01 C*03.04 A*02.01 C*07.01 A*03.01 A*0	24.9 10.6 1925.8 119.8 106.6 6.3 6.9 NA 19.9 4828.2 754.6 204.5 16.9 43.4,40.0 2916.3 6.2 1721.7	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 NA 1.7 0.1 0.7 0.1 13.1 2.6,0.4 0.2 0.9 0.4	0 0 20 60 40 40 40 40 40 60 20 80 40 60 60 20 40 60 20 20 80 100 40 20 80 100 100 100 100 100 100 100	660 100 100 100 20 40 40 40 40 20 80 40 20 80 40 20 80 40 20 100 80 40 20 100 80 100 100 100 80 100 100		4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 Basal only IFN-y only 1.39 1.35	IFN-r/TN-eg/L-15 only
KRF1A, KRF1B KRF1A	0.12756.06033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.11562 0.01400 P01190/101849.P01303/P05408 P01392100 P01392(102849.P01303/P05408 P03120 P0312 P0312 P03120 P03120 P03120 P03120 P	734.743 724.743 726.805 860.868 860.868 1347.1355 1480.1488 1480.1488 1480.1488 1480.1488 1189.1487 131.81/165-1561.31.81/488-490 517.255 706.714 30.38 479.483/707.37.479.483/885-888 109.117 158.166 109.117 158.166 30.389	REAMARSVEL ATHWYTLER. LLYPPVUH GEVKGFLRV VUDTSVAVV KLSEMVIT KLSEMVIT KLSEMSTU KLSEMSTU KLSEMSTU LESPKSUL TUALSL/KLL REGDHYTL KLNKPSQLK TUALSL/KL LESPKRVQL LESPKRVQL LESPKRVQL LESPKRVQL	8*4001 CV03:04 A*02:01 A*02:01 A*02:01 A*02:01 A*02:01 A*02:01 CV03:04 A*02:01 C*07:01 A*02:01 C*07:01 A*02:01 C*07:01 A*02:01 C*07:04 B*40:01 B*40:01 B*40:01	24.9 10.6 1925.8 119.8 10.6 6.9 6.9 NA 19.9 10.9 1	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 NA 1.7 0.1 0.1 0.1 0.1 0.7 0.1 13.1 2.6,0.4 0.2 0.9 0.4	0 0 20 40 60 40 40 40 60 20 80 40 60 60 60 60 60 20 20 20 20 20 20 20 20 20 2	66 100 100 100 100 20 40 40 40 40 100 80 40 20 100 80 40 20 100 80 40 40 40 40 40 40 40 40 40 4	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4.40 Not detected IN-ry only 2.37 2.87 3.00 10-12 10-	IFN-y(Thi-a,0.1.2) only IFN-y(Thi-a,0.1.2) only 1.4 IFN-y(Thi-a,0.1.2) only 1.6 4.9 IFN-y(Thi-a,0.1.2) only 2.70
KF1-A,KF1-B KF1-A KF1	012756/06033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 001755 001755 001755 001755 001755 0017 0017	734-743 778-805 860-868 1347-1355 1460-1488 1480-1488 1517-1525 291-299 13-142/16-156(13-18/488-490 13-15/155 706-714 13-83/16-156(13-18/488-490 13-55) 706-714 13-83/16-156(13-18/488-488 109-137 15-55 109-137 109-137 154-166 109-137 154-166 109-137 154-166 109-137 154-166 109-137 154-166 109-137 154-166 109-137 154-166 109-137 154-166 109-147 1	REAMANYEL ATHWYTERA LLYPYUH GEVGETAV VIDTSVATV KISSMATT K	8*4001 C*03:04 A*03:01 8*40:01 A*02:01 A*02:01 A*02:01 A*02:01 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 B*40:01 8*40:01 8*40:01 8*40:01 8*40:01	24.9 10.6 1925.8 119.8 10.6 6.3 6.9 NA 19.9 482.8.2 204.5 16.9 43.4.40.0 2916.3 62.2 1721.7 87.0 87.0 10.6 10.9 10.6 10.9 10.6 10.9 10.6 10.9 10.6 10.9 10.6 10.9 10.6 10.9 10.6 10.9 10.6 10.9 10.6 10.9 10.6 10.9 10.9 10.6 10.9 10.9 10.9 10.6 10.9 1	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 8.3 6.2 8.3 0.1 0.1 0.1 0.1 13.3 2.6,0.4 0.2 0.9 0.4 1.0 1.0 1.0 1.0	0 0 20 60 40 40 40 40 40 60 60 60 60 200 100 20 0 20 0 80 100 20 0 20 0 80 100 100 100 100 100 100	66 60 100 100 100 100 100 100 10	0 600 600 200 200 200 0 0 0 0 0 0 0 0 0 0 0 0	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 Basal only IFN-y	IFN+/TN-eg/L-18 only IFN+/TN-eg/L-18 only IFN+/TN-eg/L-18 only IFN+/TN-eg/L-18 only O-54 S02 IFN+(TN-eg/L-18 only IFN+/TN-eg/L-18 only
KIF1A,KIF18 KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A MV3A NCURCO1 NV/VFTNR,NPV/SCG5 PCSK1 PCSK1 PCSK1 PCSK1 PCSK1 PCSK1 PCSK2 PCSK	0.12756.00033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12755 0.12755 0.12755 0.12755 0.12755 0.12755 0.12755 0.11552 0.1014 0.11562 0.0140 0.11562 0.0140 0.11551 0.11519 0.1551 0.1551 0.155 0.155 0.155 0.155 0.155 0.155 0.155 0.15 0.1	734.743 724.743 726.805 860.868 860.868 1347.1355 1480.1488 1480.1488 1480.1488 1480.1488 1189.1487 131.81/165-563.13.81/488-490 517.255 706.714 30.38 479.433/085.888 109.117 158.166 1880.3582 142.4120 159.106 180.358 142.4120 159.106 150.358 159.35 159.35 1	AEANANSVEL ATHWYTLER. LLYPPVUH GEVKGFLRV VUDTSVANV KLSEMVIT KLSEMVIT KLSEMSUT LGDURSGLL TLALSLKL RKODHVTL KLNNFSQLK TTNHFLE IPSTG/RPLL LEFNRVQL FEDPORREY HEYYAQNGI SLAQMVPKL SLAQMVPKL	8*40:01 C*03:04 A*02:01 A*02:01 A*02:01 A*02:01 NA 8*40:01 C*03:04 A*02:01 C*03:04 A*02:01 C*03:04 B*40:01	24.9 105 1075.8 1198 106.6 6.3 6.9 6.9 6.9 482.8 204.5	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 NA 1.7 0.1 0.1 0.7 0.1 13.1 13.1 2.6,0.4 0.2 0.9 0.4 1.0 4.2	0 0 20 40 40 40 40 40 20 20 20 20 20 20 20 20 20 2	66) 100 100 100 100 100 20 20 20 20 20 20 20 20 20	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4.40 Not detected If N=y only 2.37 2.87 3.01 10 - 2.01 If N=y only 1.32 1.32 1.32 1.33 If N=y only 1.33 If N=y only 1.33 If N=y only 1.33	IFN-y(Thi-a(1):1) only IFN-y(Thi-a(1):1) only 1.6 1.6 1.6 IFN-y(Thi-a(1):1) only 1.6 IFN-y(Thi-a(1):1) only
NF1A_KF18 NF1A NF1A NF1A OF1A OF1A NF1A NF1A OF1A OF1A NF1A NF1A NF1A NF1A NF1A NF2A	012756,06033 012756 012756 012756 012756 012755 012755 012755 012755 012755 012755 012755 012755 012755 012755 000HC9 0014C9 000000000000000000000000000000000000	734-743 778-505 560-868 1347-1355 1460-1488 1480-1488 1480-1488 1517-1525 131-1516-158(13-18/488-490 13-18/166-158(13-18/488-490 13-15/155 13-555	REAWARSVEL ATHWYTERA LLYPPUH GEWGELAV VIDTSVATV KISSMATT	8*40:01 (************************************	24.9 10.6 1925.8 119.8 10.6 6.3 6.9 NA 19.9 482.8.2 754.6 204.5 16.9 43.4.40.0 2016.3 6.2 6.2 173.1.7 87.0 22.0 140.5 140	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.2 NA 1.7 0.1 0.1 0.1 13.3 2.6,0.4 0.2 0.9 0.4 1.0 1.0 2.6,0.4 0.2 0.9 0.4 1.0 1.0 1.0 2.0 0.2 0.2 0.6 0.2 0.6 0.7 8.3 6.2 0.6 0.7 8.3 6.2 0.6 0.7 8.3 6.2 0.6 0.7 8.3 6.2 0.6 0.7 8.3 6.2 0.6 0.7 8.3 6.2 0.6 0.7 8.3 6.2 0.6 0.7 8.3 0.6 0.7 8.3 0.6 0.7 8.3 0.6 0.7 0.7 0.7 0.7 0.7 0.7 0.7 0.7 0.7 0.7	0 0 20 60 40 60 40 60 60 60 20 80 0 20 80 0 20 80 0 20 100 100 100 100 100 100	66 100 100 100 100 100 100 100 1	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 Basal only IFN-y only IFN-y only IFN-y only IFN-y only IFN-y only IFN-y only 1.35 1.23 1.35 1.23 IFN-y only Not detected	IFN-ryTN-erg/L-18 only IFN-ryTN-erg/L-18 only IFN-ryTN-erg/L-18 only 3.66 5.02 IFN-ryTN-erg/L-18 only -0.54 3.00 IFN-ryTN-erg/L-18 only
KIF1A,KIF18 KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A MV03A NEUROD1 NV/UFTRN,NP/SCGS PCSR1 PCSR1 PCSR1 PCSR1 PCSR1 PCSR1 PCSR1 PCSR2 PCSR	0.12756.06033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12755 0.12755 0.12755 0.12755 0.12755 0.12757 0.13562 0.0144 0.13562 0.0144 0.13562 0.0144 0.13562 0.0144 0.1356 0.014 0.1350 0.014 0.1350 0.014 0.1350 0.014 0.1350 0.014 0.0	734.743 724.743 726.805 860.868 860.868 1347.1355 1480.1488 1480.1488 1480.1488 1480.1488 1189.1487 1118.91167 1118.91167 1118.91167 1118.91167 1138.9166 1380.388 1144.152.4124.20 239.298 1289	AEANANSVEL ATHWYTLER. LLYPPVUH GEVKGFLRV VUDTSVAVV KLSEMYT KLSEMYT KLSEMST GLORESGL TLALS/KLL RKODHYTL KLNNFSQLK FTNHFLVEL PPTG/RPL LEPNNVQL FEDPORREY HEVYAONGI SLACYMPKL BLOKKMKSI REIDKOMKSI	8*40:01 C*03:304 A*02:01 A*02:01 A*02:01 A*02:01 A*02:01 A*02:01 C*03:304 A*02:01 C*03:304 A*02:01 C*03:304 B*40:01	24.9 105 1075.8 1198 106.6 6.3 6.9 6.9 8.0 19.9 482.8 204.5	336 226 02 05 83 62 62 84 17 01 01 01 131 131 131 26,04 02 09 04 10 10 10 10 10 10 10	0 0 20 60 40 60 40 60 60 60 60 60 60 20 100 60 60 0 20 0 0 0 0 0 0 0 0 0 0 0 0 0	660 1000 1000 1000 1000 200 404 402 1000 803 404 202 1000 803 1000 803 804 1000 805 805 805 805 805 805 805	C C C C C C C C C C C C C C C C C C C	4.40 Not detected	IFA-y(Thi-a,d):13 only IFA-y(Thi-a,d):15 only 1.6 1.6 1.6 IFA-y(Thi-a,d):15 only IFA-y(Thi-a,d):15 only IFA-y(Thi-a,d):15 only IFA-y(Thi-a,d):15 only IFA-y(Thi-a,d):16 only IFA-y(Thi-a,d):10 only
NRTA.KYT38 NRTA NRTA <td>012756,06033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 01007(01649,91010)/05408 971300 971000 971300 971000 9710000000000</td> <td>734.743 778.605 860.868 1347.1355 1460.1488 1140.1488 1157.1525 231.299 1154.1167 13.12/146-46313.12/488-490 157.525 706.714 80.388 479.4431/073,479-483/885-888 109-117 158.166 109-117 158.166 109-117 158.165 282.298 282.288</td> <td>REAMANSVEL ATHWITERL LLYPPUH GEWGELRV VUDTSVARV KISSMAYT KISSMA</td> <td>8*40:01 (************************************</td> <td>24.9 10.6 1925.8 119.8 10.6 6.3 6.9 NA 19.9 482.8.2 204.5 16.9 43.4.40.0 2916.3 62.2 1721.7 87.0 22.0 140.5 16.3 17.5 16.3 17.5 16.3 16.3 16.3 16.3 16.3 17.5 16.3 17.5 16.9 16.3 16.9 16.3</td> <td>3.0 2.6 0.2 0.6 0.7 8.3 6.2 0.6 1.7 0.1 0.1 0.1 0.1 0.1 0.1 0.1 0.1 1.3.1 2.6.0.4 0.2 0.9 0.4 1.0 4.2 1.1 1.00 0.1</td> <td>0 0 60 40 40 0 0 0 0 0 0 0 0 0 0 0 0 0</td> <td>66 66 100 100 100 20 40 40 40 40 80 40 20 100 80 40 100 80 80 80 80 80 80 80 80 80</td> <td>C C C C C C C C C C C C C C C C C C C</td> <td>4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 Basal only IFN-y only IFN-y only IFN-y only IFN-y only IFN-y only 1.35 1.23 IFN-y only 1.35 1.23 IFN-y only 1.35 2.24 IFN-y only 1.35 1.23 IFN-y only 1.35 1.35 1.35 1.35 1.35 1.35 1.35 1.35</td> <td>IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only 3.66 5.02 4.39 IFA-Y(Thi-ry(k-1)) only 0.54 3.00 IFA-Y(Thi-ry(k-1)) only 0.54 3.00 IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only 2.70 4.51</td>	012756,06033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 01007(01649,91010)/05408 971300 971000 971300 971000 9710000000000	734.743 778.605 860.868 1347.1355 1460.1488 1140.1488 1157.1525 231.299 1154.1167 13.12/146-46313.12/488-490 157.525 706.714 80.388 479.4431/073,479-483/885-888 109-117 158.166 109-117 158.166 109-117 158.165 282.298 282.288	REAMANSVEL ATHWITERL LLYPPUH GEWGELRV VUDTSVARV KISSMAYT KISSMA	8*40:01 (************************************	24.9 10.6 1925.8 119.8 10.6 6.3 6.9 NA 19.9 482.8.2 204.5 16.9 43.4.40.0 2916.3 62.2 1721.7 87.0 22.0 140.5 16.3 17.5 16.3 17.5 16.3 16.3 16.3 16.3 16.3 17.5 16.3 17.5 16.9 16.3 16.9 16.3	3.0 2.6 0.2 0.6 0.7 8.3 6.2 0.6 1.7 0.1 0.1 0.1 0.1 0.1 0.1 0.1 0.1 1.3.1 2.6.0.4 0.2 0.9 0.4 1.0 4.2 1.1 1.00 0.1	0 0 60 40 40 0 0 0 0 0 0 0 0 0 0 0 0 0	66 66 100 100 100 20 40 40 40 40 80 40 20 100 80 40 100 80 80 80 80 80 80 80 80 80	C C C C C C C C C C C C C C C C C C C	4.40 Not detected IFN-y only 2.37 2.87 3.01 IFN-y only 0.71 Basal only IFN-y only IFN-y only IFN-y only IFN-y only IFN-y only 1.35 1.23 IFN-y only 1.35 1.23 IFN-y only 1.35 2.24 IFN-y only 1.35 1.23 IFN-y only 1.35 1.35 1.35 1.35 1.35 1.35 1.35 1.35	IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only 3.66 5.02 4.39 IFA-Y(Thi-ry(k-1)) only 0.54 3.00 IFA-Y(Thi-ry(k-1)) only 0.54 3.00 IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only IFA-Y(Thi-ry(k-1)) only 2.70 4.51
KIF1A,KIF18 KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A KIF1A MV3A NCUROD1 NV/VFTNR,NPV/SCG5 PCSK1 PCSK1 PCSK1 PCSK1 PCSK1 PCSK1 PCSK1 PCSK1 PCSK2 PCSK	0.12756.06033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12755 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.127 0.12 0.127 0.127 0.127 0.12 0.127 0.127 0.12 0.12 0.12 0.12	734.743 724.743 726.805 860.868 860.868 1347.1355 1460.1488 1460.1488 1460.1488 1460.1488 1151.71.525 7291.299 11158.1167 11158.1167 158.1166 109.117 158.1166 109.117 158.1166 109.128 109.12	AEANANSVEL ATHWYTLER. LLYPPVUH GEVKGFLRV VUDTSVAVY KLSEM-15.99JSVTL SIGEEKRAP AEFERNVAF GIDLRSGLL TLALSL/KLL RKODHVTL KLNNFSQLK TTNHFLEE IPSTG/RPL LEEPNRVQL FEDPORBEV HEVYAQNGI SLAQVNPKL MELGARARK RRTAQVRL GLEDLANKSI RLGARARK	8*40:01 C*03:04 A*03:01 8*40:01 A*02:01 A*02:01 A*02:01 A*02:01 A*02:01 C*03:04 B*40:01 C*03:04 B*40:01 B*4	24.9 105 1075.8 1193.8 106.6 6.3 6.9 6.9 8.0 19.9 4828.2 204.5 16.3 62.2 1731.7 87.0 2216.3 140.5 140.5 140.5 140.5 140.5 140.5 140.5 1731.7 87.0 22.0 140.5 16.3 48.0 22.5 16.3 16.3 16.3 16.3 16.3 173.7 16.5 173.7 16.5 173.7 16.5 173.7 16.5 173.7 173.7 16.5 173.7 173.7 174.5	3 2 6 2 2 6 0 2 0 6 0 7 8 3 6 2 0 7 8 3 6 2 0 7 0 7 0 1 0 7 0 1 0 7 0 1 1 3 1 1 3 1 2 6, 0 4 0 2 0 9 0 4 1 0 1 0 1 1 1 1 1 1 1 0 1 0 1 1 1 1	0 0 60 40 40 40 0 60 20 80 100 60 20 100 0 0 0 0 0 0 0 0 0 0 0 0	660 1000 1000 1000 1000 202 404 404 1000 803 404 1000 803 1000 803 803 803 803 803 803 803	C C C C C C C C C C C C C C C C C C C	4.40 Not detected Not detected Not detected Not detected 2.37 3.01 Not only Not not Not not Not not 1.32 1.33 1.33 1.33 1.33 1.33 1.33 1.33	IFR×yTNR-d(L-1) B only IFR×yTNR-d(L-1) B only 1.64 1.05 IFR×yTNR-d(L-1) B only 0.14 IFR×yTNR-d(L-1) B only 0.14 IFR×yTNR-d(L-1) B only FR×yTNR-d(L-1) B only FR×yTNR-d(L-1) B only FR×yTNR-d(L-1) B only IFR×yTNR-d(L-1) B only IFR×yTNR-
NCF1A, KCF1B NCF1A NC03A NCU001 NVCL11 NVCVC05 PCSR1 PCSR1 PCSR2 PCSR2 PCSR3 PCSR3 PMAA2 PMAA2 PRMA2 PRM4 PCPH PCPH	012756,060333 012756 012757 00000000000000000000000000000000000	734.743 778.605 860.868 1347.1355 1460.1488 1140.1488 1157.1525 291.299 1158.1167 13.18/166.168,13.18/488.490 157.525 706.714 30.388 473.483/7073,479-483/885.888 109.117 158.166 139.388 144.152,412.470 289.298 50.5	REAWARSVEL ATHWYTERK LLYPPVLWH GEWGELRY VLDTSVARV KLSEMSYTL KLSEMSYTL KLSEMSYTL GEDKRAGL GEDKRAGL GEDKRAGL EFNRYAGL EEPNRYAGL EEPNRYAGL EEPNRYAGL EEPNRYAGL EEPNRYAGL EEPNRYAGL EEPNRYAGL EEPNRYAGL EEPNRYAGL EEPNRYAGL EEPNRYAGL	8*40.01 C*03.34 A*03.01 8*40.01 A*02.01 A*02.01 A*02.01 A*02.01 A*02.01 C*03.34 B*40.01 C*03.34 B*40.01 B*40.01 B*40.01 B*40.01 B*40.01 B*40.01 B*40.01 A*02.01 A*0	24.9 10.6 1925.8 119.8 10.6 6.3 6.9 0.9 19.9 482.8.2 204.5 16.9 484.4.00 2216.3 62.2 1731.7 87.0 220.5 16.3 48.0 72.0 16.3 48.0 72.5 55.8	3 3 6 2 6 0 2 0 6 0 6 0 7 7 8 3 6 2 8 3 6 2 8 3 6 2 8 3 6 2 8 3 0 1 0 1 0 1 1 7 0 1 1 3.1 2 6,04 0 2 0 9 0.4 1.0 1.0 0.1 0.1 0.1 0.1 0.2 0.2 0.1 0.1 0.1 0.1 0.2 0.1 0.1 0.1 0.1 0.2 0.2 0.1 0.1 0.1 0.1 0.1 0.2 0.2 0.4 0.1 0.1 0.1 0.1 0.1 0.2 0.2 0.4 0.1 0.1 0.1 0.2 0.4 0.1 0.1 0.1 0.2 0.4 0.1 0.1 0.1 0.1 0.1 0.1 0.1 0.1	0 0 0 0 0 0 0 0 0 0 0 0 0 0	66 60 100 100 100 202 40 40 40 40 80 40 202 100 80 40 202 100 80 80 80 80 80 80 80 80 80	C C C C C C C C C C C C C C C C C C C	4.40 Not detected INV only 2.37 3.00 0.71 9.00 0.71 9.00 0.71 9.00 0.71 9.00 0.71 9.00 0.71 0.00 0.71 0.00 0.71 1.35 1.39 IFN y only 1.35 1.23 IFN y only 1.35 1.35 1.35 1.35 1.35 1.35 1.35 1.35	IFN-Y(TNF-y(L-1)) only IFN-Y(TNF-y(L-1)) only 3.66 5.02 4.39 IFN-Y(TNF-y(L-1)) only 0.54 3.00 IFN-Y(TNF-y(L-1)) only IFN-Y(TNF-y(L-1)) only IFN-Y(TNF-y(L-1)) only IFN-Y(TNF-y(L-1)) only 1.09 6.78 3.49 IFN-Y(TNF-y(L-1)) only 2.77 4.21 IFN-Y(TNF-y(L-1)) only 2.70 4.71 IFN-Y(TNF-y(L-1)) only 2.70 4.51 IFN-Y(TNF-y(L-1)) only 2.70 4.51 IFN-Y(TNF-y(L-1)) only 2.75
KRF1A,KF718 KRF1A PCSR1 P	0.12756.06033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12755 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.127 0.12 0.127 0.12 0.12 0.127 0.12 0.12 0.12 0.12 0.12 0.12 0.12 0.12	734.743 724.743 726.805 860.868 860.868 1347.1355 1460.1488 1460.1488 1460.1488 1460.1488 1153.1157.155 291.299 11158.1167 11158.1167 1158.1167 105.11	REAWGAYEL ALTWYTERL LLTWYCH LLTWYCH MODSANT MODSANT SCHWIGTSONT SC	8*4001 (*03304 4*0301 8*4001 4*0201 4*0201 4*0201 8*4001 8*4001 4*0201 (*03304 4*0201 8*4000000000000000000000000000000000000	24.9 10.6 1925.8 119.8 10.6 6.3 6.9 6.9 6.9 10.9 4828.2 754.6 204.5 16.9 4828.2 754.6 204.5 16.3 87.0 220.1 140.5 16.3 48.0 7.2 59.8 326.7 25.8 26.7 26.9 26.9 27.8 26.9 27.8 27.0 27.8 27.	3.0 2.6 0.2 0.7 8.3 6.2 6.2 8.3 6.2 8.3 8.4 8.4 8.4 9.4 0.1 0.1 0.1 0.1 0.1 13.1 13.1 13.1 13.1	0 0 600 400 400 400 200 800 1000 200 800 8	66 66 100 100 100 100 20 20 40 40 40 100 80 100 80 100 80 100 80 100 80 100 10	C C C C C C C C C C C C C C C C C C C	4.40 Not detected	IFN-y/TN-ext-15 only
KRTLAKF18 KRTLA KRTLAKER KRTLAKER<	0.12756,00033 0.12756 0.127570 0.127570 0.127570 0.127570 0.127570 0.127570 0.127570 0.127570 0.1275700 0.1275700000000000000000000000000000000000	734.743 734.743 736.805 860.868 1347.1355 1480.1488 1140.1488 1157.1525 291.299 1158.1167 13.18/166.158,13.18/488.490 157.525 706.714 30.388 473.483/70.73,479.483/885.888 109.117 158.166 189.388 144.152,412.420 289.298 50.58 50.58 50.516 156.164 50.516 50.516 50.516 50.516 50.505 50.516 50.516 50.516 50.505 50.516 50.516 50.505 50.516 50.505 50.516 50.516 50.505 50.516	REAMARYEL ATHWITER LLYPPUH GYKGFLW VUDTSWAVY KLSEMYTL KLSEMYTL KLSEMSTL GJDLRSGLL TLALS/KLL RRGDH/TL KLNKYGLK FTNHFLE EPPNRVGL EEPNRVGL EEPNRVGL EEPNRVGL EEPNRVGL EEPNRVGL EEPNRVGL EEPNRVGL EEPNRVGL EEPNRVGL EEPNRVGL EEPNRVGL	8*4001 (*03344 4*03301 8*4001 4*0201 4*0201 4*0201 4*0201 4*0201 4*0201 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201	24.9 10.5 10.5 11.5 10.5 11.5 10.6 11.5 10.6 10.5 1	3 3 6 2 6 0 2 0 6 0 7 0 7 0 7 0 7 0 7 0 1 0 7 0 1 1 3 1 2 6 0 4 0 2 0 9 0 4 1 0 1 0 1 1 1 0 0 1 1 0 0 1 1 0 0 1 0 1	0 0 0 0 0 0 0 0 0 0 0 0 0 0	660 1000 1000 1000 1000 202 404 402 1000 803 404 202 1000 803 803 1000 803 803 803 803 803 803 803	C C C C C C C C C C C C C C C C C C C	4.40 Not detected	IFN-Y(TNF-y(L-1)) only IFN-Y(TNF-y(L-1)) only 3.66 Y-Y(TNF-y(L-1)) only 3.67 Y-Y(TNF-y(L-1)) only 9.54 3.00 IFN-Y(TNF-y(L-1)) only 9.54 3.00 IFN-Y(TNF-y(L-1)) only IFN-Y(TNF-y(L-1)) only IFN-Y(TNF-y(L-1)) only 7.77 4.21 IFN-Y(TNF-y(L-1)) only 2.70 4.21 IFN-Y(TNF-y(L-1)) only 2.70 4.91 IFN-Y(TNF-y(L-1)) only 6.35 3.31 IFN-Y(TNF-y(L-1)) only 6.35 3.31 IFN-Y(TNF-y(L-1)) only
KF1A,KF1B GF1A G	012756/06033 012756 012756 012756 012756 012756 012755 012755 012755 012755 012755 001454 001452 000450 000450 000450 000450 000450 000450 000450 000450 000450 000450 000450 00050 000450 00050 000450 00050 00045 000450 00050 00045 000	724.743 724.743 724.743 724.743 724.743 660.868 72.865 1347.1355 1440.1488 1480.1488 1480.1488 151.715.25 13.147.061 13.147.061 13.147.061 13.147.061 13.147.061 13.147.061 13.147.061 13.147.061 13.147.061 130.348 109.117 13.147.061 130.348 109.117 130.3 109.11 14.152.112.420 289.298 109.12 17.139 17.139 17.139 17.139 17.135 154.144 159.3156 154.154 159.3156 154.154 159.3156 154.154 159.3156 154.154 159.3156 155.154 155.155 155.154 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155.155 155	REAWGRYEE AFHWYTERL LLYWYCH LLYWYCH KERSON K	8*4001 8*4001 4*0304 4*0301 4*0201 4*0201 4*0201 4*0201 4*0201 8*0001 8*0001 8*0001 8*4000	24.9 10.6 102.5 110.8 110.8 10.6 6.3 6.9 6.9 8.9 10.9 482.8.2 754.6 204.5 16.9 43.4,40.0 2016.3 62.2 1751.7 87.0 22.0 140.5 16.3 48.0 7.2 59.8 32.6.7 190.8.4 21.9 22.0 140.5 16.3 17.2 19.8 17.2	3.0 2.6 0.2 0.7 8.3 6.2 6.2 6.2 6.2 6.2 8.3 8.4 NA 1.7 0.1 0.1 0.1 13.1 2.6,0.4 0.2 0.9 0.4 1.0 1.0 1.0 1.0 0.2 0.9 0.4 1.0 0.2 0.2 0.2 0.0 0.0 0.0 0.0 0.0 0.0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0	66 66 100 100 100 100 100 100 10	C C C C C C C C C C C C C C C C C C C	4.40 Not detected IN-y only IN-y only 2.87 3.01 IN-y only 0.71 Basal only IN-y only IN	IFN-y/TN-eq/L-18 only IFN-y/TN-eq/L-18 only IFN-y/TN-eq/L-18 only S.02 S.02 IFN-y/TN-eq/L-18 only
KRTLA,KFT JB KRTLA	0.12756,00033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12757 0.13562 0.204HC3 0.13562 0.204HC3 0.13562 0.204HC3 0.13562 0.204HC3 0.13562 0.204HC3	734.743 734.743 734.743 734.743 860.868 877.885 1347.1355 1460.1488 1140.1488 1517.1525 1291.299 1158.1167 13.18/166 159.33 747.433/7073,7479.433/885.888 109.117 158.166 139.338 144.152,412.470 129.298 156.58 159.58 159.58 155.156 159.516 159.555 155.550 155.55	REAMARSVEL ATHWTLERL LLYPPUH GEVKGFLUV VUDTSVARV KLSEMVTL KLSEMVTL KLSEMVTL KLSEMSTL GLDERSGLL TLALS/KLL RRODHVTL KLNOFSGLK THNFLVEL EPSTG/RPL EEPNINVGL EEVIL	8*4001 (*03364 4*03301 8*4001 4*0201 4*0201 4*0201 4*0201 4*0201 8*400 8*40018 8*40018 8*400000000000000000000000000000000000	24.9 105 105 105 105 63 69 105 63 69 105 105 105 105 105 105 105 105	3 3 6 2 6 0 2 0 6 0 7 0 7 8 3 6 2 6 2 8 3 6 2 6 2 6 2 6 2 6 2 6 3 6 2 6 2 6 2 6 3 6 2 6 2 6 2 6 3 6 2 6 2 6 2 6 2 6 3 6 2 6 2 6 2 6 2 6 2 6 2 6 2 6 2	0 0 0 0 0 0 0 0 0 0 0 0 0 0	66 66 100 100 100 100 100 100 10	C C C C C C C C C C C C C C C C C C C	4.40 Not detected IN-V only 2.37 3.01 IN-V only 3.01 IN-V only IN-	IFN-γ(ThF-γ(L-1)) only IFN-γ(ThF-γ(L-1)) only 3.6 Y-(ThF-γ(L-1)) only 3.6 4.39 IFN-γ(ThF-γ(L-1)) only 0.54 3.00 IFN-γ(ThF-γ(L-1)) only IFN-γ(ThF-γ(L-1)) only IFN-γ(ThF-γ(L-1)) only IFN-γ(ThF-γ(L-1)) only 1.09 6.78 3.49 IFN-γ(ThF-γ(L-1)) only IFN-γ(ThF-γ(L-1)) only 2.77 4.21 IFN-γ(ThF-γ(L-1)) only 2.70 4.21 IFN-γ(ThF-γ(L-1)) only 5.5 3.31 IFN-γ(ThF-γ(L-1)) only IFN-γ(ThF-γ(L-1)) only IFN-γ(ThF-γ(L-1)) only IFN-γ(ThF-γ(L-1)) only
GF1A,KF18 GF1A NCUROD1 NFCILI NFCILION NFCILI SCSC CSCR1 CSCR1 POSI PORA POPH POPEN PIPEN PIERN/PTENN PIERN/PTENN PIERN/PTENN PIERN/PTENN <td>012756,060333 012756 012756 012756 012756 012755 012755 012755 012755 012755 012755 012755 0201452 0201452 0201452 0201452 0201452 0201452 0201452 0201452 0201452 020145 02014 020145 020145 020145 020145 020145 020145 02014 020145 02014 020145 02014</td> <td>724.743 724.743 726.805 860.868 877.885 1347.1355 1440.1488 1480.1488 1480.1488 151.71.525 131.230 131.230 131.231 134.1162 131.231.23 135.235 135.23 1</td> <td>REAMARYEL ALTWYLER LLTYYPUH LLTYYPUH KESMAYT K</td> <td>8°4001 (°0304 4°0303 4°0303 4°0201 4°0201 4°0201 4°0201 4°0201 8°4000 8°4000 8°4000 8°4000 8°4000 8°4000 8°4000 8°4000 8°4000 8°</td> <td>24.9 10.6 102.5 110.8 110.8 10.6 6.3 6.9 0.9 10.9 482.8.2 754.6 204.5 16.9 43.4,40.0 201.6.3 62.2 1751.7 87.0 22.0 140.5 15.3 45.0 7.2 55.8 32.6.7 190.8.4 213.0 84.2 17.0</td> <td>3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.4 0.7 0.1 0.7 0.1 0.7 0.1 0.7 0.1 0.7 0.1 1.7 0.1 0.2 0.9 0.4 1.0 2.5 1.0 2.1 0.1 3.5 1.6 4.7</td> <td>0 0 0 0 0 0 0 0 0 0 0 0 0 0</td> <td>66 66 100 100 100 100 100 100 10</td> <td>C C C C C C C C C C C C C C C C C C C</td> <td>4.40 Not detected IN-y only 2.37 2.67 3.01 IN-y only 0.71 Basal only IN-y only 1.23 IN-y only 1.23 IN-y only 1.23 IN-y only 1.23 IN-y only 1.25 IN-y only 1.25 IN-y only 1.25 IN-y only 1.25 IN-y only IN-y only 1.25 IN-y only Basal only IN-y only I</td> <td>IFN-r/TN-eq/L-18 only IFN-r/TN-eq/L-18 only IFN-r/TN-eq/L-18 only S02 S02 IFN-r/TN-eq/L-18 only IFN-r/TN-eq/L-18 only</td>	012756,060333 012756 012756 012756 012756 012755 012755 012755 012755 012755 012755 012755 0201452 0201452 0201452 0201452 0201452 0201452 0201452 0201452 0201452 020145 02014 020145 020145 020145 020145 020145 020145 02014 020145 02014 020145 02014	724.743 724.743 726.805 860.868 877.885 1347.1355 1440.1488 1480.1488 1480.1488 151.71.525 131.230 131.230 131.231 134.1162 131.231.23 135.235 135.23 1	REAMARYEL ALTWYLER LLTYYPUH LLTYYPUH KESMAYT K	8°4001 (°0304 4°0303 4°0303 4°0201 4°0201 4°0201 4°0201 4°0201 8°4000 8°4000 8°4000 8°4000 8°4000 8°4000 8°4000 8°4000 8°4000 8°	24.9 10.6 102.5 110.8 110.8 10.6 6.3 6.9 0.9 10.9 482.8.2 754.6 204.5 16.9 43.4,40.0 201.6.3 62.2 1751.7 87.0 22.0 140.5 15.3 45.0 7.2 55.8 32.6.7 190.8.4 213.0 84.2 17.0	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.4 0.7 0.1 0.7 0.1 0.7 0.1 0.7 0.1 0.7 0.1 1.7 0.1 0.2 0.9 0.4 1.0 2.5 1.0 2.1 0.1 3.5 1.6 4.7	0 0 0 0 0 0 0 0 0 0 0 0 0 0	66 66 100 100 100 100 100 100 10	C C C C C C C C C C C C C C C C C C C	4.40 Not detected IN-y only 2.37 2.67 3.01 IN-y only 0.71 Basal only IN-y only 1.23 IN-y only 1.23 IN-y only 1.23 IN-y only 1.23 IN-y only 1.25 IN-y only 1.25 IN-y only 1.25 IN-y only 1.25 IN-y only IN-y only 1.25 IN-y only Basal only IN-y only I	IFN-r/TN-eq/L-18 only IFN-r/TN-eq/L-18 only IFN-r/TN-eq/L-18 only S02 S02 IFN-r/TN-eq/L-18 only
KRT-LA, KRT JB KRT-LA	0.12756,00033 0.12756 0.1276 0.13862 0.1070 0.13862 0.1070 0.13862 0.1070 0.10869 0.1070 0.10869 0.1070 0.10869 0.1070 0.10869 0.1070 0.10869 0.1070 0.10869 0.1070 0.10869 0.1070 0.10869 0.1086 0.108 0.10 0.10 0.10 0.10 0.10 0.10 0.1	734.743 724.743 724.743 726.805 860.868 877.885 1347.1355 1480.1488 1480.1488 1517.1525 7291.299 113.121/165.158,13.12/458.490 517.525 705.714 109.117 153.166 109.117 153.166 109.117 153.166 109.128 259.298	REAWARSVEL ATHWYTLERL LLYPPVLWH GEVKGFLUV VLDTSVARV KLSRMYTL KLSRMYTL KLSRMYTL KLSRMSTL KLSRMSTL KLSRMSTL ESCHERR BEDKARNAF ESDRARRA HEYNARNGI SLAGTRPL EEPNRYQL EEPN	8*4001 (*03304 4*03301) 8*4001 4*02010 4*02010 4*02010 4*02010 8*400100 8*40000000000000000000000000000000000	24.9 10.6 1925.8 110.8 6.3 6.9 6.9 4.0 10.9 4.0 2.0 10.5 10.5 10.5 2.0 10.5 10.5 2.0 2.0 10.5 10.5 2.0 2.0 10.5 10.5 2.0 2.0 10.5	3 5 6 2 6 0 2 0 6 0 7 0 7 0 7 8 3 6 2 6 2 6 2 0 7 0 7 0 1 1 7 1 7 1 1 1 0 1 0 1 0 1 0 2 6 0 4 0 2 0 9 0 4 1 0 1 1 1 0 1 1 1 0 1 1 1 1 1 0 0 1 1 1 1	0 0 0 0 0 0 0 0 0 0 0 0 0 0	660 1000 1	C C C C C C C C C C C C C C C C C C C	4.40 Not detected IN-V only 2.37 3.01 IN-V only 0.71 Basal only IN-V only IN	IFN γ(TDF-γ(L-1)) only IFN γ(TDF-γ(L-1)) only 3.6 Y 1FN γ(TDF-γ(L-1)) only 3.6 4.39 IFN γ(TDF-γ(L-1)) only 0.54 3.00 IFN γ(TDF-γ(L-1)) only 2.77 4.21 IFN γ(TDF-γ(L-1)) only P(H γ(TDF-γ(L-1))) only 2.70 4.21 IFN γ(TDF-γ(L-1)) only 6.35 3.31 IFN γ(TDF-γ(L-1)) only
NRTA_KPTB NRTA NRTA NRTA OFTA NCDA NEURODI NECIT NOV/TOTMANY/SCGS PCSID PCSID<	012756,00033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 013522 00UHC3 00U	724.743 724.743 726.805 860.868 877.885 1347.1355 1440.1488 1480.1488 1480.1488 151.715.25 291.299 13-142/164.154,113,14,488.490 13-12,152.55 706.714 19-130,165 117 154.165 117 154.165 117 154.165 159 159 159 159 159 159 159 159 159 15	REAMANYEL ATHWITERL LLYPPUH SCHORTWICK KISMANT	8*4001 (*0304 4*0301 4*0201 4*0201 4*0201 4*0201 8*4001 8*4001 8*4001 8*4001 (*0304 4*0201 (*0304 8*4001 8*4000000000000000000000000000000000000	24.9 10.6 1025.8 110.8 110.8 6.3 6.9 10.9 482.8.2 754.6 204.5 16.9 43.4,40.0 2016.3 62.2 1731.7 87.0 22.0 140.5 16.3 48.0 7.2 59.8 32.6.7 1908.4 213.0 84.2 17.0 37.7 1513.4	3.0 2.6 0.2 0.6 0.7 8.3 6.2 6.4 NA 1.7 0.1 0.7 0.1 0.7 0.1 0.2 0.2 0.4 1.0 4.2 1.1 100 0.1 2.5 1.0 2.1 0.1 3.5 1.6 4.7 0.4	0 0 200 60 40 40 60 60 200 80 800 60 1000 60 60 0 1000 100 1000 200 800 0 1000 200 800 0 1000 200 200 800 1000 1000 1000 0 1000 0	660 100 100 100 100 100 100 100	C C C C C C C C C C C C C C C C C C C	4.40 Not detected	IFN-Y[TN-eql(1-1)] only IFN-Y[TN-eql(1-1)] only IFN-Y[TN-eql(1-1)] only 3.6 5.02 IFN-Y[TN-eql(1-1)] only -0.54 3.00 IFN-Y[TN-eql(1-1)] only
KF1A,KF1B KF1A KF1A KF1A KF1A KF1A KF1A KF1A KF1A KF1A MCDAD NCUROD1 NPCI11 NPC/TFTRN,NPV/SCGS PCSK1 PCSK1 PCSK1 PCSK2 PCSK1 PCSK1 PCSK1 PCSK2 PCSK1 PCSK2 PCSK1 PCSK2 PCSK1 PCSK2 PCSK3 PCSK3 PCSK4 PCSK2 PCSK1 PCSK2 PCSK3 PCSK4 PCSK5 PCSK1 PCSK2 PCSK3 PCSK4 PCSK5 PCSK5 PCSK5 PCSK5 PCSK5 PCSK5 PCSK6	0.12756,00033 0.12756 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.1275 0.127 0.127 0.1284 0 0.127 0.1284 0 0.127 0.1284 0 0.127 0.1284 0 0.127 0.1284 0 0.127 0.1284 0 0.127 0.1284 0 0.127 0 0.128 0 0.127 0 0.128 0 0.127 0 0.128 0 0 0.127 0 0.128 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	734.743 724.743 724.743 726.805 860.868 1347.1355 1480.1488 1480.1488 1517.1525 291.299 1158.1167 13.12/165.158,13.12/458.490 517.525 705.714 30.388 144.152,412.420 289.728 50.58 247.245 259.728 50.58 247.255 151.146 50.595 155.154 50.59 155.154 50.59 155.154 50.59 155.154 50.59 155.154 50.59 155.154 50.59 155.15 50.5	AEJAMUSVEL ATHWTLERL LLYPPUH GEVKGFLUV VUDTSVAVV KLSEMVTL KLSEMVTL KLSEMVTL KLSEMSTL SIGLERKRP AEFENTVAF GIDLESGLL TLASL/KLL REGOLIVTL KLNWFSQLK TTNRFLVEL EPSTQ/RPL EEPNRVQL	8*4001 (*03304 4*03301) 8*4001 4*02010 4*02010 4*02010 4*02010 8*400000000000000000000000000000000000	24.9 10.6 1925.8 119.8 10.6 6.3 6.9 6.3 19.9 48.2.2 75.4.6 20.4.5 20.4.5 20.4.5 20.4.5 20.5	3 5 6 2 6 0 2 0 6 0 7 0 7 8 3 6 2 6 2 17 0 1 17 0 1 10 17 10 10 10 11 13 1 2 6 0 4 0 2 0 4 10 0 1 10 11 11 10 0 7 10 11 11 10 0 7 10 11 11 10 10 0 7 10 11 11 10 10 10 10 10 10 10	200 200 400 400 400 400 400 400	660 1000 1	C C C C C C C C C C C C C C C C C C C	4.40 Not detected	IFN-y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only 1FN-y/TM-zy/L-13 only 1S0 4.39 IFN-y/TM-zy/L-13 only 0.51 3.00 IFN-y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only 1.77 4.71 IFN-y/TM-zy/L-13 only 2.77 4.21 IFN-y/TM-zy/L-13 only 2.73 3.49 IFN-y/TM-zy/L-13 only Y/TM-zy/L-13 only Y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only Y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only IFN-y/TM-zy/L-13 only Y/TM-zy/L-13 only
KF1A,KF18 KF1A KF1A KF1A KF1A KF1A KF1A GF1A GF1A GF1A GF1A GF1A KF1A GF1A GF1A GF1A GF1A GF1A KF1A MF3A NCUROD1 NF2 NFCI1 PK2 PCSA1 PCSA2 PCSA2 PK3A3 PCSA2 PK3A3 PMA2A PMA2A PMA2A PMA2A PMA2A PMA2A PFRN PFPRN PFPRN PFPRN PFPRN PFPRN PFPRN PFPRN SC63 SC65 SC65 SC65	012756,06033 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 012756 013562 0100 0100 0100 0100 0100 0100 0100 01	724.743 724.743 726.805 860.868 860.868 1347.1355 1347.1355 1480.1488 1517.1555 231.239 131.81/167 131.81/167 131.81/167 131.81/167 131.81/167 131.81/167 131.81/167 132.818 148 132.81 134.132.142.0 135.156 135.15 135.156 135.15 135.156 135.15 135.156 135.15 135.156 135.15 135.156 135.15 135.156 135.15 135.15 135.156 135.15	REAMANYEL REAMANYEL ATHWYELER LLYPYELH RESKOFLAV VIDTSVARV KISSMAYT KISSMAY	8*4001 (*0304 4*0301 4*0201 4*0201 4*0201 4*0201 4*0201 4*0201 (*0304 4*0201 (*0304 4*0201 (*0304 4*0201 (*0304 4*0201 (*0304 4*0201 8*4000000000000000000000000000000000000	24.9 10.6 102.5 110.8 110.8 110.8 6.3 6.9 110.9 482.8.2 754.6 204.5 16.9 43.4,40.0 2016.3 62.2 1731.7 87.0 22.0 140.5 16.3 44.0 7.2 59.8 23.6.7 190.8.4 23.7 190.8.4 23.7 153.4 153.4 153.8	3.5 2.6 0.2 0.6 0.7 6.2 8.3 6.2 NA 1.7 0.1 1.3.1 2.6,0.4 0.2 0.4 1.0 4.2 1.1 100 0.1 2.5 1.0 2.5 1.0 2.1 0.1 3.5 1.6 4.7 4.7 4.7 4.7 0.2 1.7	200 200 400 400 400 400 400 400	660 100 100 100 100 100 100 100	C C C C C C C C C C C C C C C C C C C	4.40 Not detected	IFN-Y(TN-cyl):15 only IFN-Y(TN-cyl):15 only 3.66 5.02 4.39 IFN-Y(TN-cyl):15 only 0.54 3.00 IFN-Y(TN-cyl):15 only IFN-Y(TN-cyl):15 only IFN-Y(TN-cyl):15 only IFN-Y(TN-cyl):15 only IFN-Y(TN-cyl):15 only IFN-Y(TN-cyl):10 only IFN-Y(TN-cyl):10 only IFN-Y(TN-cyl):10 only 2.77 4.71 IFN-Y(TN-cyl):10 only 2.76 3.83 IFN-Y(TN-cyl):10 only 6.35 3.31 IFN-Y(TN-cyl):10 only 0.71/16-cyl):10 only 0.71/16-cyl):10 only 0.71/16-cyl):10 only 0.71/16-cyl):10 only 0.71/16-cyl):10 only 0.71/17-cyl):10 only 0.71/17-cyl):10 only 0.71/17-cyl):10 only 0.71/17-cyl):10 only 0.71/17-cyl):10 only 0.71/17-cyl):10 only 0.72 0.72 0.72 0.72
KF1A,KF1B KF1A MV03A NEUROD1 NPC111 NPC/TFTRN, NP/SCGS PCSK1 PCSK1 PCSK2 PCS10 PK333 PMMA2 PMMA2 PMMA2 PPRN PTPRN SCG5 SCG5 SCG5 SCG5 SCG5 SCG5 SCG5	0.12756,00033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12757 0.13862 0.2014C 0.13862 0.2014C 0.13862 0.2014C 0.13862 0.2014C 0.1380 0.1270 0.1380 0.1270 0.128 0.127 0.128 0.12 0.128 0.127 0.128 0.127 0.128 0.127 0.128 0.127 0.128 0.127 0.128 0.127 0.128 0.127 0.128 0.127 0.128 0.127 0.128 0.12 0.128 0.127 0.128 0.12 0.128 0.12 0.12 0.12 0.12 0.12 0.12 0.12 0.12	734.743 724.743 724.743 724.743 726.805 860.868 1347.1355 1480.1488 1480.1488 1517.1525 291.299 113.12/165.168,13.12/485.490 517.525 705.714 30.38 479.483/70.73,479.483/885.888 109.117 158.166 380.388 144.152,412.420 289.298 50.58 289.298 50.58 5	AEAWARSVEL ATHWYTLERL LLYPPVLWH GEYKGFLUV VLDTSVARV KLSEMVTL KLSEMVTL KLSEMVTL KLSEMVTL KLSEMSTL KLSEMSTL KLSEMSTL EGDERSGL EFKRVAF GLOLRSGL EFFNRVQL EEPRRV	8*4001 (*03344 4*03301) 8*4001 4*02010 4*02010 4*02010 4*02010 4*02010 4*02010 4*02010 (*03304 8*40010 4*02010 (*03304 8*40010 8*40010 8*40010 8*40010 8*40010 8*40010 8*40010 8*40010 8*40010 8*40010 4*020000000000000000000000000000000000	24.9 10.6 1925.8 119.8 10.6 6.3 6.9 6.9 48.2 204.5 204.	3 2 6 2 2 6 0 2 0 6 0 7 8 3 6 2 6 2 10 10 0 7 0 1 13 1 2 6 0 4 0 2 0 4 10 0 1 10 0 13 1 13 1 2 6 0 4 0 2 0 4 10 0 1 10 0 10 1 13 1 13 1 2 6 0 4 0 2 10 0 10 1 10 0 10 0	200 200 400 400 400 400 400 400	660 1000 1	C C C C C C C C C C C C C C C C C C C	4.40 Not detected Not detec	IFN-y(TNI-sy(L-1)) only IFN-y(TNI-sy(L-1)) only IFN-y(TNI-sy(L-1)) only 1.4.2 1.4.3 1.5.4 1.5.5 1.5.7
KF1A,KF18 KF1A KF1A KF1A KF1A KF1A KF1A GF1A GF1A GF1A GF1A GF1A KF1A GF1A GF1A GF1A KF1A MC3A NEUROD1 NPC NEURO1 NPC11 VPCSTPRN/PCSCS PCSR/ PCSR1 PCSR2 PCSR2 PCSR2 PG10 PEG10 PFR38, PFR31 PPMA2 PPMA2 PPMA2 PPMA2 PPMA2 PPMA PPMA2 PPMA PPMA2 PPMA PPMA2 PPMA PPMA2 PPMA PPMA2 PPMA PPMA2 PPMA1 PPMA2 PPMA2 PPMA2 PPMA2 PPMA3 PPMA2 PPMA3 PPMA1 PPMA2 PPMA1 PPMA2 PPMA2 PPMA3 PPMA3	012756,00033 012756 01275 012756 01275 012756 01275 012756 01275 01	724.743 726.805 860.868 860.868 1347.1355 1440.1488 1480.1488 1480.1488 1517.1525 231.239 131.81/165.458,13.18/488.4590 151.525 706.714 98.38 479.431/073,479.483/885.488 109.117 158.166 109.117 158.166 109.137 158.164 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.144 158.258 158.145 158.	REAMANYEL REAMANYEL ATHWYELER LIYPYEW KISSMAT	8*4001 (*0304 **0301 **0304 **0301 **0201 **01 **	24.9 10.6 102.5 110.8 110.8 110.6 6.3 6.9 110.9 423.2 754.6 204.5 16.9 43.4,40.0 2016.3 62.2 1731.7 87.0 22.0 140.5 16.3 45.0 72.2 1908.4 23.6,7 1908.4 213.0 84.2 17.0 15.3 84.2 17.0 15.3 15.3 15.3 15.5 1	3 5 6 2 6 0 2 0 6 0 7 8 3 6 2 NA 1.7 0.1 1.3.1 2.6,0.4 0.2 0.9 0.4 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0	200 200 400 400 400 400 400 400 400 400	660 100 100 100 100 100 100 100	C C C C C C C C C C C C C C C C C C C	4.40 Not detected Not detected Not detected Not detected Not detected Not detected 2.37 0.01 0.01 0.01 0.01 0.01 0.01 0.01 0.0	IFA-Y(ThE-g/L-1) only IFA-Y(ThE-g/L-1) only 1FA-Y(ThE-g/L-1) only 3.66 5.70 4.30 IFA-Y(ThE-g/L-1) only 0.54 3.00 IFA-Y(ThE-g/L-1) only IFA-Y(ThE-g/L-1) only IFA-Y(ThE-g/L-1) only IFA-Y(ThE-g/L-1) only IFA-Y(ThE-g/L-1) only 1.09 6.78 3.49 IFA-Y(ThE-g/L-1) only 2.77 4.21 IFA-Y(ThE-g/L-1) only 2.76 4.31 IFA-Y(ThE-g/L-1) only 6.35 3.31 IFA-Y(ThE-g/L-1) only 169 9.71 133 IFA-Y(ThE-g/L-1) only 169 170-71 161 171 180 180 190 191 191 191 192 191 191
KRF1A, KRF1B KRF1A NCUROD1 NPCL11 PCSK1 PCSK1 PCSK2 PCSK2 PCSK3 PCSK3 PR3383 PMMA2 PMMA2 PRPH PRPN PTPRN PTPRN <t< td=""><td>0.12756/06033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12757 0.13562 0.1270 0.13562 0.1070 0.13562 0.1070 0.13562 0.1070 0.1356 0.1070 0.1350 0.1070 0.135 0.101 0.1580 0.111 0.158 0.111 0.158 0.111 0.158 0.111 0.158 0.111 0.158 0.111 0.158 0 0.111 0.15 0 0.11 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0</td><td>724.743 724.743 726.805 860.868 877.885 1347.335 1480.1488 1480.1488 1140.1488 1157.1525 291.299 113.12/165.163,1.12/488.490 517.525 705.714 109.117 158.166 130.388 144.152,412.412 282.298 25.58 25.</td><td>AEAWARSVEL ATHWTERL LLYVPUH LLYVPUH LLYVPUH LLYVPUH LLYVPUH LLYVPUH LLYVPUH SCHOLTS SC</td><td>8*4001 (*03344 4*03301 8*4001 4*0201 4*0201 4*0201 4*0201 8*4001 8*4001 8*4001 6*0304 4*0201 (*03344 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 4*0201 8*4001 8*4001 4*0201 8*4001 4*0201 8*4001 4*0200</td><td>24.9 105 1055 1198 1065 63 69 69 69 69 69 69 82 2045 2058</td><td>336 02 02 03 83 62 83 62 83 62 83 62 83 62 83 62 83 83 62 83 83 62 83 83 83 83 83 83 83 83 83 83 83 83 83</td><td>200 200 400 400 400 400 400 400</td><td>660 1000 1</td><td>C C C C C C C C C C C C C C C C C C C</td><td>4.40 Not detected Not detected Not detected Not detected Not detected 2.37 2.87 3.01 Not detected Not detected Not detected Not detected 1.33 Not detected 2.04 4.33 Not detected 4.33 Not detec</td><td>IFA-y(Thi-a,2L-1) only IFA-y(Thi-a,2L-1) only IFA-y(Thi-a,2L-1) only 1.4 0.54 4.59 IFA-y(Thi-a,2L-1) IFA-y(Thi-a,2L-1) only IFA-y(Thi-a,2L-1) only <t< td=""></t<></td></t<>	0.12756/06033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12757 0.13562 0.1270 0.13562 0.1070 0.13562 0.1070 0.13562 0.1070 0.1356 0.1070 0.1350 0.1070 0.135 0.101 0.1580 0.111 0.158 0.111 0.158 0.111 0.158 0.111 0.158 0.111 0.158 0.111 0.158 0 0.111 0.15 0 0.11 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	724.743 724.743 726.805 860.868 877.885 1347.335 1480.1488 1480.1488 1140.1488 1157.1525 291.299 113.12/165.163,1.12/488.490 517.525 705.714 109.117 158.166 130.388 144.152,412.412 282.298 25.58 25.	AEAWARSVEL ATHWTERL LLYVPUH LLYVPUH LLYVPUH LLYVPUH LLYVPUH LLYVPUH LLYVPUH SCHOLTS SC	8*4001 (*03344 4*03301 8*4001 4*0201 4*0201 4*0201 4*0201 8*4001 8*4001 8*4001 6*0304 4*0201 (*03344 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 4*0201 8*4001 8*4001 4*0201 8*4001 4*0201 8*4001 4*0200	24.9 105 1055 1198 1065 63 69 69 69 69 69 69 82 2045 2058	336 02 02 03 83 62 83 62 83 62 83 62 83 62 83 62 83 83 62 83 83 62 83 83 83 83 83 83 83 83 83 83 83 83 83	200 200 400 400 400 400 400 400	660 1000 1	C C C C C C C C C C C C C C C C C C C	4.40 Not detected Not detected Not detected Not detected Not detected 2.37 2.87 3.01 Not detected Not detected Not detected Not detected 1.33 Not detected 2.04 4.33 Not detected 4.33 Not detec	IFA-y(Thi-a,2L-1) only IFA-y(Thi-a,2L-1) only IFA-y(Thi-a,2L-1) only 1.4 0.54 4.59 IFA-y(Thi-a,2L-1) IFA-y(Thi-a,2L-1) only IFA-y(Thi-a,2L-1) only <t< td=""></t<>
NRTA_KYT38 NRTA PRA	012756,060333 012756 01275 012756 01275 012756 01275 012756 01275 012756 01275 0127 01275 0127 0127 0127 0127 0127 0127 0127 0127	734-743 778-705 860-868 1347-1355 1440-1488 1140-1488 1157-1525 123-1239 1159-1167 13-127146-1463-134/488-490 15-525 706-714 105-117 158-1166 105-117 158-166 105-117 158-166 105-117 158-165 105-125	REAMANYEL REAMANYEL ATHWYELER LLYPYEVH GEWGELRY VIDTSVARY KISSMYTL	8*4001 (*0304 4*0301 4*0201 4*0201 4*0201 4*0201 4*0201 4*0201 (*0304 4*0201 (*0304 4*0201 (*0304 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 8*4001 4*0201 4*	24.9 105.6 1975.8 1108.4 1066.6 6.9 106.6 6.9 106.6 1075.8 106.6 1075.8 10	3 3 6 2 8 0 2 0 6 0 6 0 7 7 7 8 3 6 2 8 3 6 2 8 3 6 2 8 3 6 2 8 3 17 0 1 0 1 0 1 0 1 0 1 0 1 0 2 0 0 0 4 0 2 0 9 0 4 10 0 1 10 0 1 11 10 0 1 0 1	0 0 200 660 400 660 400 600 400	660 1000 1	C C C C C C C C C C C C C C C C C C C	4.40 Not detected	IFN-yTM-s-yL-15 only IFN-yTM-s-yL-15 only IFN-yTM-s-yL-15 only 3.65 Y-YTM-s-yL-16 only 0.54 3.00 IFN-yTM-s-yL-16 only 2.77 4.21 IFN-yTM-s-yL-16 only 7.76 3.83 IFN-yTM-s-yL-16 only 6.55 3.31 IFN-yTM-s-yL-16 only (6.55 3.31 IFN-yTM-s-yL-16 only (1.63 IFN-yTM-s-yL-16 only (1.61 0.78 Basal only 1.60 1.61 1.61 IFN-yTM-s-yTM-s-yTM-s-yTM-s-yTM-s
KF1A,KF18 GF1A G	0.12756/00033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12757 0.13562 0.1270 P01301/005408 P01301/005408 P01301/005408 P01301/005408 P01301/00540 0.1570 0.1580 0.1570 0.1580 0.1570 0.158 0.158 0.158 0.158 0.158 0.158 0.158 0.158 0.158 0.158 0.158 0.158 0.15 0.158 0.15 0.158 0.15 0.158 0.15 0.15 0.15 0.15 0.15 0.15 0.15 0.15	734.743 734.743 734.743 734.743 660.868 677.865 660.868 1347.1355 1460.1484 1480.1484 1480.1484 151.745.75 151.154.1166 153.116/165.131.8/488.400 151.16/152.134.7479.443.7858.88 109.117 137.385 156.166 380.388 144.52.412.400 289.298 247.255 171.129 171.379 173.385 156.164 509.516 509.58 247.255 171.129 173.385 156.164 509.516 509.516 509.5975 575.580.708-711 150.129 156.174 158.192 158.194 154.194 155.155 155.155 155.155 155.1	AEAMANYEL ALTHYVTERL LLTYVPUH LLTYVPUH VIDTSANN VIDTSANN SEGNATIS	8*4001 (*0304 4*0301 4*0301 4*0301 4*0201 4*0201 4*0201 4*0201 4*0201 4*0201 5*0304 4*0201 5*0304 4*0201 5*0304 5*0301 5*0304 5*0301 5*05 5*05 5*05 5*05 5*05 5*05 5*05 5*	24.9 105 1075.8 119.8 105.6 6.3 6.9 119.8 105.6 6.3 6.9 NA 19.9 482.8 204.5 205.5 20	336 02 05 05 83 62 62 84 17 01 01 01 131 26,04 02 03 04 10 01 131 131 12 0,04 02 03 04 10 01 10 10 01 10 10 01 10 10 01 10 10	0 0 232 0 243 0 244 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 1000 0 1000 0 0 <	66 100	C C C C	4.40 Not detected Not detec	IPA-(Thi-q/L-1) only IPA-(Thi-q/L-1)
NRTA_KYT38 NRTA PEGIO PEGIO PEGIO PEGIO PRARAZ PMMA2 PMMA2 PRMA2 PRENOPTRN PRENOPTRN </td <td>012756,00033 012756 01275 012756 01275 012756 01275 0127 0127 0127 0127 0127 0127 0127 0127</td> <td>734.743 734.743 736.805 860.868 1347.1355 1460.1488 1140.1488 1157.1525 231.239 131.81/164-543.134/458-490 131.81/164-543.134/458-490 131.81/164-543.134/458-490 131.81/164-543.134/458-490 131.81/164 136.138 136.138 136.138 136.138 136.138 136.138 136.138 137.138 136.13</td> <td>KEAMADYEL ATHWYTLER. LLYPPYLH GEWGFLAV VIDTSVATV KISEMAYTL KISEMAY</td> <td>8*4001 6*4001 4*0301 4*0201 4*0201 4*0201 4*0201 4*0201 4*0201 4*0201 0*0304 4*0201 (*0304 (*0304 4*0201 (*0304</td> <td>24.9 24.9 105.6 1975.8 1108.2 1066.6 6.9 10.0 10.0</td> <td>3.5 2.6 0.2 0.6 0.7 0.7 0.8 3.3 6.2 6.2 NA 1.7 0.1 1.3.1 2.6,0.4 0.2 0.9 0.4 1.0 4.2 1.1 10.0 2.5 1.0 2.1 0.1 2.5 1.0 2.1 0.1 3.5 1.6 4.7 0.4 0.2 1.6 7.3 0.4 0.2 1.7 1.3.6 7.3 0.4 2.9 1.4</td> <td>0 0 220 0 400 <t< td=""><td>660 1000 1000 1000 1000 202 404 402 404 402 404 404</td><td>C C C C</td><td>4.40 Not detected Not detected</td><td>IFN-γ(Th = γ(L = 1)) only IFN-γ(Th = γ(L = 1)) only 3.6 5.0 4.30 IFN-γ(Th = γ(L = 1)) only 9.54 3.00 IFN-γ(Th = γ(L = 1)) only 9.54 3.00 IFN-γ(Th = γ(L = 1)) only 9.54 3.00 IFN-γ(Th = γ(L = 1)) only IFN-γ(Th = γ(L = 1)) only 1.09 6.78 3.49 IFN-γ(Th = γ(L = 1)) only 2.77 4.21 IFN-γ(Th = γ(L = 1)) only 2.70 4.21 IFN-γ(Th = γ(L = 1)) only 6.35 3.31 IFN-γ(Th = γ(L = 1)) only 6.35 3.31 IFN-γ(Th = γ(L = 1)) only -0.78 Buail ority NA 1.99 1.90 1.90 1.90 1.90 1.90 1.90 1.90 1.</td></t<></td>	012756,00033 012756 01275 012756 01275 012756 01275 0127 0127 0127 0127 0127 0127 0127 0127	734.743 734.743 736.805 860.868 1347.1355 1460.1488 1140.1488 1157.1525 231.239 131.81/164-543.134/458-490 131.81/164-543.134/458-490 131.81/164-543.134/458-490 131.81/164-543.134/458-490 131.81/164 136.138 136.138 136.138 136.138 136.138 136.138 136.138 137.138 136.13	KEAMADYEL ATHWYTLER. LLYPPYLH GEWGFLAV VIDTSVATV KISEMAYTL KISEMAY	8*4001 6*4001 4*0301 4*0201 4*0201 4*0201 4*0201 4*0201 4*0201 4*0201 0*0304 4*0201 (*0304 (*0304 4*0201 (*0304	24.9 24.9 105.6 1975.8 1108.2 1066.6 6.9 10.0 10.0	3.5 2.6 0.2 0.6 0.7 0.7 0.8 3.3 6.2 6.2 NA 1.7 0.1 1.3.1 2.6,0.4 0.2 0.9 0.4 1.0 4.2 1.1 10.0 2.5 1.0 2.1 0.1 2.5 1.0 2.1 0.1 3.5 1.6 4.7 0.4 0.2 1.6 7.3 0.4 0.2 1.7 1.3.6 7.3 0.4 2.9 1.4	0 0 220 0 400 <t< td=""><td>660 1000 1000 1000 1000 202 404 402 404 402 404 404</td><td>C C C C</td><td>4.40 Not detected Not detected</td><td>IFN-γ(Th = γ(L = 1)) only IFN-γ(Th = γ(L = 1)) only 3.6 5.0 4.30 IFN-γ(Th = γ(L = 1)) only 9.54 3.00 IFN-γ(Th = γ(L = 1)) only 9.54 3.00 IFN-γ(Th = γ(L = 1)) only 9.54 3.00 IFN-γ(Th = γ(L = 1)) only IFN-γ(Th = γ(L = 1)) only 1.09 6.78 3.49 IFN-γ(Th = γ(L = 1)) only 2.77 4.21 IFN-γ(Th = γ(L = 1)) only 2.70 4.21 IFN-γ(Th = γ(L = 1)) only 6.35 3.31 IFN-γ(Th = γ(L = 1)) only 6.35 3.31 IFN-γ(Th = γ(L = 1)) only -0.78 Buail ority NA 1.99 1.90 1.90 1.90 1.90 1.90 1.90 1.90 1.</td></t<>	660 1000 1000 1000 1000 202 404 402 404 402 404 404	C C C C	4.40 Not detected	IFN-γ(Th = γ(L = 1)) only IFN-γ(Th = γ(L = 1)) only 3.6 5.0 4.30 IFN-γ(Th = γ(L = 1)) only 9.54 3.00 IFN-γ(Th = γ(L = 1)) only 9.54 3.00 IFN-γ(Th = γ(L = 1)) only 9.54 3.00 IFN-γ(Th = γ(L = 1)) only IFN-γ(Th = γ(L = 1)) only 1.09 6.78 3.49 IFN-γ(Th = γ(L = 1)) only 2.77 4.21 IFN-γ(Th = γ(L = 1)) only 2.70 4.21 IFN-γ(Th = γ(L = 1)) only 6.35 3.31 IFN-γ(Th = γ(L = 1)) only 6.35 3.31 IFN-γ(Th = γ(L = 1)) only -0.78 Buail ority NA 1.99 1.90 1.90 1.90 1.90 1.90 1.90 1.90 1.
KF1A,KF18 GF1A G	0.12756/00033 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12756 0.12757 0.13562 0.1270 P01301/02649/01301/05568 P05508 P05509 0.1010 0.101 0.1010 0.101 0.1010 0.101 0.10	724.743 724.743 724.743 724.743 724.743 724.743 724.743 134.7135 1440.148 1480.148 1480.148 1480.148 1480.148 151.7153 151.7153 151.7153 151.7153 151.715 152.71 152.71 152.71 152.71 152.71 152.71 152.72 152.75 152.72 15	AEAMANYEL ALTWYTERL LLTWYCH LLTWYCH ALTWYTERL LLTWYCH ALTWYTERL LLTWYCH ALTWYTER ALT	8*4001 (*0304 4*0301 4*0301 4*0301 4*0201 4*0201 4*0201 4*0201 4*0201 5*000 5*00 5*	24.9 105 1075.8 1193.8 105.6 6.3 6.9 1193.8 105.6 6.3 6.9 NA 19.9 482.8 204.5 16.9 204.5 16.9 204.5 16.9 204.5 16.2 1731.7 87.0 2216.3 162.1 1731.7 87.0 1098.4 205.8 1098.4 205.8 1098.4 215.8 205.8 1098.4 215.8 205.8	336 226 02 05 83 62 62 NA 17 01 01 131 26,04 02 03 04 10 01 131 131 12 26,04 02 03 04 10 01 131 131 131 100 01 03 04 20 21 21 100 01 01 100 10 25 10 21 21 21 21 21 21 21 21 21 21 21 21 21	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	66 100 100 100 100 100 100 20 40	C C C C C C C C C C C C C C C C C C C	4.40 Not detected Not detec	IPA-(Thi-q.(L-1)) only IPA-(Thi-q.(L-1

Table S1. Related to Fig 1G, 2A-E. List of the 98 HLA-I-bound peptides identified in ECN90 β cells. The bioinformatics filters depicted in Fig. 1G were applied and the β -cellenriched (n=86; conventional and with PTMs), peptide splice (n=10) and mRNA splice (n=2) aa sequences identified are listed in the alphabetical order of their source proteins. HLA-I restrictions were assigned based on the top scores of predicted affinity and stability, as NetMHCstabpan determined bv (www.cbs.dtu.dk/services/NetMHCstabpan). The percent of positive biological replicates out of 5 analyzed are listed for IFN-y-treated, IFN- $\gamma/TNF-\alpha/IL-1\beta$ -treated and untreated (basal) ECN90 β cells. The last two columns display the log₂ median fold change (FC) in peptide content for IFN-y vs. basal and IFN-y/TNF- α /IL-1 β vs. basal conditions, respectively. The color codes indicate no change (<2 log₂ FC, either positive or negative; green), an increase for cytokine-treated conditions (\geq +2 log₂ FC; red) or a decrease for cytokine-treated conditions ($\leq -2 \log_2 FC$; blue). HLA-A2restricted peptides in bold red fonts were further analyzed for their recognition by CD8⁺ T cells, with those eventually validated as recognized by the naïve CD8⁺ T-cell repertoire (see Fig. 4) highlighted in grey. The remaining HLA-A2-restricted peptides were excluded because they were either length variants of peptides with better affinity scores or β-cell epitopes already described in the literature, i.e. PPI₂₋₁₀, PPI₆₋₁₄, PPI₁₅₋₂₄, PPI₂₉₋₃₈ (INS_{B5-14}), PPI₃₄₋₄₂ (INS_{B10-18}). The PTM peptides identified as originating from ubiquitous proteins are listed separately in Table S2. NA, not available.

Source protein(s)	Accession number(s)	Amino acid positions	Sequence	HLA restriction	Predicted affinity (nM)	Predicted stability (h)	IFN-γ (%)	IFN-γ/TNF-α/IL-1β (%)	Basal (%)	IFN-y log2 FC	IFN-γ/TNF-α/IL-1β log2 FC
AAMP	Q13685	150-159	ATGDM(+15.99)SGLLK	A*03:01	253.54	2.0	40	40	20	0.09	-0.63
ACTNA	043707	442-455	AFENRIM(+15.99)VSV	R*40:01	5.4	13.5	00	40	40	Basal only	Rasal only
AP1M2	Q9Y6Q5	30-38	IEHFM(+15.99)PLLV	B*40:01, B*49:01	263.3, 147.7	0.8, 0.9	40	0	20	1.52	Basal only
APOL6	Q9BWW8	85-93	AVISGVM(+15.99)SL	C*03:04, A*02:01	98.9, 22.3	1.3, 0.3	40	0	20	2.49	Basal only
ARMCX1	Q9P291	379-387	KEW(+3.99)DREILL	B*40:01	26.5	1.4	100	80	0	IFN-y only	IFN-γ/TNF-α/IL-1β only
ASXL1	099437	327-335	GEFTHEM(+15.99)QV SM(+15.99)FGAGITV	8*40:01 4*02:01	103.4	1.1	40	40	20	3.16	Basal only
BRD8	Q9H0E9	770-778	AEFQRDIM(+15.99)L	B*40:01	18.2	1.7	20	20	100	-3.14	-4.80
BUB1B	O60566	745-753	IEDRPM(+15.99)PKL	B*40:01	141.3	0.8	80	80	0	IFN-y only	IFN-γ/TNF-α/IL-1β only
C16orf62	Q7Z3J2	929-937	RTM(+15.99)VKTLEY	A*03:01	98.7	1.8	60	40	0	IFN-y only	IFN-γ/TNF-α/IL-1β only
CCT8	P14635 P50990	204-212 296-304	KVADM(+3.99)LVQV	A*02:01 A*03:01	122.4	2.0	40	0	20	U.45 IEN-y only	Not detected
CDC73	Q6P1J9	313-321	GTYHGM(+15.99)TLK	A*03:01	12.5	13.4	0	60	0	Not detected	IFN-y/TNF-α/IL-1β only
COPG1;COPG2	Q9Y678;Q9UBF2	307-317	AVRTLNKVAM(+15.99)K	A*03:01	227.0	1.0	100	0	0	IFN-γ only	Not detected
COX11	Q9Y6N1	244-252	AEDPRM(+15.99)IKV	B*40:01, B*49:01	1308.6, 1530.9	0.7, 0.6	60	80	60	0.44	-0.53
CINNA1 CINNB1	P35221 P35222	898-906 186-194	SEFKAM(+15.99)DSI HAIM(+15.99)RSDOM	C*03:04	29.9	1.0	60	40	40	0.21	-0.68
DEGS1	015121	83-91	SM(+15.99)TLAIHEI	A*02:01	66.9	4.5	40	0	20	0.38	Basal only
DES;INA;NEFH;NEFM;VIM	Q16352;P17661;P07197;P12036;P08670	377-387	REYQDLLNVKM(+15.99)	B*40:01	25.3	1.7	40	40	40	0.62	-0.30
DHX8	Q14562	981-989	AEFPLEPM(+15.99)L	B*40:01	9.6	1.8	80	20	80	2.68	-5.37
DSP	P15924 004637	2792-2802	SM(+15.99)VEDITGLRL	A*02:01 A*02:01	69.6	0.9	20	0	40	-0.71	Basal only Basal only
FAM136A	Q96C01	8-18	RVQEAVESM(+15.99)VK	A*03:01	331.7	5.7		0	40	Basal only	Basal only
FBXO21	094952	417-425	AM(+15.99)YPDQVQL	A*02:01	32.3	1.7	20	40	0	IFN-y only	IFN-γ/TNF-α/IL-1β only
G3BP1	Q13283	106-114	RRFM(+15.99)QTFVL	C*07:01	61.4	0.2	20	40	40	-4.14	-0.64
GNAS	Q5JWF2;P63092	752-762	AM(+15.99)SNLVPPVEL	A*02:01	41.0	1.9	100	100	40	3.89	3.26 Racal only
GNB1;GNB2;GNB3	P62879;P62873;P16520	328-337	ATGSW(+3.99)DSFLK	A*03:01	221.6	1.6	40	0	20	-2.21	Basal only
GPR162	Q16538	32-41	AW(+15.99)IILSISAK	A*03:01	3880.6	0.2	100	100	40	1.24	2.03
HAUS7	Q99871	196-203	PLDM(+15.99)QPLL	A*02:01	3427.1	0.4	0	60	0	Not detected	IFN-γ/TNF-α/IL-1β only
HID1 HIA-R	U8IV36	752-760	GTAM(+15.99)WFRTY	A*03:01	1730.8	0.2	20	40	20	-0.76 Basal only	0.70
HSF2	Q03933	156-164	SLW(+3.99)KEVSEL	A*02:01	30.3	1.9	40	40	40	-0.06	0.21
HSP90AA1;HSP90AA2P;HSP90AB1;HSP90AB2P	P08238;Q14568;Q58FF8;P07900	19-27	AEIAQLM(+15.99)SL	B*40:01	8.2	2.4	40	0	20	1.92	Basal only
HSPA14	Q0VDF9	56-66	RIRNISNTVM(+15.99)K	A*03:01	44.3	6.0	40	0	40	0.66	Basal only
HSPA4;HSPA4L	095757;P34932	169-177	RLM(+15.99)NETTAV	A*02:01	7.3	27.5	80	100	40	0.49	0.18
IRAK2	043187	45-53	KI M(+15.99)AVGGGLAV	A*03:01	28.9	0.9	40	80	20	2.76	0.78
IRF6	014896	235-243	KEYGQTM(+15.99)TV	B*40:01, B*49:01	99.7, 81.9	1.4, 2.0	20	20	40	-1.07	-2.56
KPNA1;KPNA5;KPNA6	O60684;P52294;O15131	432-440	VM(+15.99)DSKIVQV	A*02:01	9.7	4.3	20	60	40	0.66	0.16
KRT19	P08727;P08727	361-369	QEYQRLM(+15.99)DI	B*49:01	120.8	0.8	40	20	0	IFN-y only	IFN-γ/TNF-α/IL-1β only
MADD MED19	Q8WXG6	389-397	RIM(+15.99)PDDVWLV	A*02:01 A*03:01	2.7	9.8	40	80	40	1.07	U.80
MKI67	P46013	540-549	VM(+15.99)HTPPVLKK	A*03:01	17.5	2.7	20	40	20	1.49	1.02
MTOR	P42345	1189-1197	QIFIPM(+15.99)VNK	A*03:01	61.5	0.7	40	40	20	2.39	-0.39
NAA20	P61599	91-99	KLM(+15.99)ELLEEI	A*02:01	2.2	51.5	40	0	20	1.65	Basal only
NDN NOMO1-NOMO2-NOMO3	Q99608 015155:P69849:05IPE7	161-169	HTM(+15.99)EFALVK SEVI PTI W/+3 991V	A*03:01 B*40:01 B*49:01	44.1	2.5	40	40	40	Not detected	IFN-γ/TNF-α/IL-1β only
NRIP1	P48552	745-753	ALSEQILM(+15.99)V	A*02:01	5.9	12.1	100	40	0	IFN-y only	IFN-γ/TNF-α/IL-1β only
OSBPL9	Q96SU4	138-147	KLFDDKLQNC(+47.98)	A*02:01	188.0	1.5	40	100	40	3.08	0.57
PARP14	Q460N5	278-288	KVLDTIM(+15.99)ATKL	A*02:01	212.1	1.3	40	0	0	IFN-y only	Not detected
PCNA	P12004	197-205	KI M(+15.99)NEPVQL	A*02:01	5.3	2.3	40	0	20	2.59	0.15 Rasal only
PCSK9	Q8NBP7	194-202	REIEGRVM(+15.99)V	B*40:01	14.2	1.7	60	0	20	1.47	Basal only
PKLR;PKM	P14618;P30613	21-29	AM(+15.99)ADTFLEH	A*03:01	365.9	1.0	0	40	0	Not detected	IFN-γ/TNF-α/IL-1β only
PLXNA4	Q9HCM2	1538-1547	KNVPC(+47.98)SHRPK	NA	NA	NA	0	0	40	Basal only	Basal only
POMP PRKAR1A:PRKAR1B	Q9Y244 P10644·P31321	124-132 244-253	SEVM(+15.99)GEPHL KM(+15.99)YEEEISKV	A*02:01	34	1.1	40	100	40	1.30 IEN-y only	0.02 Not detected
PRKDC	P78527	1302-1311	AM(+15.99)HDIIAAEK	A*03:01	69.1	4.6	20	0	60	-2.66	Basal only
PSMA2	P25787	113-121	RVASVM(+15.99)QEY	A*03:01	553.4	0.6	20	60	20	0.29	0.84
PSMB9	P28065	15-24	GEVHTGTTIM(+15.99)	B*40:01	40.3	1.1	100	0	0	IFN-y only	Not detected
REC8	095072	184-194	RTM(+15.99)SEAALVKK AFPIRM(+15.99)IFI	A*03:01 NA	44.2 NA	17.1 NA	100	40	40 20	5 38	-0.75
RPAIN	Q86UA6	128-136	AEW(+3.99)EANPLI	B*40:01, B*49:01	20.1, 35.5	1.2, 1.2	20	40	20	-1.83	-1.16
RPL19	P84098	137-145	ILM(+15.99)EHIHKL	A*02:01	2.4	39.3	40	20	20	1.00	-1.98
RPL3 RPN2	P39023	324-333 619-627	GEVTNDFVM(+15.99)L	B*40:01	16.1	1.2	100	0	0	IFN-y only	Not detected
RP529	P62273	5-13	QLYW(+15.99)SH(+15.99)PRK	A*03:01	14.3	2.2	40	40 20	20	Basal only	-0.73
SEC11C	Q9BY50	180-189	AVM(+15.99)GAYVLLK	A*03:01	26.3	12.2	40	80	60	-0.57	-0.66
SEC11C	Q9BY50	24-32	KM(+15.99)NKRQLYY	A*03:01	11.9	1.5	80	80	20	1.65	0.04
SEC14L1	Q92503 D62341	32.42	QUDKVW(+3.99)QL KM(+15.99)OVATGPULK	A*02:01 A*03:01	3.9	5.2	20	40	20	Basal only	-0.50
SEPT9	Q9UHD8	492-500	REM(+15.99)IPFAVV	B*40:01	17.8	1.7	40	40	40	IFN-y only	IFN-y/TNF-α/IL-1β only
SKA2	Q8WVK7	3-11	AEVDKLELM(+15.99)	B*40:01	121.2	1.0	100	100	20	4.21	3.13
SLC25A3	Q00325	226-234	RQIPYTM(+15.99)M(+15.99)K	A*03:01	38.2	14.0	40	40	40	1.51	-0.82
SMCHD1	A6NHR9 008945	393-402	REIQDDM(+15.99)QTL RVM(+15.99)KAI VNR	B*40:01 A*03:01	251.0	1.4	80	20	40	Basal only 2 07	-2.08
STAT1	P42224	300-308	VLW(+3.99)DRTFSL	A*02:01	3.5	8.3	20	40	0	IFN-y only	IFN-y/TNF-α/IL-1β only
STAU1	095793	416-424	GILPM(+15.99)VPEV	A*02:01	8.3	7.2	20	0	40	-3.26	Basal only
STIP1	P31948	252-262	KELDPTNM(+15.99)TYI	B*40:01, B*49:01	125.7, 135.9	0.9, 0.9	60	40	20	4.21	1.40 Receiver http://www.com/com/com/com/com/com/com/com/com/com/
TGM6	095932	617-626	M(+15.99)VGVAVTVFV	A*02:01	040.8	1.0	0 20	0	40 20	-0,42	-0,17
THOC7	Q6I9Y2	76-84	REM(+15.99)ENYEKI	B*40:01, B*49:01	22.7, 92.7	1.0, 1.4	40	60	40	-0.35	-0.89
TIMM23;TIMM23B	014925;Q5SRD1	161-169	GTMTGM(+15.99)LYK	A*03:01	16.2	9.9	20	20	40	-0.44	-0.65
TJP2	Q9UDY2	80-88	QENDRVVM(+15.99)V	B*49:01	378.0	0.6	40	0	0	IFN-y only	Not detected
TMED9	098VK6	130-139	SIVESTIQIVI(+15.99)11 SLFAGGM(+15.99)I RV	A*02:01	46.8 9.9	0.4 6.0	20	40	40 20	-0.33	-0.47 Basal only
TNPO2	014787	215-223	IEH(+15.99)LFALAV	B*40:01, B*49:01	58.2, 101.9	1.2, 0.9	20	40	20	1.94	0.15
TOP2A;TOP2B	P11388;Q02880	168-177	KLC(+47.98)NIFSTKF	NA	NA	NA	40	0	0	IFN-y only	Not detected
TRANK1	015050	1987-1996	EAH(+15.99)FLQGVIL	NA	NA 25.C	NA	100	100	40	1.90	1.48
UBL5	098ZL1	60-68	YEIHDGM(+15.99)NL	B*40:01	25.b 13.6	1.2	40	60	20	U.19 IFN-y only	-0.21 Not detected
UBR2	Q8IWV8	703-711	SM(+15.99)M(+15.99)DPNHFL	A*02:01	3.6	5.7	40	0	0	IFN-y only	Not detected
UQCRQ	014949	3-11	REFGNLTRM(+15.99)	B*40:01	33.0	1.7	80	100	40	1.76	-0.10
USP11	P51784	611-619 621-629	M(+15.99)LFGHPLLV	A*02:01	3.1	7.9	20	20	40	2.22	-1.23
UTRN	P46939	414-422	RLHDVLM(+15.99)EL	A*02:01	5.8	11.7	40	40	40	1.06	-0.44
LITON	010000			0.840.04	10.0	2.4					4.60

Table S2. Related to Fig 1G. List of the 99 HLA-I-bound PTM peptides originating from ubiquitous proteins identified in ECN90 β cells. Data representation is the same as in Table S1. NetMHCstabpan predicted affinity and stability scores refer to the unmodified aa sequence.

Source protein(s)	Accession number(s)	Aminoacid positions	Sequence	HLA restriction	Predicted affinity (nM)	Predicted stability (h)	IFN-γ/TNF-α/IL-1β log2 FC
		2-10	ALWMRLLPL	A*02:01	26.7	5.6	Basal only
		2-12*	ALWMRLLPLLA	A*02:01	83.8	2.8	-2.16
		5-13	MRLLPLLAL	B*39:01	71.4	0.6	1.71
		5-13	M(+16)RLLPLLAL	B*39:01	71.4	0.6	Basal only
		15-24	ALWGPDPAAA	A*02:01	95.5	1.8	-3.26
INS (n=12)	R01209	15-24	ALW(+16)GPDPAAA	A*02:01	95.5	1.8	-3.58
1103 (11-12)	F01308	15-24	ALW(+4)GPDPAAA	A*02:01	95.5	1.8	Basal only
		17-24*	WGPDPAAA	NA	NA	NA	-3.87
		29-38	HLCGSHLVEA	A*02:01	521.1	1.5	6.64
		33-41	SHLVEALYL	B*39:01	283.8	0.4	IFN-γ/TNF-α/IL-1β only
		34-42	HLVEALYLV	A*02:01	3.3	17.8	Basal only
		57-67*	EAEDLQVGQVE	NA	NA	NA	0.13
		344-352	KMDQLAKEL	A*02:01	543.8	0.6	0.04
	P10645	344-354	KMDQLAKELTA	A*02:01	823.2	0.6	-1.68
	10045	381-390	YGFRGPGPQL	C*12:03, C*14:02	150.2, 139.9	0.2, 0.2	-0.37
		383-390	FRGPGPQL	B*39:01	380.7	0.7	-0.90
		791-800	DQKNGATHYW	NA	NA	NA	-1.08
KIF1A (n=3)	Q12756	1058-1066	EVTKSFIEY	A*25:01	370.6	1.3	-1.86
		1319-1327	NRVTGVYEL	B*39:01	19.8	0.7	-0.72
		136-144	DTAEFSREF	A*25:01	187.4	0.7	4.61
SCG5 (n=3)	P05408	186-195*	YLQGQRLDNV	A*02:01	44.8	2.2	1.77
		186-196	YLQGQRLDNVV	A*02:01	151.8	1.7	-1.19
		429-437	YFMSDTREE	C*14:02	1857.7	0.1	Basal only
CHGB (n=3)	P05060	429-437	YFM(+16)SDTREE	C*14:02	1857.7	0.1	IFN-γ/TNF-α/IL-1β only
		440-448	FLGEGHHRV	A*02:01	4.4	8.6	-3.35
PCSK2 (n=1)	P16519	26-34	ERPVFTNHF	C*14:02	3731.5	0.1	0.74
IAPP (n=1)	P10997	23-30	TPIESHQV	B*51:01	1423.2	1.1	1.84
SLC7A2 (n=1)	P52569	650-658	IFHEKTSEF	C*14:02	27.4	0.3	0.33
UNC13A (n=1)	Q9UPW8	1234-1243	DIISKDFASY	A*25:01	73.9	0.8	-0.58
IAPP/IAPP;PTPRN/IAPP	P10997/P10997;Q16849/P10997	15-17/5-10;596-598/5-10	VAL/KLQVFL	A*02:01, C*12:03	3599.1, 3763.0	0.5, 0.1	-0.58
SCG5/PCSK2;SLC30A8/PCSK2	P05408/P16519;Q8IWU4/P16519	113-115/565-570;168-170/565-570	YPD/RGTWTL	B*39:01	10.0	0.8	IFN-γ/TNF-α/IL-1β only
PTPRN/IAPP	Q16849/P10997	795-802/44-48	WQM(+16)VWESG/RLANF	B*39:01, C*14:02	1487.1, 1801.9	0.3, 0.2	1.84
WARS-035	WARS-035	18-26	SVAQAGVHW	NA	NA	NA	IFN-γ/TNF-α/IL-1β only

Table S3. Related to Fig 2F. List of the 33 HLA-I-bound peptides identified in primary human islets. The same bioinformatics filters used for ECN90 β cells (Fig. 1G) were applied, barring the inter-sample reproducibility and HLA-I enrichment filters due to the analysis of a single split islet preparation (cytokine-treated or not), without mock immunopurification. As sequences are listed according to the number of peptides identified for each source protein. HLA-I restrictions were assigned based on the predicted affinity and stability (NetMHCstabpan). The last column displays the log2 median FC in peptide content for the IFN- γ /TNF- α /IL-1 β *vs.* basal condition (color codes as in Table S1). Source proteins and/or peptides previously identified or not in ECN90 β cells are in blue and red, respectively, with peptide length variants marked with an asterisk. The 3 HLA-A2-restricted peptides retained for CD8⁺ T-cell studies are highlighted in yellow: a SCG5₁₈₆₋₁₉₅ shorter variant with a higher affinity than the previously identified SCG5₁₈₆₋₁₉₆ peptide, a newly identified CHGB₄₄₀₋₄₄₈ peptide and an IAPP₁₅₋₁₇/IAPP₅₋₁₀ or PTPRN₅₉₆₋₅₉₈/IAPP₅₋₁₀ peptide splice as sequence (VAL/KLQVFL) previously identified in ECN90 β cells, which was finally assigned an HLA-A2 restriction based on the HLA-I haplotype of the islet preparation (HLA-A*02:01/25:01, -B*39:01/51:01, -C*12:03/14:02, i.e. sharing only HLA-A2 with ECN90 β cells). NA, not available.

Study phase	Group	Case ID	Age (yrs)	Gender (M/F)	T1D duration (wks)	GADA	IA-2A	ZnT8A	Therapy
		H004N	44	М	NA	NA	NA	NA	NA
		H005N	36	F	NA	NA	NA	NA	NA
		H017N	35	F	NA	NA	NA	NA	NA
		H079O	36	F	NA	NA	NA	NA	NA
		H087N	34	М	NA	NA	NA	NA	NA
		H106S	47	F	NA	NA	NA	NA	NA
Sereening	Healthy	H297S	28	М	NA	NA	NA	NA	NA
Screening	(n=14)	H312C	36	М	NA	NA	NA	NA	NA
		H314C	26	М	NA	NA	NA	NA	NA
		H315C	43	F	NA	NA	NA	NA	NA
		H316C	27	F	NA	NA	NA	NA	NA
		H354C	26	F	NA	NA	NA	NA	NA
		H356C	28	М	NA	NA	NA	NA	NA
		H372C	24	F	NA	NA	NA	NA	NA
			34.5	43% M					
			(24-47)	57% F					
		D216P	27	М	10.9	+	-	-	Insulin
		D217Db	34	М	19.0	+	NA	NA	Insulin
		L264D	28	F	32.0	+	+	+	Insulin
		D267T	25	F	1.0	+	NA	NA	Insulin
	T1D (n=10)	D287T	30	М	4.1	+	NA	NA	Insulin
		D314D	44	F	0.6	+	+	+	Insulin
		D324D	28	F	11.3	+	+	NA	Insulin
		D325D	19	М	0.3	-	-	+	Insulin
		D327V	34	F	0.7	+	-	NA	Insulin
		D339V	29	М	0.4	+	+	NA	Insulin
Validation			28.5	50% M	2.6				
Valluation			(19-44)	50% F	(0.3-32.0)				
		H015T	30	М	NA	NA	NA	NA	NA
		H170S	34	F	NA	NA	NA	NA	NA
		H172O	33	F	NA	NA	NA	NA	NA
		H192C	25	F	NA	NA	NA	NA	NA
	Healthy	H193C	45	F	NA	NA	NA	NA	NA
	(n=10)	H227C	26	М	NA	NA	NA	NA	NA
		H230C	29	F	NA	NA	NA	NA	NA
		H245C	22	М	NA	NA	NA	NA	NA
		H314C	26	М	NA	NA	NA	NA	NA
		H354C	26	F	NA	NA	NA	NA	NA
			27.5	40% M					
			(22-45)	60% F					

Table S4. Related to Fig. 4-5. Characteristics of HLA-A2⁺ study subjects for ex-vivo MMr studies on PBMCs. The healthy donors analyzed in Fig. 4 (screening phase) and the T1D and age/sex-matched healthy donors analyzed in Fig. 5 (validation phase) are listed. The distribution of age, gender and T1D duration is shown at the bottom of each list (median and range for numerical variables). GADA, IA-2A, ZnT8A, anti-GAD, -IA-2 and -ZnT8 aAbs, respectively; NA, not available or not applicable.

	nPOD	0	Age	T1D	Positive	C-peptide			Pancr	eas MMr	+		PLN MMr ⁺			
	case	Sex	(yrs)	(yrs)	aAbs	(ng/ml)	IAPP	ISL1	UCN3	ZnT8	MelanA	IAPP	ISL1	UCN3	ZnT8	MelanA
	6070	F	23	7	IA-2/mIAA	<0.05			39	74	0			32	0	
	6161	F	19	7	IA-2/mIAA	<0.05	110	59		124	0	0	0		1176	0
	6211	F	24	4	GAD/IA-2/ZnT8/mIAA	<0.05	29	32		30	0	103	45		60	54
(6=	6212	М	20	5	mIAA	<0.05	19	29		0		0	0		0	
D (n=	6237	F	18	12	GAD/mIAA	<0.05			8	267	0			0	0	
1	6242	М	39	19	IA-2/mIAA	<0.05	33	9		66	0	0	0		101	0
	6243	М	13	5	mIAA	0.42	47	17		0		72	34		209	0
	6258	F	39	37	mIAA	<0.05			19	118	0			0	299	0
	6325	F	20	6	GAD/IA-2	0.14			28	28	0			35	0	0
	6080	F	69	NA	GAD/mIAA	1.84	38	39		55	25	43	41		50	61
	6101	М	65	NA	GAD	26.18	36	30		0		0	0		0	
	6123	F	23	NA	GAD	2.01	22	48		0		0	0		0	
6=	6151	М	30	NA	GAD	5.49			9	28	0					
ü ±	6154	F	49	NA	GAD	<0.05			21	64	0					
aAb	6171	F	4	NA	GAD	8.95	13	9		37	0	0	0		0	
	6347	М	9	NA	mIAA	3.26	15	7		33	0	66	60		60	191
	6388	F	25	NA	GAD/mIAA	1.38			0	34	0			0	35	0
	6397	F	21	NA	GAD	12.8			6	42	0			0	189	0
	6103	М	2	NA	—	0.98			0	55	0			0	0	
	6174	М	21	NA	—	3.00	0	0				28	0			
	6179	F	20	NA	_	2.74			0	96	0			67	150	0
	6182	М	3	NA	—	2.28			4	23	0			89	163	65
=12)	6227	F	17	NA	—	2.75			18	3	0			0	0	
es (r	6234	F	20	NA	—	6.89			0	6	0			0	0	
iabet	6254	М	38	NA	—	6.43	0	0		0		0			0	
No d	6271	М	17	NA	—	11.47	0	5		0		0	0		0	0
	6287	F	57	NA	—	4.75	7	8		4	0	0	0		0	
	6289	М	19	NA	—	8.05	0	6		0		55	0		0	
	6357	М	5	NA	_	8.82	9	29		0		32	0		0	
	6366	F	21	NA	—	0.41	14	24		0		55	0		0	
	6028	М	33	17		22.40		25		0			0		0	
(n=4)	6059	F	19	0.3	_	10.68	12			0		0			107	0
T2D (6273	F	45	2	_	3.17	13	7				0	0			
	6275	М	48	2	—	3.46	0			0		133	117		0	

Table S5. Related to Fig. 6. nPOD cases analyzed by *in-situ* **tissue MMr staining.** The clinical characteristics of each case are reported along with the counts $(x10^{-3})$ of MMr⁺ cells/mm² pancreas and PLN section area for each of the indicated peptides. Positive sections are marked in red. Case #6287 (presenting a circumscribed neuroendocrine tumor in the pancreatic pan-body region; pan-tail region analyzed here) was classified as a non-diabetic control. NA, not applicable; mIAA, micro-insulin aAbs.

Lasers and filters		Photo	omultiplier t	tubes (PM	Ts)	
488 nm blue laser	С	В	А			
Long pass (LP) filter	١	502	655			
Band pass (BP) filter	488/10	530/30	695/40			
Fluorochrome	SSC	FITC				
561 nm yellow-green laser	Е	D	С	В	А	
Long pass (LP) filter	١	600	635	685	735	
Band pass (BP) filter	582/15	610/20	660/20	710/50	780/60	
Fluorochrome	PE	PE-CF594			PE-Cy7	
633 nm red laser	С	В	А			
Long pass (LP) filter	١	710	750			
Band pass (BP) filter	660/20	730/45	780/60			
Fluorochrome	APC		APC-H7			
405 nm violet laser	F	E	D	С	В	А
Long pass (LP) filter	١	502	595	630	675	750
Band pass (BP) filter	450/50	525/50	605/20	655/8	710/50	780/60
Fluorochrome	BV421	L/D Aqua		BV650	BV711	BV786

Table S6. Related to STAR Methods. Configuration of the flow cytometer used forHLA-A2 MMr assays. L/D, Live/Dead.