

A retrospective study of factors associated with treatment decision for nontuberculous mycobacterial lung disease in adults without altered systemic immunity

Judith Provoost, Florent Valour, Delphine Gamondes, Sandrine Roux, Nathalie Freymond, Emilie Perrot, Pierre-Jean Souquet, Lize Kiakouama-Maleka, Christian Chidiac, Gérard Lina, et al.

▶ To cite this version:

Judith Provoost, Florent Valour, Delphine Gamondes, Sandrine Roux, Nathalie Freymond, et al.. A retrospective study of factors associated with treatment decision for nontuberculous mycobacterial lung disease in adults without altered systemic immunity. BMC Infectious Diseases, 2018, 18 (1), pp.659. 10.1186/s12879-018-3559-x. inserm-02043709

HAL Id: inserm-02043709 https://inserm.hal.science/inserm-02043709

Submitted on 21 Feb 2019 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Open Access



A retrospective study of factors associated with treatment decision for nontuberculous mycobacterial lung disease in adults without altered systemic immunity

Judith Provoost¹, Florent Valour^{2,3,4}, Delphine Gamondes⁵, Sandrine Roux², Nathalie Freymond¹, Emilie Perrot¹, Pierre-Jean Souquet^{1,4}, Lize Kiakouama-Maleka¹, Christian Chidiac^{2,4}, Gérard Lina^{3,4,6}, Oana Dumitrescu^{3,4,6}, Agathe Sénéchal¹ and Florence Ader^{2,3,4*}

Abstract

Background: Nontuberculous mycobacteria (NTM) lung diseases are increasingly recognized as chronic opportunistic infections, occurring in individuals with a wide variety of underlying conditions. In the absence of systemic immunodeficiency, decision of NTM lung disease treatment must relies on a careful risk/benefit assessment, given the requirement of long-term administration of multidrug therapies supported by limited evidence. The primary objective was to identify the factors associated with anti-NTM treatment initiation. Clinical and radiological outcome upon treatment were studied.

Methods: This retrospective, single center study (2013–2016, 45 months) addressed the criteria supporting treatment decision among adults with NTM lung disease without systemic immunodeficiency at our institution, with the assigned goal to harmonize the practice. All patients matched the current international definitions of NTM lung disease according to the American Thoracic Society criteria. Factors associated with anti-NTM treatment were investigated by conditional logistic regression. Clinical and radiological outcomes of treated and untreated NTM-disease cases were examined. Mortality rate was assessed. An expert radiologist conducted a blinded computed tomography (CT)-scan review of the treated and untreated patients.

(Continued on next page)

* Correspondence: florence.ader@chu-lyon.fr

²Département des Maladies infectieuses et tropicales, Hôpital de la Croix-Rousse, Hospices Civils de Lyon, 103, Grande-Rue de la Croix-Rousse,

³CIRI-Centre International de Recherche en Infectiologie, Inserm, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, Ecole Normale Supérieure de Lyon, Univ Lyon, F-69007 Lyon, France Full list of author information is available at the end of the article



© The Author(s). 2018 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

^{69317,} cedex 04 Lyon, France

(Continued from previous page)

Results: Among 51 cases of NTM lung diseases, 25 (49%) received anti-NTM treatment. In univariate analysis, a body mass index (BMI) < 18 kg/m² (odds ratio (OR), 4.2 [95% confidence interval (Cl) 1.2–15.2]; p = 0.042), hemoptysis (OR, 11.8 [95% Cl 1.35–12.9]; p = 0.026), excavation(s) (OR, 4.8 [95% Cl 1.4–16.4], p = 0.012), prior anti-NTM treatment (OR, 5.65 [95% Cl 1.06–29.9]; p = 0.042), *Aspergillus* spp. co-infection (OR, 6.3 [95% Cl 1.8–22.2]; p = 0.004) were associated with treatment initiation. In multivariate analysis, *Aspergillus* spp. co-infection was the only independent determinant of treatment initiation (OR, 5.3 [95% Cl 1.1–25.4]; p = 0.036). Twenty-one (81%) patients received ≥3 anti-NTM drugs. Median treatment duration and follow-up were 36.3 (interquartile range [IQR], 13.1–64.4) weeks and 17.1 (IQR, 8.7–27.1) months, respectively. Regarding radiological outcome, 85 CT-scans were reviewed, showing similar rates of regression or stabilization in treated and untreated patients. Overall mortality rate was not different in treated and untreated patients.

Conclusion: The most relevant variable associated with anti-NTM treatment initiation was *Aspergillus* spp. co-infection. Radiological regression or stabilization of pulmonary lesions was not different between the treated and untreated patients.

Keywords: Aspergillus spp., Bronchiectasis, Chronic obstructive pulmonary disease, Hemoptysis, Mycobacteria, *Mycobacterium avium complex*, Nontuberculous mycobacteria

Background

Nontuberculous mycobacteria (NTM) are ubiquitous bacteria of environmental origin including a widely diverse number of species (> 160), some of which cause disease in humans [1–3]. Prevalence of NTM lung diseases is unexpectedly increasing in industrialized countries, as consistently uncovered by recent studies [3, 4]. The key issue remains to determine whether NTM are the true and single promoter of an evolving lung disease or chronic airway colonizers, among others. To standardize the diagnosis of NTM lung disease, the guidelines for NTM diagnosis of the American Thoracic Society (ATS)/Infectious Disease Society of America (IDSA) and the British Thoracic Society (BTS) require isolation and growth of the same NTM strain on at least two separate samples from the patient [5, 6].

Human host and pathogenic NTM relationship is still poorly understood, as NTM virulence is highly variable from one species to another. NTM lung disease is strongly associated with pre-existing pulmonary conditions such as chronic obstructive pulmonary disease (COPD), cystic fibrosis, idiopathic bronchiectasis, prior active tuberculosis or pneumoconiosis [6]. It is also frequently associated with genetic or acquired systemic immune deficiency such as defects in the pathways of inflammatory cytokines interleukin (IL)-12, tumor necrosis factor (TNF)- α or interferon (IFN)- γ , immunosuppressive treatments (including anti-TNF- α therapy or corticosteroids), solid-organ transplantation, or acquired immune deficiency syndrome (AIDS)/human immunodeficiency virus (HIV) infection [1-7]. However, it may also occur in individuals without recognized severe immune local or systemic deficiency. In the absence of patent predisposition, NTM diseases are overrepresented among the specific morphotype of slender women with a low body fat [8].

Treatment decision for NTM lung disease is challenging. There is debate as to which patients should benefit the most from treatment according to medical background, comorbidities, clinical status, radiologic features and causal NTM strain. Assessment of clinical, microbiologic, and radiologic response to treatment is not standardized as well. We focused on NTM lung diseases in adults without systemic immunodeficiency that met the ATS criteria guidelines. Based on the comparison of a group of treated and untreated patients, the primary objective was to identify the factors associated with physician decision of initiating anti-NTM treatment. Secondary objectives were to study the outcome upon treatment and to propose a standardized evaluation for the diagnosis and decision making to treatment of NTM lung diseases in adults without systemic immunodeficiency.

Methods

Study design and patient population

We conducted a retrospective, observational, single-center study between January 2013 and February 2016 (45 months) among adults (\geq 18 year-old) without systemic immunosuppression presenting NTM lung disease. Exclusion criteria were HIV infection, cystic fibrosis, primary ciliary dyskinesia, active malignant disease, solid-organ transplantation or ongoing immunosuppressant treatments such as TNF inhibitor or high-dose corticosteroid (\geq 1 mg/kg more than 21 days). Case identification was based on cross-referencing the databases of the mycobacteria laboratory and the departments of infectious and pulmonary diseases. Patients eligible for inclusion in the NTM lung disease cohort were those who matched the

criteria previously defined by the ATS/IDSA and the BTS guidelines with the minimum requirement of clinical and microbiologic following criteria: (i) pulmonary symptoms associated with multifocal bronchiectasis with multiple small nodules on computed tomography (CT)-scan; (ii) proper exclusion of other diagnoses; (iii) NTM-positive culture results from at least two separate expectorated sputum samples or a NTM-positive culture result from at least one bronchial wash or lavage [5, 6]. Patients' characteristics at diagnosis were collected in order to perform analysis on 146 selected variables: demographics; history of predisposing factors; underlying pulmonary diseases; comorbidities; pulmonary function testing; respiratory bacterial or mycological co-infection(s), which definition was similar to NTM criteria, namely positive culture isolation of the same species from at least two separate expectorated sputum samples or a positive culture result from at least one bronchial wash or lavage; immunologic status; nutritional status; clinical features; microbiologic assessment through identification of NTM species on positive NTM cultures and sample culture conversions; radiologic features on high-resolution CT-scans (fibrocavitary disease or nodular/bronchiectasis disease); prior treatment for NTM lung disease, treatment combination and duration; outcome. Because of the retrospective observational nature of the study and the lack of any modification in patients' management, the need for informed consent was waived with the authorization of the Ethics Committee of Lyon University Hospital (Comité d'Ethique, Hospices Civils de Lyon), which approved the study under the number 17-207.

Radiologic assessment

An independent expert chest radiologist, blinded to the patient information, retrospectively reviewed the CT-scans performed without injection of intravenous contrast media, assigned in random order at diagnosis and six to 24 months after treatment or during the follow up of the untreated patients. The number and size of cavity(ies) and their wall thickness were evaluated in the lung window setting. Nodular opacity(ies) (\geq 10 mm), cluster(s) of small nodules (\leq 5 mm), the tree-in-bud pattern, the presence of bronchiectasis in any of the lobes or multifocal bronchiectasis were evaluated. Based on the number and size of the lesions, the expert classified the lesions as improved, stable or worsening.

Endpoints

The primary endpoint was to identify the factors significantly involved in the decision of initiating anti-NTM treatment by patient referent physician. Secondary endpoints were the assessment of clinical and radiological outcomes upon anti-NTM treatment in comparison with no treatment. Based on these findings, a standardized appraisal was proposed to assist diagnosis management and treatment decision for NTM lung diseases in adults without altered systemic immunity.

Statistical analysis

Descriptive data were used to estimate the frequencies of the study variables. There were expressed as count (percentage, %) for dichotomous variables and as medians (interquartile range [IQR]) for continuous values. The number of missing values was excluded from the denominator. Non-parametric statistical methods Fisher exact test, x2 test and Mann-Whitney U test were used to compare groups, where appropriate. The probability of treatment initiation over time was evaluated by Kaplan-Meier survival curve analysis, with group comparison using the log-rank (Mantel-Cox) test. Stepwise binary logistic regression analysis was used to assess the determinants for treatment initiation, expressed as odd ratios (OR) with 95% confidence intervals (95% CI). After checking the variables for interactions, variables with medical meaning and with *p*-values obtained in the univariate analysis of < 0.15 were included in the final multivariate model. A value of p < 0.05 was considered significant. All analyses were performed using SPSS software version 24.0 (SPSS. Chicago. IL).

Results

Out of 149 patients eligible to ATS/IDSA NTM lung disease criteria, 51 patients were included in the study, of who 25 (49%) received and 26 (51%) did not receive anti-NTM treatment (Fig. 1). The median age was 68 (interquartile range [IQR], 59-75) year-old with a male/ female ratio of 0.76. Etiologic NTM agents were Mycobacterium avium (n = 17, 33.3%), M. chimaerae (n = 14, 27.5%), M. xenopii (n = 9, 17.6%), M. intracellulare (n =9, 17.6%), M. simiae (n = 3, 5.9%), M. kansasii (n = 1, 2%), and *M. abscessus* (n = 1, 2%), with three patients having ≥ 2 concomitant NTM lung diseases. Importantly, the evidence of NTM lung disease has led to diagnose six underlying chronic lung diseases, of which a genetically documented cystic fibrosis in a 61 year-old women. On descriptive analysis, patient's characteristics did not significantly differ, to the exception of a lower BMI (p =0.023) and a higher number of previously known NTM lung disease (p = 0.038) in treated versus untreated patients (Table 1). Notable percentages of missing data at diagnosis have to be acknowledged for active tobacco smoking (64.7%, n = 33), respiratory functional testing (23.5%, n = 12) with a very few patients having a 6-min walk test, baseline arterial blood oxygenation levels (58.8%, n = 30), CT-scan follow up within 24 months after diagnosis in untreated patients (46%, n = 12).

Factors leading the patient referent physician to initiate anti-NTM treatment were assessed using bivariate analysis.



Table 1 Patients' demographic and baseline descriptive characteristics

	Treated patients ($n=25$)	Untreated patients ($n=26$)	P-value
Age, median (IQR) years	65 (54-74)	70 (60.8-76)	0.227
Gender (male)	9 (36)	13 (50)	0.4
Active smokers	6 (24)	6 (23)	1
Cumulative tobacco smoking (IQR) pack/year	32.5 (28.8-52.5)	40 (21.3-60)	0.79
BMI (IQR) kg/m ²	17.9 (16.8-19.9)	19.9 (18.3-23.8)	0.023
History of tuberculosis	6 (24)	5 (19.2)	0.74
History of NTM lung disease	8 (32)	2 (7.7)	0.038
History of croup during infancy	1	1	1
Gastroesophageal reflux disease	4 (16)	2 (7.7)	0.42
History of cured malignant solid tumor	1 (4)	5 (19.2)	0.19
Diabetes mellitus	0	2 (7.7)	0.49
Chronic heart failure	0	4 (16)	0.11
Asthma or atopic disease	6 (24)	2 (7.7)	0.14
Chronic obstructive pulmonary disease	4 (16)	9 (34.6)	0.19
Bronchiectasis	14 (56)	14 (53.8)	1
Chronic interstitial lung disease	2 (8)	5 (19.2)	0.42
Chronic rheumatic disease	6 (24)	2 (7.7)	0.14
Autoimmune disease	3 (12)	4 (15.4)	1
History of thoracic surgery	5 (20)	3 (11.5)	0.46
Corticosteroid inhaled therapy	6 (24)	7 (27)	1

NOTE Data are No. (%) of patients, unless otherwise indicated. Variables were compared using Mann–Whitney U test, $\chi 2$ test or the Fisher exact test, where appropriate

Abbreviations: IQR, interquartile range

Among the factors significantly associated with treatment were the BMI < 18 kg/m² (odds ratio (OR), 4.2 [95% confidence interval (CI) 1.2–15.2]; p = 0.042), prior anti-NTM treatment (OR, 5.65 [95% CI 1.06–29.9]; p = 0.042), hemoptysis (OR = 11.8 [95% CI 1.35– 102.9]; p = 0.026), the presence of excavation(s) on lung CT-scan (OR, 4.8 [95% CI 1.4–16.4], p = 0.012), *Aspergillus* spp. co-infection (OR, 6.3 [95% CI 1.8– 22.2]; p = 0.004) (Table 2). In multivariate analysis,

Aspergillus spp. co-infection was the single independent factor associated with anti-NTM treatment (OR, 5.3 [95% CI 1.1–25.4]; p = 0.036) (Table 3).

The probability of treatment initiation over time according to the presence or not of targeted variables was investigated. The probability was significantly higher in case of BMI < 18 vs. > 18 kg/m² (p = 0.006), of hemoptysis vs. no hemoptysis (p = 0.009), of *Aspergillus* spp. co-infection vs. no co-infection (p = 0.029), of pulmonary excavation(s) vs.

Table 2 Bivariate analysis of the factors associated with treatment initiation in adult patients with nontuberculous mycobacterial lung disease without systemic immunodeficiency

Tested variables	Treated patients (n=25)	Untreated patients (n=26)	P-value	OR [95% CI]
Clinical criteria				
$BMI < 18 (kg/m^2)$	13/24 (54.2)	5/23 (21.7)	0.026	4.25 [1.18-15.2]
Weight loss > 5%	17 (68)	10 (38.5)	0.05	3.4 [1.07-10.77]
fatigue	13 (52)	7 (27)	0.071	2.94 [0.9-9.46]
chronic cough	16 (64)	14 (54)	0.572	1.52 [0.49-4.69]
night sweats	5 (20)	3 (11.5)	0.465	1.92 [0.41-9.05]
hemoptysis	8 (32)	1 (3.8)	0.026	11.8 [1.3-102.8]
Microbiological criteria				
AFB-positive sputum sample on direct examination	13 (52)	8 (30.8)	0.160	2.44 [0.77-7.65]
NTM-positive culture NTM on lower respiratory tract sample	2 (1-2.3)	1 (1-1.8)	0.044	1.79 [1-3.2]
Functional criteria				
FEV1 (L)	1.6 (1.2-2.2)	1.9 (1.2-2.5)	0.554	0.72 [0.34-1.51]
Immunological and nutritional status				
total lymphocyte count (G/L)	1.3 (1.2-1.7)	1.4 (1.1-2.1)	0.388	0.95 [0.86-1.04]
CD4+ lymphocyte count (%)	54 (4.5-61.5)	40 (37.8-44.3)	0.031	1.07 [0.95-1.21]
serum albumin (mg/L)	39.5 (36-43)	39 (33-42.5)	0.481	1.05 [0.97-1.14]
serum pre-albumin (mg/L)	0.21 (0.15-0.24)	0.22 (0.15-0.25)	0.974	1.25 [0-1.48]
vitamin D deficiency	6/10 (60)	6/7 (86)	0.338	0.25 [0.02-2.95]
Co-infection(s)				
Mycobacterium tuberculosis	1 (4)	1 (4)	0.35	1.04 [0.06-17.6]
Pseudomonas spp.	2 (8)	3 (11.5)	1	0.68 [0.1-4.4]
Aspergillus spp.	15 (60)	5 (19.2)	0.004	6.3 [1.8-22.2]
Radiologic features at diagnosis				
nodule(s) (≥ 10mm)	12 (48)	7/23 (30)	0.25	2.11 [0.65-6.9]
cluster(s) of micronodules (< 5mm)	16 (64)	10/23 (44)	0.246	2.31 [0.72-7.4]
cavitation(s)	18 (72)	8/23 (35)	0.019	4.82 [1.42-16.4]
bronchiectasis	14 (56)	11/23 (48)	0.773	1.39 [0.45-4.3]
emphysema	9 (36)	11/23 (48)	0.559	0.61 [0.19-1.95]
Radiologic review				
regression	8/23 (35)	4/13 (31)	1	1.2 [0.28-5.15]
stabilization	9/23 (39)	6/13 (46)	0.736	0.75 [0.19-2.97]
worsening	6/23 (26)	3/13 (23)	1	1.18 [0.24-5.8]

NOTE Data are No. (%) of patients, unless otherwise indicated. Denominators have been stipulated for variables with missing data. Odds ratios (OR) were determined by conditional logistic regression with 95% confidence interval (95% CI)

Abbreviations: AFB, acid-fast bacilli; BMI, body mass index; CI, confidence interval; FEV1, forced expiratory volume in 1 second; NTM, nontuberculous mycobacteria; OR, odds ratio

Independent variables	OR [95% CI]	<i>P</i> -value
Aspergillus spp.	5.33 [1.1-25.5]	0.036
weight loss > 5%	2.44 [0.6-10.2]	0.22
hemoptysis	3.2 [0.28-36.3]	0.35
excavation(s)	3.54 [0.83-15.1]	0.087

Abbreviations: CI, confidence interval; OR, odds ratio

no excavation (p = 0.005) (Fig. 2a, b, c and d, respectively). In the treated group, 21 (84%) received ≥3 anti-NTM drugs, among whom 23 (92%) received a macrolide as part of a first-line regimen. Median anti-NTM treatment duration was 36.3 (IQR, 13.1–64.4) weeks. Thirteen (52%) of the treated patients developed at least one antimicrobial-related adverse event, among which gastrointestinal disorders (n = 6), impaired blood cell count (n = 5) and cholestatic and/or cytolytic hepatitis (n = 4). According to physician judgment, clinical stabilization and improvement was obtained for 8 (34.8%) and 12 (52.2%) of treated patients, respectively. Patient's median follow-up from diagnosis assessment was 17.1 (IQR, 8.7-27.1) months. Eighty-five CT-scans were reviewed for comparative assessment after treatment or over the course of follow-up in the untreated subset. The median interval between the initiation of NTM treatment and the date of the assessed images was 56.7 (IQR, 43.3-97.7) weeks. Although 34.8% (n = 8) and 39.1% (n = 9)experienced regression or stabilization of their pulmonary lesions in the treated patients, these rates did not differ from the untreated patients with 30.8% (n = 4) and 46.2% (n = 6) of regression and stabilization over the course of their follow-up, respectively (Table 2). Finally, all-cause mortality was not different between treated and untreated groups, although lost to follow-up was high in the untreated group (n = 10, 38.5%). Regarding the four patients that deceased (n = 3 in the treated group and n = 1 in the untreated group), the cause of mortality was linked to the underlying diseases rather than NTM-related mortality.

Discussion

In the present study, treated patients were characterized by BMI < 18 kg/m², presence of hemoptysis and excavation(s), *Aspergillus* spp. co-infection, and prior anti-NTM treatment. *Aspergillus* spp. co-infection was the only independent factor associated with treatment initiation. A single study from five English centers has recently addressed the factors that influence anti-NTM treatment initiation using similar retrospective design of treated and untreated cohort comparison (2010–2014), to the exception of different



Fig. 2 Overall probability of anti-nontuberculous mycobacteria treatment initiation according to the presence or the absence of the following variables: **a** body mass index < 18 kg/m^2 ; **b** hemoptysis; **c** *Aspergillus* spp. co-infection; **d** lung excavation(s). Overall survival probabilities were determined by the Kaplan-Meier method and were compared by the log-rank test

Table 4 Standardized appraisal for patients with nontuberculous
mycobacterial lung diseases without systemic immunodeficiency
referred at our institution

Table 4 Standardized appraisal for patients with nontuberculous
 mycobacterial lung diseases without systemic immunodeficiency referred at our institution (Continued)

Comments			Comments		
Exhaustive history record infant respiratory	croup, Influenza, measle	lung HRCT scan	Micronodules (≤ 5mm), nodules (≥ 10 mm), pulmonary consolidations,		
diseases		Cytology and microbiologic	tests on respiratory samples		
gastro-oesophageal reflux disease		BALF cytological composit	ion		
atopic dermatitis		sputum/BALF direct examination			
eating disorders	anorexia, bulimia	sputum/BALF	Non-fermenting gram-negative bacilli		
family history of respiratory disease		collection for bacterial culture	(Pseudomonas aeruginosa)		
prior tuberculosis		sputum/BALF collection for	filamentous fungi		
prior anti-NTM treatment		mycological culture	human Pospiratony syncytial virus		
Environmental exposure		collection for	parainfluenza virus, metapneumovirus,		
passive tobacco		respiratory virus PCR screening	rhinovirus, and coronavirus		
active tobacco	Packs-per-vear	Specific tests depending on the setting			
smocking		α -1 antitrypsin level	lung emphysema		
other toxic/drug		sweat test	cystic fibrosis		
exposure		CFTR gene screening			
chronic alcoholism	micro-inhalations	airway cilia sample	primary ciliary dyskinesia		
habitat characteristics Blood tests	domestic water system, allergens	IFN-γ and IL-12 plasma levels	host predisposition to NTM diseases		
serum albumin	nutritional status	blood anti-IFN-y			
serum pre-albumin		autoantibodies			
vitamin D (25-OH D3)		IFN-γ or IL-12 gene			
HIV serology	cellular immune defect		oalveolar lavage fluid: CETP, cystic fibrosis		
T CD4+ and CD8+ lymphocyte counts		ADDreviations: BALF, pronchoaiveolar lavage fluid; CFTR, cystic fibrosis transmembrane conductance regulator, FEV1, forced expiratory volume in 1 second; HRCT, high-resolution computed tomography; IFN-γ, interferon camma: In immunoclobulins: II. interdeukin			
total IgG serum level	hypogammaglobulinemia	garrina, ig, inimanogiobalin	, i.e. interreducti		
total eosinophil count	Aspergillus-related allergic lung diseases	inclusion critoria y	which allowed non AIDS/HIV im		
total IgE serum level		munosuppressed patients (36 and 33% in the treated and untreated subsets, respectively) to be evaluated.			
Aspergillus-specific IgE					
Aspergillus serology		In the multivariate	e analysis, patients had increased		
galactomannan antigenemia	invasive aspergillosis	odds of anti-NTM treatment in case of cavitation on CT imaging, night sweats and weight loss [9]. Here, three out of four criteria are part of guideline cri- teria leading to decision of anti-NTM treatment. They all			
Interferon-γ reactive assay	latent tuberculosis infection				
Lung function testing		reflect a degree of sev	rerity of NTM lung disease linked with		
spirometry including FEV1	lung volumes	progression of an impaired respiratory condition. The current problematic of NTM lung diseases shifts from dis-			
plethysmography		tinguishing colonizati	on from infection toward differentiat-		
pulse oxymetry	oxygen level	ing stable, poorly active vs. progressive active NTM disease,			

arterial blood gas tests

6-minutes walk test

stress test

Imagery

chest X-ray

11 h e 2, the latter being responsible for further structural lung damage(s). The risk/benefit analysis includes prescribing recommended long-term multidrug regimens with concerns over suboptimal cure rates and frequently reported drug-related side effects. To support these arguments, others have already shown that physician ATS/IDSA guideline adherence is moderate confirming such a trend of intuitive clinical behavior mostly based on evidence of severity and/ or patent progression of the disease. A multicenter study has reported a sample of 349 US physicians treating 915 patients with M. avium complex (MAC) lung disease. Overall, 55% (n = 411) of patients with MAC lung disease received some type of anti-NTM treatment from their physician and 13% prescribed antibiotic regimens were consistent with ATS/IDSA guidelines [10]. Another study conducted in France has shown that among a cohort of 31 NTM lung diseases, only 12 (38.7%) received appropriate treatment matching ATS/IDSA guidelines [11]. Inappropriate prescriptions were mostly related to shorter treatment duration (6 months or less) and/or off-recommendation regimen, particularly those excluding macrolide from the combination or those using a single-drug macrolide regimen [10, 11]. It has to be acknowledged that guidelines specify that treatment for MAC-associated lung disease in HIV-negative individuals can be a three-times-weekly drug regimen upon culture conversion while on therapy for 1 year, which may favor treatment compliance [5, 12, 13].

Patients with bronchiectasis and NTM lung disease have a higher prevalence of being sensitized to *Aspergillus* than patients with NTM-free bronchiectasis [12]. Allergic airway manifestations in response to Aspergillus are termed Aspergillus-related lung diseases with a spectrum going from Aspergillus-induced hypersensitivity to the severe allergic bronchopulmonary aspergillosis (ABPA) [14]. By itself, the coexistence of NTM and Aspergillus in lung airway justifies the need for testing Aspergillus serology, total serum immunoglobulin (Ig)E and Aspergillus-specific IgE levels as well as mycological direct examination and culture of sputum or bronchial aspirates for presence of filamentous fungi in the diagnosis algorithm of patients with NTM lung diseases. Active co-infections with NTM and Aspergillus spp. have also been described, in which patients with NTM lung disease develop chronic forms of pulmonary aspergillosis which definitions and management have been revisited in recent updated guidelines [14–17].

Radiographic improvement may be hampered by concomitant lung disease and the limited potential for resolution of consolidated radiologic abnormalities. A previous study has investigated radiologic response to treatment showing consistent results with those found in the present study. Although anti-NTM treatment led to an improvement or stabilization of lesions for a majority of patients, these modifications were not significantly different from the untreated group who went through CT-scan follow up indicating that anti-NTM treatment did not lead to radiological abnormalities reversion [9]. The present study has strengths and limitations. The strength is the study of the largest cohort so far of NTM lung diseases in patients without systemic immunodeficiency with exhaustive data collection and blinded radiological assessment. We acknowledge the biases that contribute to mitigate conclusions from the study such as being conducted in a single center, the important differences in physician's management resulting in lack of consistency in treatment decision making, the number of missing data. In addition, the CT-scans were not performed at fixed intervals, particularly in the untreated subset of patients. Finally, treatment duration and outcome criteria were not standardized, which prevent to properly assess treatment efficacy. Useful consensus definitions for key outcome parameters to be used in the treatment of NTM lung diseases have been released very recently, which should harmonize data collection regarding NTM treatment [18].

Future researches are necessary to better define criteria associated with progressive active NTM disease in the immunocompetent setting. Concretely, this preliminary study has led to implement in our institution a standardized appraisal for the diagnosis of NTM lung diseases in this particular setting (Table 4). The aim is to provide a future basis for the development of a diagnosis scoring system supporting anti-NTM treatment decision. Future studies should focus as well on the most relevant CT imaging variables associated with response to treatment over time that may be applied in future clinical trials to assess treatment outcome.

Conclusions

In summary, the main factors supporting anti-NTM treatment decision in immunocompetent were low BMI, hemoptysis, lung excavation(s), prior anti-NTM treatment and *Aspergillus* pp. co-infection, the latter being the only independent factor. Anti-NTM treatment did not achieve radiological abnormalities reversion, as pulmonary lesions assessment showed no difference between the treated and the untreated patients. A diagnosis of NTM lung disease in an immunocompetent patient requires investigating the presence of a chronic pulmonary underlying disease.

Abbreviations

AIDS: Acquired immune deficiency syndrome; ATS: American thoracic society; BTS: British thoracic society; COPD: Chronic obstructive pulmonary disease; CT: Computed tomography; HIV: Human immunodeficiency virus; IDSA: Infectious disease society of America; IFN: Interferon; IL: Interleukin; NTM: Non tuberculous mycobacteria; TNF: Tumor necrosis factor

Acknowledgements

The authors gratefully acknowledge Pr Gilles Devouassoux for helpful insights.

Funding

No external funding was received for this study.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

JP contributed to conception and design of the study, acquisition of the data, interpretation of the data, drafted the manuscript and approved the final version; FV carried out the statistical analysis, participated in revision of the paper for important intellectual content, and approved the final version; DG carried out the blinded review of 85 CT-scans with the examination of each imagery, she participated in revision of the paper for important intellectual content, and approved the final version; GL and OD are in charge of the Mycobacteria laboratory and have provided NTM strain identification and drug susceptibility tests. They participated in revision of the paper for important intellectual content, and approved the final version; SR, NF, EP, PJS, LKM, and CC contributed to acquisition of the data, revision of the paper for important intellectual content, and approved the final version; FA and AS are the project initiators, contributed to conception and design of the study, drafted the manuscript and approved the final version. FA coordinated the project until the submission of the article. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The Ethics Committee of Lyon University Hospital (Comité d'Ethique, Hospices Civils de Lyon) approved the study under the number 17–207. Because of the retrospective observational nature of the study and the lack of any modification in patients' management, the need for informed consent was waived with the authorization of the Ethics Committee of Lyon University Hospital (Comité d'Ethique, Hospices Civils de Lyon), which approved the study under the number 17–207.

Consent for publication

Not applicable.

Competing interests

The authors have no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. No writing assistance was utilized in the production of this manuscript.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Département de Pneumologie, Hospices Civils de Lyon, Lyon, France. ²Département des Maladies infectieuses et tropicales, Hòpital de la Croix-Rousse, Hospices Civils de Lyon, 103, Grande-Rue de la Croix-Rousse, 69317, cedex 04 Lyon, France. ³CIRI-Centre International de Recherche en Infectiologie, Inserm, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, Ecole Normale Supérieure de Lyon, Univ Lyon, F-69007 Lyon, France. ⁴Université Claude Bernard Lyon 1, Lyon, France. ⁵Département de Radiologie, Hospices Civils de Lyon, Lyon, France. ⁶Institut des Agents Infectieux, Hospices Civils de Lyon, Lyon, France.

Received: 1 July 2018 Accepted: 26 November 2018 Published online: 14 December 2018

References

- Wassilew N, Hoffmann H, Andrejak C, Lange C. Pulmonary disease caused by non-tuberculous mycobacteria. Respiration. 2016;91:386–402.
- Aksamit TR, Philley JV, Griffith DE. Nontuberculous mycobacterial lung disease: the top ten essentials. Respir Med. 2014;108:417–25.
- Raju RM, Raju SM, Zhao Y, Rubin EJ. Leveraging advances in tuberculosis diagnosis and treatment to address nontuberculous mycobacterial disease. Emerg Infect Dis. 2016;22:365–9.
- Winthrop KL, McNelley E, Kendall B, et al. Pulmonary nontuberculous mycobacterial disease prevalence and clinical features: an emerging public health disease. Am J Respir Crit Care Med. 2010;182:977–82.
- Griffith DE, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. Am J Respir Crit Care Med. 2007;175:367–416.
- Haworth CS, Banks J, Capstick T, et al. British Thoracic Society guidelines for the management of non-tuberculous mycobacterial pulmonary disease (NTM-PD). Thorax. 2017;72:ii1–ii64.

- Larsson LO, Polverino E, Hoefsloot W, Codecasa LR, Diel R, Jenkins SG, Loebinger MR. Pulmonary disease by nontuberculous mycobacteria-clinical management, unmet needs and future perspectives. Exp Rev Respir Med. 2017;11:977–89.
- Kartalija M, Ovrutsky AR, Bryan CL, et al. Patients with nontuberculous mycobacterial lung disease exhibit unique body and immune phenotypes. Am J Respir Crit Care Med. 2013;187:197–205.
- Rawson TM, Abbara A, Kranzer K, et al. Factors which influence treatment initiation for pulmonary non-tuberculous mycobacterium infection in HIV negative patients; a multicentre observational study. Respir Med. 2016;120:101–8.
- Adjemian J, Prevots DR, Gallagher J, Heap K, Gupta R, Griffith D. Lack of adherence to evidence-based treatment guidelines for nontuberculous mycobacterial lung disease. Ann Am Thorac Soc. 2014;11:9–16.
- El-Zeenni A, Chanoine S, Recule S, Stahl JP, Maurin M, Camara B. Are guidelines on the management of non-tuberculous mycobacteria lung infections respected and what are the consequences for patients? A French retrospective study from 2007 to 2014. Eur J Clin Microbiol Infect Dis. 2018;37:233–40.
- Griffith DE, Brown BA, Cegielski P, Murphy DT, Wallace RJ Jr. Early results (at 6 months) with intermittent clarithromycin including regimens for lung disease due to *Mycobacterium avium* complex. Clin Infect Dis. 2000;30:288–92.
- Jeong BH, Jeon K, Park HY, et al. Intermittent antibiotic therapy for nodular bronchiectatic Mycobacterium avium complex lung disease. Am J Respir Crit Care Med. 2015;191:96–103.
- Kunst J, Wickremasinghe M, Wells A, Wilson R. Nontuberculous mycobacterial disease and *Aspergillus*-related lung disease in bronchiectasis. Eur Respir J. 2006;28:352–7.
- Hafeez I, Muers MF, Murphy SA, Evans EG, Barton RC, McWhinney P. Nontuberculous mycobacterial lung infection complicated by chronic necrotising pulmonary aspergillosis. Thorax. 2000;55:717–9.
- Kobashi Y, Fukuda M, Yoshida K, Miyashita N, Niki Y, Oka M. Chronic necrotizing pulmonary aspergillosis as a complication of pulmonary *Mycobacterium avium* complex disease. Respirology. 2006;11:809–13.
- Denning DW, Cadranel J, Beigelman-Aubry C, Ader F, Chakrabarti A, Blot S, Ullman A, Dimopoulos G, Lange C. Chronic pulmonary aspergillosisrationale and clinical guidelines for diagnosis and management. Eur Resp J. 2016;47(1):45–68.
- van Ingen J, Aksamit T, Andrejak C, et al. Treatment outcome definitions in nontuberculous mycobacterial pulmonary disease: an NTM-NET consensus statement. Eur Respir J. 2018;51:1800170.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- · thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

