



HAL
open science

Targeting Oxidative Stress With Auranofin or Prima-1Met to Circumvent p53 or Bax/Bak Deficiency in Myeloma Cells

Benoit Tessoulin, Géraldine Descamps, Christelle Dousset, Martine Amiot, Catherine Pellat-Deceunynck

► To cite this version:

Benoit Tessoulin, Géraldine Descamps, Christelle Dousset, Martine Amiot, Catherine Pellat-Deceunynck. Targeting Oxidative Stress With Auranofin or Prima-1Met to Circumvent p53 or Bax/Bak Deficiency in Myeloma Cells. *Frontiers in Oncology*, 2019, 10.3389/fonc.2019. inerm-02055153

HAL Id: inerm-02055153

<https://inerm.hal.science/inerm-02055153>

Submitted on 3 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author's Proof









Carefully read the entire proof and mark all corrections in the appropriate place, using the Adobe Reader commenting tools ([Adobe Help](#)). Do not forget to reply to the queries.







We do not accept corrections in the form of edited manuscripts.

In order to ensure the timely publication of your article, please submit the corrections within 48 hours.

If you have any questions, please contact health.production.office@frontiersin.org.

Author Queries Form

Q1	The citation and surnames of all of the authors have been highlighted. Please check all of the names carefully and indicate if any are incorrect. Please note that this may affect the indexing of your article in repositories such as PubMed.	
Q2	Confirm that the email address in your correspondence section is accurate.	
Q3	Please ask the following authors to register with Frontiers (at https://www.frontiersin.org/Registration/Register.aspx) if they would like their names on the article abstract page and PDF to be linked to a Frontiers profile. Please ensure to provide us with the profile link(s) when submitting the proof corrections. Non-registered authors will have the default profile image displayed. "Benoit Tessoulin" "Christelle Dousset."	
Q4	If you decide to use previously published, copyrighted figures in your article, please keep in mind that it is your responsibility, as the author, to obtain the appropriate permissions and licenses and to follow any citation instructions requested by third-party rights holders. If obtaining the reproduction rights involves the payment of a fee, these charges are to be paid by the authors.	
Q5	Ensure that all the figures, tables and captions are correct.	
Q6	Verify that all the equations and special characters are displayed correctly.	
Q7	Ensure to add all grant numbers and funding information, as after publication this is no longer possible.	
Q8	Ensure, if it applies to your study, the ethics statement is included in the article.	

Q9	<p>Please ensure that any supplementary material is correctly published at this link: https://www.frontiersin.org/articles/10.3389/fonc.2019.00128/full#supplementary-material (you may need to copy-paste the link directly in your browser).</p> <p>Please provide new files if you have any corrections. Note that ALL supplementary files will be deposited to FigShare and receive a DOI. Notify us of any previously deposited material.</p>	
Q10	<p>Could you please confirm if all author affiliations are fine as listed?</p>	
Q11	<p>Please reduce short running title to maximum of five words.</p>	
Q12	<p>Please confirm whether the keywords are fine. Please keep the keywords list to a maximum of eight.</p>	
Q13	<p>Please provide the volume number, page range and doi for "(3)."</p>	
Q14	<p>Please update and add "Benaniba et al., submitted" to the reference list.</p>	



Targeting Oxidative Stress With Auranofin or Prima-1^{Met} to Circumvent p53 or Bax/Bak Deficiency in Myeloma Cells

Benoit Tessoulin^{1,2,3,4}, Geraldine Descamps^{1,2,3}, Christelle Dousset^{1,2,3}, Martine Amiot^{1,2,3} and Catherine Pellat-Deceunynck^{1,2,3*}

¹ CRCINA, INSERM, CNRS, Université d'Angers, Université de Nantes, Nantes, France, ² L'Héma-NexT, i-Site NexT, Université de Nantes, Nantes, France, ³ SIRIC ILIAD, Angers, Nantes, France, ⁴ Service d'Hématologie Clinique, Unité d'Investigation Clinique, CHU de Nantes, Nantes, France

OPEN ACCESS

Edited by:

Massimo Libra,
Università degli Studi di Catania, Italy

Reviewed by:

Himanshi Bhatia,
Washington University in St. Louis,
United States
Nadim Mahmud,
University of Illinois at Chicago,
United States

*Correspondence:

Catherine Pellat-Deceunynck
catherine.pellat-deceunynck
@univ-nantes.fr

Specialty section:

This article was submitted to
Hematologic Malignancies,
a section of the journal
Frontiers in Oncology

Received: 09 November 2018

Accepted: 13 February 2019

Published: xx February 2019

Citation:

Tessoulin B, Descamps G, Dousset C, Amiot M and Pellat-Deceunynck C (2019) Targeting Oxidative Stress With Auranofin or Prima-1^{Met} to Circumvent p53 or Bax/Bak Deficiency in Myeloma Cells. *Front. Oncol.* 9:128. doi: 10.3389/fonc.2019.00128

Prima-1^{Met} (APR-246) was previously shown to be dependent on glutathione inhibition and on ROS induction in cancer cells with mutated or deleted *TP53*. Because this ROS induction was, at least in part, due to a direct interference with the thioredoxin reductase enzyme, we investigated whether activity of Prima-1^{Met} could be mimicked by auranofin, an inhibitor of the thioredoxin reductase. We thus compared the activity of auranofin and Prima-1^{Met} in 18 myeloma cell lines and in 10 samples from patients with multiple myeloma or plasma cell leukemia. We showed that, similar to Prima-1^{Met}, the activity of auranofin was not dependent on either *TP53* status or p53 expression; was inhibited by N-acetyl-L-cysteine, a ROS scavenger; displayed a dramatic synergy with L-buthionine sulfoximine, an irreversible inhibitor of glutathione synthesis; and induced cell death that was not dependent on Bax/Bak expression. These data showed that auranofin and Prima-1^{Met} similarly overcome cell death resistance in myeloma cells due to either p53 deficiency or to mitochondrial dysfunction.

Keywords: Prima-1^{Met}, APR-246, auranofin, ROS, venetoclax

INTRODUCTION

Mutations and/or deletions of *TP53* are associated with resistance to treatments in multiple myeloma as in most B-cell malignancies (1). *TP53* mutation and deletion are associated in B-cell malignancies suggesting that bi-allelic alterations of *TP53* are involved in resistance, although overall survival of patients with lymphoma or myeloma appears more significantly related to mutations than to deletion (2, 3). Mutations of *TP53* induced a more rapid development of spontaneous tumors than the deletion of *TP53* (4). Moreover, some mutations are characterized by a gain of function, making mutant forms of the p53 protein interesting therapeutic targets (5). Given the importance of folding for p53 activity and the existence of temperature-dependent mutations, chemical molecules able to change the conformation of p53 and to restore its transcriptional activity were screened (6). During the last 15 years, several molecules were isolated for their efficacy to induce cell death in *TP53* mutated cells and some of these molecules were shown to interact with the mutant p53 protein (7, 8). However, the p53 dependency of several p53 reactivating molecules, such as RITA and Prima-1^{Met}, is debated as both drugs killed cancer cells independently from *TP53* status and p53 expression (9, 10). Indeed, it was recently demonstrated


115 using CRISPR/Cas9 technology that the cell response to RITA,
 116 which is a DNA damaging drug, depended on FANCD2
 117 expression (11). On the other hand, Prima-1^{Met} has been
 118 shown to decrease glutathione and to induce ROS independently
 119 from p53 expression or mutations (10, 12, 13), at least
 120 by directly interfering with thioredoxin reductase, a central
 121 enzyme of the detoxifying redox pathway (14). These results
 122 prompted us to further investigate whether auranofin, an
 123 inhibitor of the thioredoxin reductase, displayed a Prima-
 124 1^{Met}-like activity. We therefore investigated activity and death
 125 mechanism of auranofin in myeloma cell lines and primary
 126 myeloma cells characterized for *TP53* status. We showed that
 127 activity of auranofin and Prima-1^{Met} correlated in myeloma

172 cells and that both drugs induced a Bax/Bak-independent
 173 cell death.
 174
 175
 176
 177
 178

MATERIALS AND METHODS

Human Myeloma Cell Lines (HMCLs) and Primary Samples

179 Eighteen HMCLs used for this study, i.e., 7 HMCLs with
 180 a wild-type *TP53* status (MDN, NCI-H929, NAN9, NAN11,
 181 XG3, XG6, XG7), 8 HMCLs with a missense *TP53* mutation
 182 (JIM3, KMS12PE, LP1, NAN10, OPM2, U266, XG2, XG5) and
 183 3 HMCLs with a *TP53* indel leading to the lack of mRNA and/or
 184 protein expression (JJN3, L363, NAN7). All HMCLs have been
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222



Q4
 Q5

extensively characterized (10, 15, 16). *TP53* status was performed by direct sequencing of RT-PCR products (16) and by whole exon sequencing (17). After obtaining informed consent, blood or bone marrow samples from patients with MM were collected at the Department of Hematology of the Nantes University Hospital (MYRACLE cohort, ethical approval NCT03807128, Benaniba et al., submitted). Plasma cells were obtained after gradient density centrifugation and FISH was performed as previously described (9, 18).

Reagents and Antibodies

Prima-1^{Met} was purchased from Santa Cruz Biotechnology (CliniSciences, Nanterre, France), L-buthionine sulfoximine (BSO), auranofin and N-acetyl-L-cysteine were purchased from Sigma-Aldrich (Saint-Quentin Fallavier, France). Anti-CD138-PE monoclonal antibody was purchased from Beckman Coulter (Villepinte, France), Annexin V-FITC was purchased from ImmunoTools (Friesoythe, Germany), anti-Bax and anti-actin were purchased from BD-Biosciences (Le Pont de




FIGURE 2 | Auranofin and Prima-1^{Met} induced ROS-dependent BAX/BAK-independent cell death. **(A)** L363 and OPM2 cell lines (500 000 cells/ml) were incubated for 2 days with auranofin (2000 and 1000 nM, respectively) or Prima-1^{Met} (80 and 40 μM, respectively) with or without NAC (0.5 mM). Cell death was assessed by Annexin V staining. The data represent 3 independent experiments. The statistical analyses were performed using the Wilcoxon matched-pairs signed-rank test. **(B)** LP1 cells were transfected with siRNA against *BAX* or *BAK1* for 2 days and sensitivity to each drug was assessed. The statistical analyses were performed using the two-way ANOVA test with multiple comparisons. **(C)** Auranofin and Prima-1^{Met} LD₅₀ values were determined in XG5 parental cells and in venetoclax resistant XG5-199R cells. Cells were incubated with increasing concentrations of auranofin or Prima-1^{Met} with or without 500 μM BSO for 2 days, and cell death was assessed by Annexin V staining.

Claix, France), Cell Signaling (Ozyme, Montigny-le-Bretonneux, France) and Millipore (Guyancourt, France), respectively.

siRNA Experiments

Transient *BAX* or *BAK* silencing was performed in LP1 myeloma cells (100 pmol siRNA/3 × 10⁶ cells) using lipofectamine RNAiMax (Thermo Fischer Scientific, Saint-Herblain, France), as previously reported (19).

Cell Death Assay

The cell lines or mononuclear cells from patients' samples (500,000 cells/ml) were incubated with Prima-1^{Met} or auranofin with different concentrations as indicated within the legends of the figures. Cell death was assessed by Annexin V staining in cell lines and by the loss of CD138 staining in primary myeloma cells (10, 19–21). The fluorescence acquisition and analysis were performed using a FACsCalibur with Cell Quest




FIGURE 3 | Primary myeloma cells were similarly sensitive to Prima-1^{Met} and to auranofin. **(A)** Mononuclear cells from patient #4 with myeloma were incubated for 24 hours with or without (-) Prima-1^{Met} (5 or 10 µM) or auranofin (5 or 10 µM) and stained with anti-CD138-PE. Viability was assessed by CD138 expression. **(B)** Mononuclear cells from 10 samples were incubated for 24 hours with Prima-1^{Met} (5 or 10 µM) or auranofin (5 or 10 µM) with or without BSO (250 µM), and stained with anti-CD138-PE. The statistical analyses were performed using the Wilcoxon matched-pairs signed-rank test. **(C)** Cell death induced by Prima-1^{Met} (5 µM, samples #3,4,5,6,8,9, and 10), auranofin (50nM, samples #3,4,5,6,7,9, and 10) or BSO (250 µM, samples #3-10) was compared to cell death induced by combinations (combo) of Prima-1^{Met} or auranofin with BSO. The statistical analyses were performed using the Wilcoxon matched-pairs signed-rank test. **(D)** Cell death induced by auranofin (50 nM) or Prima-1^{Met} (5 µM) was analyzed according to del(17p) status. The statistical analyses were performed using the Mann-Whitney test. **(E)** Myeloma cell death induced by 5 µM of Prima-1^{Met} was plotted against cell death induced by 50 nM auranofin. Correlation was performed using the Pearson test.

TABLE 1 | Prima-1^{Met} and Auranofin induced cell death in primary myeloma cells.

N°	Sample	Status	del(17p)	CD138 ⁺ cells (%)										
				Prima-1 ^{Met} 5 μM	Prima-1 ^{Met} 10 μM	Auranofin 50 nM	Auranofin 500 nM	BSO 250 μM	Prima-1 ^{Met} 5 μM + BSO	Prima-1 ^{Met} 10 μM + BSO	Auranofin 50 nM + BSO	Auranofin 500 nM + BSO		
1	MM		nd	0.1	0.1	nd	0.4	nd	nd	nd	nd	nd	nd	nd
2	MM	-	-	0.3	0.3	nd	0.1	nd	nd	nd	nd	nd	nd	nd
3	PCL	+	+	nd	nd	6.9	nd	22.4	3.3	nd	nd	0.6	nd	nd
4	MM	+	+	1.0	1.0	6.0	0.0	7.9	0.5	0.2	0.4	0.2	0.2	0.2
5	MM	-	-	4.0	4.0	16.4	0.3	16.1	3.0	0.3	0.3	0.1	0.1	0.1
6	MM	-	-	5.0	5.0	19.9	12.5	20.0	3.6	0.1	12.5	0.1	0.1	0.1
7	MM	-	-	0.1	0.1	5.6	1.1	6.6	0.1	0.1	0.3	0.2	0.2	0.2
8	PCL	-	-	0.2	0.2	18.2	0.4	50.1	0.2	0.2	nd	0.2	0.2	0.2
9	PCL	+	+	16.1	16.1	40.0	32.3	35.7	8.9	1.4	19.6	1.0	1.0	1.0
10	MM	-	-	0.4	0.4	49.2	36.6	48.0	0.6	0.4	33.2	0.4	0.4	0.4

Peripheral blood or bone marrow samples were obtained from patients with plasma cell leukemia (PCL) or Multiple Myeloma (MM).

Deletion of the short arm of chromosome 17, del(17p), was performed by FISH.

Mononuclear cells were incubated overnight with Prima-1^{Met}, Auranofin with or without BSO, as indicated within the table. nd, not done.

(Becton Dickinson) or FlowJo (Ashland, OR, USA) software, Cytozell core facility (SFR Bonamy, Nantes, France).

Statistical Analyses

The statistical analyses were performed using GraphPad Prism 7.

RESULTS

Sensitivity of Myeloma Cell Lines to Prima-1^{Met} and Auranofin Correlated

We assessed the efficacy of auranofin, an inhibitor of thioredoxin reductase, in comparison with Prima-1^{Met} in 18 HMCLs. We determined the lethal dose 50 (LD₅₀) of auranofin and Prima-1^{Met} to HMCLs using Annexin V staining at day 2, as illustrated in **Figure 1A**. **Figure 1B** (left panel) shows a positive correlation between LD₅₀ values for auranofin and Prima-1^{Met} ($p = 0.042$, $r^2 = 0.2327$, Pearson test); however, auranofin displayed higher activity (about 80-fold) compared to Prima-1^{Met} (median values were 0.4 and 32.5 μM, respectively). Notably, the correlation was essentially supported by HMCLs displaying abnormal *TP53* status (**Figure 1B**, middle and right panels, Pearson test), although activity of auranofin and Prima-1^{Met} was not dependent on p53 mutations or expression (**Figure 1C**). Using the CellMinerCDB portal (<https://discover.nci.nih.gov/cellminerfdb/>), which provides pharmacologic sensitivity of cancer cell lines, we confirmed that activity of another inhibitor of the thioredoxin reductase (PX-12) also correlated with activity of Prima-1^{Met} in myeloma cell lines (**Figure S1**).

Because it was previously shown that Prima-1^{Met} synergized with BSO, an irreversible inhibitor of GSH synthesis, we determined whether auranofin also synergized with BSO (10, 12). Six HMCLs (JJN3, MDN, OPM2, XG5, XG6, and U266) were cultured with increasing concentrations of Prima-1^{Met} or auranofin with or without BSO (500 μM) and LD₅₀ values were determined. Both Prima-1^{Met} and auranofin strongly synergized with BSO, and LD₅₀ values were decreased by ~ 4- and 10-fold, respectively (**Figure 1D**).

Auranofin and Prima-1^{Met} Induced ROS-Dependent Bax/Bak-Independent Cell Death

We previously demonstrated that Prima-1^{Met} induced ROS-dependent cell death in myeloma cells that was prevented by N-acetyl-L-cysteine (NAC) (10). Because auranofin has been shown to induce ROS production, we thus assessed whether NAC was also able to inhibit cell death induced by auranofin (22). As shown in **Figure 2A** in L363 and OPM2, the addition of NAC inhibited cell death induced by auranofin and Prima-1^{Met} by 86% ($p = 0.03$) and 95% ($p = 0.03$), respectively. Prima-1^{Met} was shown to induce apoptosis that was independent from Bax/Bak: Prima-1^{Met} induced lipid peroxidation that mediated mitochondrial permeabilization, cytochrome C release and activation of caspases (13). To determine the role of Bax/Bak in auranofin-induced cell death, we performed *BAX/BAK1* silencing in LP1 cells. A decrease in Bax or Bak expression did not significantly modify cell responses to either drug (**Figure 2B**).

Q5

Moreover, the venetoclax-resistant XG5-199R HMCL, in which the expression of both apoptosis executors was lost, remained as sensitive as the parental XG5 cell line to both drugs (Figure 2C) (23). These results showed that auranofin and Prima-1^{Met} induced ROS-dependent, Bax/Bak-independent cell death.

Sensitivity of Primary Myeloma Cells to Prima-1^{Met} and Auranofin Correlated

We assessed the activity of each drug with or without BSO in 10 samples from patients with either multiple myeloma (MM) or plasma cell leukemia (PCL) with or without the deletion of the short arm of chromosome 17, del(17p). Mononuclear cells were incubated with different concentrations of Prima-1^{Met} or auranofin with or without BSO (250 μM). Cell death was determined by the loss of CD138 expression, Figure 3A. Indeed, cell death could not be monitored by Annexin V staining as the loss of myeloma viability was associated with the loss of expression of the plasma cell specific CD138 expression, as illustrated in Figure S2 (10, 19, 20). Both drugs significantly induced myeloma cell death (Figure 3B; Table 1). The median values of cell death induced by Prima-1^{Met} (5 or 10 μM) were 68% ($p = 0.0020$) and 97% ($p = 0.0039$), respectively, and the median values of cell death induced by auranofin (50 or 500 nM) were 20% ($p = 0.0391$) and 97% ($p = 0.0039$), respectively. Although BSO (250 μM) induced a weak decrease in cell viability (median cell death 7.5%, $p = 0.0312$, Figure 3B), it significantly synergized with both Prima-1^{Met} (5 μM) and auranofin (50 nM). The cell death median values of the combination of BSO with Prima-1^{Met} or with auranofin vs. the sum of cell death induced by Prima-1^{Met} or auranofin plus BSO were 85.2% vs. 62.8% ($n = 7$, $p = 0.0156$, 1.24-fold increase) and 95.6% vs. 12.7% ($n = 7$, $p = 0.0156$, 6.41-fold increase), respectively (Figure 3C). The sensitivity of samples to both drugs was not different in samples with or without the deletion of the short arm of chromosome 17, Figure 3D. Auranofin (50 nM) and Prima-1^{Met} (5 μM) displayed correlated activity in myeloma samples ($n = 8$, $r^2 = 0.5963$, $p = 0.0306$), Figure 3E.

CONCLUSION

In this paper, we showed that the sensitivity of myeloma cells to auranofin correlated with sensitivity to Prima-1^{Met} in both HMCLs and in primary myeloma cells from patients

REFERENCES

1. Tessoulin B, Eveillard M, Lok A, Chiron D, Moreau P, Amiot M, et al. p53 dysregulation in B-cell malignancies: more than a single gene in the pathway to hell. *Blood Rev.* (2017) 31:251–9. doi: 10.1016/j.blre.2017.03.001
2. Eskelund CW, Dahl C, Hansen JW, Westman M, Kolstad A, Pedersen LB, et al. TP53 mutations identify younger mantle cell lymphoma patients who do not benefit from intensive chemoimmunotherapy. *Blood* (2017) 130:1903–10. doi: 10.1182/blood-2017-04-779736
3. Walker BA, Mavrommatis K, Wardell CP, Ashby TC, Bauer M, Davies F, et al. A high-risk, double-Hit, group of newly diagnosed myeloma identified by genomic analysis. *Leukemia* (2018).

with MM or PCL, and that auranofin was more efficient than Prima-1^{Met}. Although p53 competent HMCLs were very sensitive to both drugs, the activity of Prima-1^{Met} or auranofin was not dependent on TP53 status, highlighting that targeting the ROS/GSH pathway was efficient in cells expressing mutant p53 protein or lacking p53 expression. Prima-1^{Met} and auranofin induced Bak/Bax-independent cell death and were efficient in myeloma cells resistant to the Bcl2-specific BH3-mimetic venetoclax. These data showed that auranofin, as Prima-1^{Met}, overcomes resistance to cell death mediated by either p53 deficiency or by mitochondrial loss of priming in myeloma cells. Both drugs appear thus of particular interest for resistance *in vivo*. Prima-1^{Met} (APR-246) is under clinical evaluation in ovarian cancers with mutated TP53 or in refractory (TP53-mutated) myeloid neoplasms, alone or in combination. Auranofin, which was used to treat patients with arthritis, was recently shown to be able to eliminate side populations and to enhance ibrutinib efficacy in solid cancer cells (22, 24). These recent findings are in favor of assessing auranofin in myeloma patients resistant to current therapies as well as resistant to the Bcl2-specific BH3 mimetics venetoclax.

AUTHOR CONTRIBUTIONS

BT designed the study, performed experiments, and participated in the writing of the article. GD performed experiments. CD and MA provided XG5 venetoclax-resistant cells. CP-D designed the study and wrote the article. All authors approved the manuscript.

ACKNOWLEDGMENTS

This work was supported by grants from the Ligue Régionale Contre le Cancer, DHU Oncogreff, AF3M, Actions Cancer 44, SIRIC ILIAD (INCa-DGOS-Inserm_12558), and i-Site NexT (ANR-16-IDEX-0007). BT was supported by Inserm and ARC (poste d'accueil).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fonc.2019.00128/full#supplementary-material>

4. Xu J, Qian J, Hu Y, Wang J, Zhou X, Chen H, et al. Heterogeneity of Li-Fraumeni syndrome links to unequal gain-of-function effects of p53 mutations. *Sci Rep.* (2014) 4:4223. doi: 10.1038/srep04223
5. Hanel W, Marchenko N, Xu S, Yu SX, Weng W, Moll U. Two hot spot mutant p53 mouse models display differential gain of function in tumorigenesis. *Cell Death Differ* (2013) 20:898–909. doi: 10.1038/cdd.2013.17
6. Bykov VJN, Eriksson SE, Bianchi J, Wiman KG. Targeting mutant p53 for efficient cancer therapy. *Nat Rev Cancer* (2018) 18:89–102. doi: 10.1038/nrc.2017.109
7. Zhang Q, Bergman J, Wiman KG, Bykov VJN. Role of Thiol Reactivity for Targeting Mutant p53. *Cell Chem Biol.* (2018) 25:1219–30 e3. doi: 10.1016/j.chembiol.2018.06.013

- 685 8. Zhang Q, Bykov VJN, Wiman KG, Zawacka-Pankau J. APR-246 reactivates
686 mutant p53 by targeting cysteines 124 and 277. *Cell Death Dis.* (2018) 9:439.
687 doi: 10.1038/s41419-018-0463-7
- 688 9. Surget S, Descamps G, Brosseau C, Normant V, Maiga S, Gomez-Bougie P,
689 et al. RITA (Reactivating p53 and Inducing Tumor Apoptosis) is efficient
690 against TP53 abnormal myeloma cells independently of the p53 pathway. *BMC*
691 *Cancer* (2014) 14:437. doi: 10.1186/1471-2407-14-437
- 692 10. Tessoulin B, Descamps G, Moreau P, Maiga S, Lode L, Godon C,
693 et al. PRIMA-1Met induces myeloma cell death independent of
694 p53 by impairing the GSH/ROS balance. *Blood* (2014) 124:1626–36.
695 doi: 10.1182/blood-2014-01-548800
- 696 11. Wanzel M, Vishedyk JB, Gittler MP, Gremke N, Seiz JR, Hefter M, et al.
697 CRISPR-Cas9-based target validation for p53-reactivating model compounds.
698 *Nat Chem Biol.* (2016) 12:22–8. doi: 10.1038/nchembio.1965
- 699 12. Lambert JM, Gorzov P, Veprintsev DB, Soderqvist M, Segerback D,
700 Bergman J, et al. PRIMA-1 reactivates mutant p53 by covalent binding
701 to the core domain. *Cancer Cell* (2009) 15:376–88. doi: 10.1016/j.ccr.2009.
702 03.003
- 703 13. Liu DS, Duong CP, Haupt S, Montgomery KG, House CM, Azar WJ,
704 et al. Inhibiting the system xC(-)/glutathione axis selectively targets
705 cancers with mutant-p53 accumulation. *Nat Commun.* (2017) 8:14844.
706 doi: 10.1038/ncomms14844
- 707 14. Peng X, Zhang MQ, Conserva F, Hosny G, Selivanova G, Bykov VJ, et al.
708 APR-246/PRIMA-1MET inhibits thioredoxin reductase 1 and converts the
709 enzyme to a dedicated NADPH oxidase. *Cell Death Dis.* (2013) 4:e881.
710 doi: 10.1038/cddis.2013.417
- 711 15. Maiga S, Brosseau C, Descamps G, Dousset C, Gomez-Bougie P, Chiron D,
712 et al. A simple flow cytometry-based barcode for routine authentication of
713 multiple myeloma and mantle cell lymphoma cell lines. *Cytometry A.* (2015)
714 87:285–8. doi: 10.1002/cyto.a.22643
- 715 16. Moreaux J, Klein B, Bataille R, Descamps G, Maiga S, Hose D, et al. A high-risk
716 signature for patients with multiple myeloma established from the molecular
717 classification of human myeloma cell lines. *Haematologica* (2011) 96:574–82.
718 doi: 10.3324/haematol.2010.033456
- 719 17. Tessoulin B, Moreau-Aubry A, Descamps G, Gomez-Bougie P, Maiga S,
720 Gaignard A, et al. Whole-exon sequencing of human myeloma cell lines
721 shows mutations related to myeloma patients at relapse with major hits in
722 the DNA regulation and repair pathways. *J Hematol Oncol.* (2018) 11:137.
723 doi: 10.1186/s13045-018-0679-0
- 724 18. Lode L, Eveillard M, Trichet V, Soussi T, Wulleme S, Richebourg
725 S, et al. Mutations in TP53 are exclusively associated with
726 del(17p) in multiple myeloma. *Haematologica* (2010) 95: 1973–6.
727 doi: 10.3324/haematol.2010.023697
- 728 19. Lok A, Descamps G, Tessoulin B, Chiron D, Eveillard M, Godon C, et al. p53
729 regulates CD46 expression and measles virus infection in myeloma cells. *Blood*
730 *Adv.* (2018) 2:3492–505. doi: 10.1182/bloodadvances.2018025106
- 731 20. Surget S, Chiron D, Gomez-Bougie P, Descamps G, Menoret E, Bataille R, et al.
732 Cell death via DR5, but not DR4, is regulated by p53 in myeloma cells. *Cancer*
733 *Res.* (2012) 72:4562–73. doi: 10.1158/0008-5472.CAN-12-0487
- 734 21. Gomez-Bougie P, Maiga S, Tessoulin B, Bourcier J, Bonnet A, Rodriguez
735 MS, et al. BH3-mimetic toolkit guides the respective use of BCL2 and
736 MCL1 BH3-mimetics in myeloma treatment. *Blood* (2018) 132:2656–69.
737 doi: 10.1182/blood-2018-03-836718
- 738 22. Hou GX, Liu PP, Zhang S, Yang M, Liao J, Yang J, et al. Elimination of stem-
739 like cancer cell side-population by auranofin through modulation of ROS and
740 glycolysis. *Cell Death Dis.* (2018) 9:89. doi: 10.1038/s41419-017-0159-4
- 741 23. Dousset C, Maiga S, Gomez-Bougie P, Le Coq J, Touzeau C, Moreau P, et al.
742 BH3 profiling as a tool to identify acquired resistance to venetoclax in multiple
743 myeloma. *Br J Haematol.* (2017) 179:684–8. doi: 10.1111/bjh.14251
- 744 24. Hu J, Zhang H, Cao M, Wang L, Wu S, Fang B. Auranofin Enhances Ibrutinib's
745 Anticancer Activity in EGFR-Mutant Lung Adenocarcinoma. *Mol Cancer*
746 *Ther.* (2018) 17:2156–63. doi: 10.1158/1535-7163.MCT-17-1173
- 747 **Conflict of Interest Statement:** The authors declare that the research was
748 conducted in the absence of any commercial or financial relationships that could
749 be construed as a potential conflict of interest.
- 750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798