



HAL
open science

Can we learn from coupling EEG-fMRI to enhance neuro-feedback in EEG only?

Claire Cury, Pierre Maurel, Rémi Gribonval, Christian Barillot

► To cite this version:

Claire Cury, Pierre Maurel, Rémi Gribonval, Christian Barillot. Can we learn from coupling EEG-fMRI to enhance neuro-feedback in EEG only?. OHBM 2019 - Annual Meeting Organization for Human Brain Mapping, Jun 2019, Rome, Italy. pp.1. inserm-02074623

HAL Id: inserm-02074623

<https://inserm.hal.science/inserm-02074623v1>

Submitted on 5 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Can we learn from coupling EEG-fMRI to enhance neuro-feedback in EEG only?

C. Cury^{1,2}, P. Maurel¹, R. Gribonval² and C. Barillot¹

1 – CNRS, Inserm, Inria, Univ Rennes, IRISA, Empenn team

2 – CNRS, Inria, Univ Rennes, IRISA, Panama team



PANAMA
Partnership and New Algorithms
for Audio and Signal Modeling

EMPENN




INTRODUCTION

Neuro-feedback (NF) : Learn to control your brain with your brain.

EEG and fMRI, grounds solutions in the context of **brain rehabilitation** protocols.

EEG and fMRI provide complementary information.

EEG is easy to use, fMRI is a costly and exhausting for patients modality




Bi-modal NF :

- Records and synchronises EEG and fMRI signals, in real time (Mano et al).
- Combines NF-EEG and NF-fMRI advantages
- Improve the quality of NF sessions (Perronnet et al).
- It is not portable or easy to use, due to the fMRI modality.

→ Can we enhance NF in EEG only, from a previous bi-modal NF session ?

METHOD

- **Design and strategy** : Machine learning mechanism based on bimodal NF scores and EEG signals.




- **Model** : Non linear structured design matrix X

$$X = [X_0; X_3; X_4; X_5], \in \mathbb{R}^{T \times 4E \times B}, \text{ with } X_i \in \mathbb{R}^{T \times E \times B}$$

$$X_0(t, e, b) = \text{Freq}(EEG(e, I_t), F_b), \forall t \in \{1, \dots, T\} \text{ and } \forall b \in \{1, \dots, B\}$$

$$\left. \begin{aligned} X_3(\cdot, e, b) &= X_0(\cdot, e, b) * \text{HRF}(3) \\ X_4(\cdot, e, b) &= X_0(\cdot, e, b) * \text{HRF}(4) \\ X_5(\cdot, e, b) &= X_0(\cdot, e, b) * \text{HRF}(5) \end{aligned} \right\} \forall e \in \{1, \dots, E\}, \text{ and } \forall b \in \{1, \dots, B\}$$



- **Optimisation** : structured sparse regularisation following 3 conditions:

1. Spatial sparsity
2. Smooth across frequency bands
3. Group selection of frequency bands.

$$\hat{\alpha} = \arg \min_{\alpha} \sum_{t=1}^T \frac{1}{2} (NF(t) - \langle X(t), \alpha \rangle)^2 + \lambda \underbrace{\|\alpha\|_{21}}_{\text{Cond 1. and 2.}} + \rho \underbrace{\|\alpha\|_1}_{\text{Cond 3.}}$$




RESULTS

- **SIGNIFICANT INFORMATION FROM NF-fMRI CAN BE CAPTURED BY THE MODEL, AND ENHANCE EEG ONLY NEUROFEEDBACK.**
- PREDICTION WITH **NF-PREDICTOR 5** WITH A MEDIAN CORRELATION OF 0.74

• Method tested on 17 subjects with 3 bimodal neuro-feedback sessions of motor imagery tasks.

• We tested 5 NF-predictors:

1. $\tilde{y}_{\alpha_c}(t) = \langle X, \hat{\alpha}_c \rangle$, learned from X and $NF_c = NF\text{-EEG} + NF\text{-fMRI}$
2. $\tilde{y}_{\alpha_e}(t) = \langle X, \hat{\alpha}_e \rangle$, learned from X_0 and NF-EEG
3. $\tilde{y}_{\alpha_f}(t) = \langle X, \hat{\alpha}_f \rangle$, learned from X_d and NF-fMRI
4. $\tilde{y}_{\alpha_e}(t) + \tilde{y}_{\alpha_f}(t)$
5. $y_e(t) + \tilde{y}_{\alpha_f}(t)$, with $y_e(t) = NF\text{-EEG}(t)$



References :

Abreu, R., Leal, A., Figueiredo, P. EEG-Informed fMRI: A Review of Data Analysis Methods. *Frontiers in Human Neuroscience*, no. 12 (Feb 2018)

Gramfort, A., Strohmeier, D., Haueisen, J., Hamalainen, M., Kowalski, M.: *Functional Brain Imaging with M/EEG Using Structured Sparsity in Time-Frequency Dictionaries*. IPMI, vol. 6801, pp. 600–611. (2011)

Mano, M., L'ecuyer, A., Bannier, E., Perronnet, L., Noorzadeh, S., Barillot, C.: How to Build a Hybrid Neurofeedback Platform Combining EEG and fMRI. *Frontiers in Neuroscience* 11 (2017)

Meir-Hasson, Y., Kinreich, S., Podlipsky, I., Hendler, T., Intrator, N.: An EEG Finger-Print of fMRI deep regional activation. *NeuroImage* 102, 128–141 (2014)

Noorzadeh, S., Maurel, P., Oberlin, T., Gribonval, R., Barillot, C.: Multi-modal EEG and fMRI Source Estimation Using Sparse Constraints. *MICCAI* (2017)

Perronnet, L., L'ecuyer, A., Mano, M., Bannier, E., Lotte, F., Clerc, M., Barillot, C.: Unimodal Versus Bimodal EEG-fMRI Neurofeedback of a Motor Imagery Task. *Frontiers in Human Neuroscience* 11 (Apr 2017)

Zotev, V., Self-regulation of human brain activity using simultaneous real-time fMRI and EEG neurofeedback. *NeuroImage* 85, pp. 985–995 (Jan 2014)