



## Association between genetic obesity susceptibility and mother-reported eating behavior in children up to 5 years

Blandine de Lauzon-Guillain, Yves Akoli Koudou, Jérémie Botton, Anne Forhan, Sophie Carles, Véronique Pelloux, Karine Clément, Ken K. Ong, Marie Aline Charles, Barbara Heude

### ► To cite this version:

Blandine de Lauzon-Guillain, Yves Akoli Koudou, Jérémie Botton, Anne Forhan, Sophie Carles, et al.. Association between genetic obesity susceptibility and mother-reported eating behavior in children up to 5 years: genetic obesity risk and child's eating behavior. *International Journal of Pediatric Obesity*, 2019, 14 (5), pp.e12496. 10.1111/ijpo.12496 . inserm-02093066

**HAL Id: inserm-02093066**

**<https://inserm.hal.science/inserm-02093066>**

Submitted on 8 Apr 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Association between genetic obesity susceptibility and mother-reported eating behavior in children up to 5 years

Blandine de Lauzon-Guillain, PhD<sup>1,2,8</sup>, Yves Akoli Koudou, MSc<sup>1</sup>, Jérémie Botton, PhD<sup>1,3</sup>, Anne Forhan, MSc<sup>1,2</sup>, Sophie Carles, PhD<sup>1</sup>, Véronique Pelloux, PhD<sup>4,5,6</sup>, Karine Clément, PhD<sup>4,5,6</sup>, Ken K. Ong, FRCPCH<sup>7</sup>, Marie Aline Charles, MD, PhD<sup>1,2</sup>, Barbara Heude, PhD<sup>1,2</sup>; on behalf of the EDEN Mother-Child Cohort Study Group\*

**\* Members of the EDEN Mother-Child Cohort Study Group:** I. Annesi-Maesano, JY.

Bernard, J. Botton, M.A. Charles, P. Dargent-Molina, B. de Lauzon-Guillain, P. Ducimetière, M. de Agostini, B. Foliguët, A. Forhan, X. Fritel, A. Germa, V. Goua, R. Hankard, B. Heude, M. Kaminski, B. Larroque†, N. Lelong, J. Lepeule, G. Magnin, L. Marchand, C. Nabet, F. Pierre, R. Slama, M.J. Saurel-Cubizolles, M. Schweitzer, O. Thiebaugeorges.

### *Affiliations*

<sup>1</sup> INSERM, UMR1153 Epidemiology and Biostatistics Sorbonne Paris Cité Center, Early ORigin of the Child's Health and Development Team (ORCHAD), Paris, France

<sup>2</sup> Paris Descartes University, France

<sup>3</sup> Univ. Paris-Sud, Université Paris-Saclay, Faculty of Pharmacy, F-92296, Châtenay-Malabry, France;

<sup>4</sup> Institute of Cardiometabolism and Nutrition, ICAN, F-75013, Paris, France;

<sup>5</sup> INSERM, UMRS 1166, Nutriomic team 6, Paris, F-75013 France;

<sup>6</sup> Sorbonne Universités, UPMC Université Paris 06, UMRS1166, Paris, F-75013 France;

<sup>7</sup> Medical Research Council Epidemiology Unit & Department of Paediatrics, University of Cambridge, Addenbrooke's Hospital, Cambridge, England

<sup>8</sup> INRA, U1125 Epidemiology and Biostatistics Sorbonne Paris Cité Center (CRESS), Early ORigin of the Child's Health and Development Team (ORCHAD), Paris, F-75014 France.

***PubMed indexing.***

de Lauzon-Guillain, Koudou, Botton, Forhan, Carles, Pelloux, Clément, Ong, Charles, Heude;  
on behalf of the EDEN Mother-Child Cohort Study Group

***Running head***

genetic obesity risk and child's eating behavior

***Keywords***

obesity, genetic, eating behaviour, growth, birth cohort, childhood

***Corresponding author***

de Lauzon-Guillain Blandine

INSERM CRESS – Eq6 ORCHAD

16 av. Paul Vaillant Couturier, 94807 Villejuif Cedex, FRANCE

Tel: +33145595019; Fax: +33147269454; E-mail: [blandine.delauzon@inserm.fr](mailto:blandine.delauzon@inserm.fr)

***Abbreviations***

BMI-GRS: combined obesity risk-allele score

SNP: single-nucleotide polymorphisms

## **Abstract**

**Background.** Many genetic polymorphisms identified by genome wide association studies for adult BMI have been suggested to regulate food intake.

**Objective.** The objective was to study the associations between a genetic obesity risk score, appetitive traits and growth of children up to age 5 years, with a longitudinal design.

**Methods.** In 1,142 children from the EDEN birth cohort, a combined obesity risk-allele score (BMI-GRS) was related to appetitive traits (energy intake up to 12 months, a single item on appetite from 4 months to 3 years, a validated appetite score at 5 years) using Poisson regressions with robust standard errors. The potential mediation of appetitive traits on the association between BMI-GRS and growth was assessed by the Sobel test.

**Results.** Children with a high BMI-GRS were more likely to have high energy intake at 1 year and high appetite at 2 and 5 years. High energy intake in infancy and high appetite from 1 year were related to higher subsequent BMI. High 2-y appetite seemed to partially mediate the associations between BMI-GRS and BMI from 2 to 5 years (all  $p \leq 0.05$ ).

**Conclusions.** Genetic susceptibility to childhood obesity seems to be partially explained by appetitive traits in infancy, followed by an early childhood rise in BMI.

## 18 **Introduction**

19 Genome-wide-association studies (GWAs) have identified several genes associated with an  
20 increased risk of obesity (1, 2). Time has now come to take advantage of these major findings  
21 in order to better understand the mechanisms of lifecourse obesity development from infancy  
22 to adulthood. Many birth cohorts have shown that scores of genetic susceptibility to obesity  
23 are associated with early growth, as early as 2 years in some of them, but not with foetal  
24 growth (3, 4, 5, 6).

25 A great part of the genes implicated by GWAs, and especially those from monogenic studies  
26 on severe childhood or early onset obesity, are purportedly involved in the central regulation  
27 of food intake (7). Furthermore, the genetic component of eating behavior has clearly been  
28 established through heritability studies, in adults (8) and also in children (9, 10). Eating  
29 behavior even in infancy may influence later adiposity development (11), which raises the  
30 question whether the association between the genetic score and adiposity is mediated by  
31 eating behaviour in childhood. Cross-sectional studies, conducted among 8-11 years children,  
32 have already shown that increased BMI of children homozygous for the at-risk allele of a  
33 FTO SNP may be partially mediated by appetitive traits (12, 13). Similarly, a cross-sectional  
34 analysis of the Twins Early Development Study highlighted that satiety responsiveness  
35 partially mediated the association between a polygenic risk score and adiposity in children  
36 aged 8-11 years(14). However, such a cross-sectional design is insufficient to establish a  
37 causal pathway. More recently, this causal pathway has been tested longitudinally within a  
38 Norwegian cohort following children from the age of 4 years to the age of 8 years (15). In that  
39 study, higher genetic risk for obesity was associated with appetitive traits at 6 years, but these  
40 appetitive traits were not related to BMI gain up to 8 years. As a previous study has shown  
41 that appetitive traits assessed at age 3 months predicted weight at 9 months, it would be  
42 important to test the potential mediating effect of infant or toddler's appetitive traits on

subsequent growth. A prospective design would also offer the opportunity to highlight a potential window of opportunity for this mediating effect.

In this context, we aimed at investigating within the French EDEN mother-child cohort, how genetic predisposition to obesity may influence eating behavior very early life and whether this relationship might explain the accelerated growth in children with higher genetic risk for obesity. In the present study, we focused on three aspects of child's eating behavior that could be assessed longitudinally, from 4 months to 5 years: energy intake and appetite.

## **Material and methods**

### ***Study population***

The EDEN mother-child study is a prospective cohort whose aim is to assess pre- and post-natal determinants of child growth, development and health. It has been described in more detail elsewhere (16). In brief, 2002 pregnant women were recruited from 2003 to 2006 in two French university hospitals, before 24 weeks of amenorrhea. Exclusion criteria were multiple pregnancies, known diabetes prior to pregnancy, illiteracy, plan to move outside the region within the next three years. The study was approved by the Ethics Committee of the University Hospital of Kremlin-Bicêtre on December 12, 2002 and data files were declared to the National Committee for Processed Data and Freedom. Written consent was obtained from both parents.

The primary endpoints of the cohort were pre and post-natal growth, blood pressure, allergies, infectious diseases, mental health and cognitive development. Dietary behavior was initially collected as one of the main exposure factors in the cohort and was considered as a potential mediating factor in the present study.

Data collected during pregnancy and at birth, include sociodemographic variables, maternal smoking, gestational diabetes, parental anthropometric measurements, and newborn characteristics (sex, gestational age, birthweight) (17). At four, eight and twelve months after birth, mothers completed mailed questionnaires that provided detailed information on the feeding method (exclusive breast-feeding, exclusive formula-feeding or mixed-diet) between birth and four months, then at eight and twelve months as well as the date of breastfeeding cessation. Breastfeeding duration was derived from these data (18). Clinical exams were also conducted at birth, 1 year, 3 years and 5 years and were dedicated to collection of biological sample, cognitive assessments by psychologists and measurement of anthropometric and clinical parameters.

#### ***Genotyping and BMI Genetic Risk Score (BMI-GRS) generation***

DNA was extracted from cord blood samples collected at birth. Genotypes at 16 single-nucleotide polymorphisms (SNPs) were measured at the Medical Research Council Epidemiology Unit, Cambridge (iPLEX platform; Sequenom), as previously described (19). Among the 32 loci identified by Speliotes et al. as having genome-wide significant associations with BMI in adults (2), we considered in the present study the 16 SNPs also showing associations with childhood BMI either in that original report (2) or in subsequent data (4); they lie in or near NRXN3 (rs10146997), SLC39A8 (rs13107325), TNNT3 (rs1514175), PTBP2 (rs1555543), MC4R (rs17782313), FLJ35779 (rs2112347), NEGR1 (rs2568958), RPL27A (rs4929949), TMEM18 (rs6548238), RBJ/POMC (rs713586), CADM2 (rs7640855), TRA2B/ETV5 (rs7647305), BDNF (rs925946), TFAP2B (rs987237), FTO (rs9941349), and ZNF608 (rs4836133). The summary of genotyped SNPs and allele frequencies is presented in **supplementary table S1**. All variants passed genotyping quality control criteria (call rate, >95%; Hardy-Weinberg equilibrium,  $P > .01$ ).

Combined obesity risk-allele scores, indicating genetic susceptibility to obesity, were calculated for each participant as the sum of risk alleles (0, 1 or 2 at each locus) associated with higher BMI across the 16 SNP loci. The score ranged from 0 to 32. To minimize dropout due to missing genotype data, infants with missing genotype data at 4 (25%) or fewer loci were imputed with the mean number of susceptibility alleles in their cohort for each locus. In sensitivity analyses, we also computed a weighted BMI-GRS, where each risk allele was multiplied by the ‘European only sex combined’ effect estimate for the BMI-increasing allele reported by Locke and colleagues (1).

### ***Child’s eating behavior***

Maternal perception of infant’s appetite was assessed, at ages 4, 8 and 12 months, by one question, translated as follow: “Usually, you would say that your baby: 1/ is always hungry or demanding to feed 2/ demands to feed the same as other babies of the same age 3/ needs to be stimulated to eat”, at ages 2 and 3 years, by one question: “Usually, you would say that your child: 1/ is always hungry or asking to eat 2/ has a normal appetite compared with other children of the same age 3/ is not often hungry”, and at 5 years by the Low Appetite scale from the Children’s Eating Difficulties Questionnaire (20). From 4 months to 4 years, a high appetite was defined as the category of children “always hungry or claiming to eat” and at 5 years as a Low Appetite score below the 10<sup>th</sup> percentile.

Infant diet was assessed by food records on three non-consecutive days (two weekdays and one weekend day), when the infant was not sick, at 4, 8 and 12 months. Data were checked and computerized a posteriori by a dietician. Nutrient intake was then calculated based on two food composition databases, one specific to ready-prepared baby foods from the French baby foods industry group (SFAE 2005) and one for common foods from the French Observatory of Food Nutritional Quality (CIQUAL 2006) (21). Total nutrient intake was calculated only among infants who were not breastfed during the three days of the dietary records, given that



the amount of breast milk intake could not be measured in breastfed infants. At each age, high energy intake was defined as the highest quintile of energy intake. Unfortunately, energy intake was not available in 2-5-year-old children, as diet was collected from a 24-item food frequency questionnaire from 2 years instead of 3-d dietary records.

### ***Child's growth***

At each clinical examination, child's weight and height were measured. In between, weight and height data were collected from self-administered questionnaires and from measurements noted in the child's health booklet by health professionals. Children had on average 22 weight measurements (interquartile range 16–26) from birth to 5–6 years. Individual growth curves were obtained for weight and height using the Jenss-Bayley growth curve model (22). This model allows us to predict weight, length/height and then calculate BMI at any given age (22). We used these model-based values of weight, length/height and BMI at 1, 2, 3, 4 and 5 y. At each age, the WHO growth standards were used for the calculation of weight-for-age, length/height-for-age and BMI-for-age z-scores. The WHO Anthro SAS macro (WHO Anthro, SAS Macro, Geneva: World Health Organization; <http://www.who.int/childgrowth/software/en/>) was used for this calculation.

### ***Sample selection***

Of the 2002 women who were recruited, 76 women were excluded because they left the study before or at the time of delivery, 24 because of miscarriages, intra-uterine death, or discontinuation of pregnancy for medical reasons, and 9 because they delivered outside the study hospitals. Data on birthweight were available for 1899 newborns. Among them, genotyping data were available for 1324 children. Children without any data on eating behavior from 4 months to 5 years (n=123) as well as those with missing data on age, sex, growth, breastfeeding duration or maternal smoking (n=59) were excluded from the analyses. Among the 1899 newborns with data on birth weight, the 1142 participants included in our

analyses were quite similar to those excluded concerning gender ( $p=0.9$ ) and maternal pre-gestational BMI ( $p=0.12$ ) but had older mothers (29.7 vs 29.1 years at delivery,  $p=0.007$ ) and were born with higher gestational age (39.4 vs. 39.0 weeks of amenorrhea,  $p<0.0001$ ) and higher birth weight (3324 g vs. 3210 g,  $p<0.0001$ ).

### ***Statistical analyses***

Student *t*-tests,  $\chi^2$ -tests and 1-way ANOVA were used to test differences between included and excluded populations.

Associations between the obesity risk-allele score (BMI-GRS) and child's eating behavior were tested by Poisson regression models with robust standard errors, adjusted for recruitment center, infant's gender, and age at eating behavior assessment. Additional analyses were conducted to adjust further for breastfeeding duration and maternal smoking during pregnancy.

Associations between infant/child eating behavior and predicted child's weight-for-age, length/height-for age or BMI-for-age WHO z-scores were tested by linear regression models adjusted for recruitment center, infant's sex, age at eating behavior assessment. Additional analyses were conducted to adjust further for breastfeeding duration, maternal smoking during pregnancy and parental height.

Associations between BMI-GRS and predicted child's weight, length/height or BMI z-scores were tested by linear regression models adjusted for recruitment center and infant's sex.

If an eating behaviour was associated with both the independent variable (BMI-GRS, path a, Supplementary Figure S1) and the dependent variable (WHO z-score, path b), we tested for the presence of mediation. Mediation is said to occur when a third variable lies on the causal pathway between an exposure (in this case, the BMI-GRS) and an outcome (in this case, WHO z-scores). To test for mediation, the linear regression of genetic risk score on each WHO z-score was adjusted for the considered eating behaviour. The presence of mediation

was established using the Sobel test (23, 24) and quantified by the mediation ratio  $((\beta - \beta') / \beta)$ , where  $\beta$  is the initial coefficient for BMI from the model  $WHO\ z\text{-score} \sim BMI\text{-GRS}$  (path c) and  $\beta'$  is the coefficient for WHO z-score after the model is additionally adjusted for the eating behaviour (path c'). As high score on appetitive traits were considered as binary variables, we used the method adapted by Jasti et al. to calculate the Sobel test (25). We conducted sensitivity analyses using a weighted genetic risk score for BMI (WtBMI-GRS) instead of the crude genetic risk score. To calculate this weighted score, at each locus, the number of BMI increasing variants was multiplied by the effect estimate for the BMI-increasing variant from Speliotes et al (2). All analyses were carried out using SAS V9.3 (SAS, Cary, NC).

## **Results**

Among the 1142 mothers-child pairs with both genotyping data and eating behavior assessment, mothers were aged on average 29.7 (SD=4.8) years, had a pre-pregnancy BMI of 23.1 (SD=4.3) kg/m<sup>2</sup> and breastfed for 3.3 (SD=3.7) months. Children carried on average 13.7 (SD=2.5) obesity risk alleles, with a range of 5 to 22 obesity risk alleles. At 5 years, 6.6 % of children were considered overweight or obese using IOTF definition (26). Children's BMI and appetitive traits are described in **Supplementary table S2**.

### ***BMI-GRS related to eating behaviours***

BMI-GRS was not significantly related to high energy intake at 4 mo and 8 mo but was positively related to energy intake at 1 year (in infants who were not breastfed at time of energy intake assessment) (**Figure 1**). Among all children, BMI-GRS was positively related to maternal perception of high appetite between 8 months and 5 years, although not significant in the first year or at 3 years. Further adjustment for breastfeeding duration and maternal smoking during pregnancy did not substantially modify the associations (data not

shown). In sensitivity analyses, when considering the weighted BMI-GRS, results were similar (data not shown).

### *Child's eating behavior and growth parameters*

Among children not breastfed during the 3-d dietary records, high energy intakes at 4 and 8 months were positively related to weight-for-age, length/ height-for-age and BMI-for-age z-scores (**Table 1**) from 1 to 5 years. High energy intake at 12 months was positively related to weight-for-age z-score from 3 to 5 years but was not related to length/ height-for-age or BMI-for-age z-scores.

High appetite before 1 year was not clearly related to subsequent WHO's z-scores. High appetite was not related to length/height-for-age z-score, whatever the age considered. High appetite at 1 year was positively related to weight-for-age or BMI-for-age z-scores up to 5 years but the strength of the association decreased with age and was no more significant for BMI-for-age at 5 years. High appetite at 2 or 3 years was positively related to child's subsequent BMI-for-age z-score and, to a lesser extent, weight-for-age z-score. These associations were mitigated by further adjustment for breastfeeding duration, maternal smoking during pregnancy and parental height, the associations but remained significant, except the association between high energy intake at 12 months and weight-for-age z-score from 3 to 5 years (supplementary table S3)

### *Mediation analysis*

The association between BMI-GRS and child's size increased with age and became significant from 3 years onwards for child's weight (**Figure 2A**) and 4 years onwards for child's BMI (**Figure 2B**). High appetite at 2 years was the only eating behavior related to both BMI-GRS and BMI, and therefore considered as a potential mediator in the association between BMI-GRS and BMI. These associations were attenuated after adjustment for high appetite at 2 years (Figure 2A and 2B). The Sobel test for mediation was significant but the

mediation was only partial with a mediation ratio decreasing with time from 47% for 2-y BMI z-score ( $p=0.03$ ) and 35% for 3-y BMI z-score ( $p=0.03$ ) to 28% for 4-y BMI z-score ( $p=0.05$ ) and 24% for 5-y BMI ( $p=0.05$ ). Similar results were found with the weighted BMI-GRS (data not shown).

## **Discussion**

In our study, the score of genetic predisposition to develop obesity was associated with a higher risk of high energy intake at 1 year, and high parental-perceived appetite at 2 and 5 years. The associations between the BMI-GRS and BMI-for-age or, to a lesser extent, weight-for-age z-scores were attenuated by further adjustment for high appetite at 2 years, whereas BMI-GRS was not related to length/height-for-age z-score up to 5 years. Finally, among infants not breastfed during the food records, energy intakes at 4 and 8 months were positively related to child's WHO z-scores until 5 years, but were not related to the BMI-GRS.

The association between genetic susceptibility to obesity and appetitive traits has been previously examined in childhood but not so early in infancy. Within 8-11 years olds twins of the TEDS study, Llewellyn et al. (14) showed that a BMI-GRS (based on 28 SNPs) was related to both BMI and satiety sensitivity, and that the latter mediated the association with BMI. However, in a study conducted by Steinsbeek et al. within Norwegian children aged from 4 to 6 years at inclusion (15), the BMI-GRS (based on 32 SNPs) was related to BMI from 4 to 8 years, to BMI change from 4 to 6 years and from 6 to 8 years, but was associated only with slowness in eating (negative association) at 6 years and not with other eating behaviours (i.e. food responsiveness, emotional overeating, enjoyment of food or satiety responsiveness). In that study, the association between the BMI-GRS and BMI was not mediated by any appetitive trait assessed at age 6 years.

238 If evidence of a mediating effect of appetitive traits on the association between the BMI-GRS  
239 and BMI remains scarce, many studies examined the influence of appetitive traits on BMI.  
240 The associations shown in EDEN are in agreement with previous studies in spite of the  
241 heterogeneity of ages and methods of investigation. In the GUSTO study (27), food  
242 responsiveness at 3 months was related to higher BMI up to 15 months, whereas appetitive  
243 traits assessed at 12 months were not related to BMI z-score in toddlers. Moreover, satiety  
244 responsiveness in early childhood was found related to lower subsequent BMI (28, 29).  
245 Similar results were found in the Gemini study (11), as all appetitive traits measured at 3  
246 months of age were related to later weight, with stronger association for general appetite and  
247 satiety responsiveness. In that study, the possibility of reverse causality was suggested by the  
248 observation that weight at 9 months was also related to appetitive traits in toddlers, with  
249 stronger association for food responsiveness, satiety responsiveness and general appetite.  
250 Despite the positive associations between infant appetite and subsequent weight gain and also  
251 between infant size and subsequent, the prospective influence of appetite on weight appeared  
252 to be stronger than the reverse.

253 In our study, energy intake in the first months of life was related to growth throughout early  
254 childhood. High energy intake could probably be considered as an indicator of rapid growth  
255 during this period and rapid growth in infancy is related to higher risk of overweight later in  
256 childhood in this and other cohorts (30). The influence of high energy intake on child's 5-y  
257 BMI is similar to the influence of high genetic predisposition to obesity (data not shown).  
258 However, notably energy intake in infancy was not related to the BMI-GRS, and might be  
259 more explained by prenatal or early postnatal exposures. We have to acknowledge that energy  
260 intake was assessed only among infants who were not breastfed during the three days of the  
261 dietary records, given that the amount of breast milk intake could not be measured in

breastfed infants. Moreover, from 2 years onwards, the dietary data collection was unfortunately not designed to derive energy intake.

The main strength of this study is that eating behaviour and BMI were assessed repetitively and prospectively from very early in infancy (as early as birth for BMI and four months for eating behaviour), which permitted us to describe in detail the longitudinal patterns of associations between the BMI-GRS, eating behaviour and BMI. However, appetite was assessed only by one item, whereas other assessments, maybe based on more complex scales, would have allowed us to check for consistency for this trait. Moreover, we have to acknowledge that eating behaviour up to 3 years was not assessed by a validated tool up to 3 years as such tools were not developed when the EDEN study was designed. A validated tool became available and was introduced only at the 5-y follow-up. Concerning the item on appetite, a similar item was used by previous studies (10, 11) and associations with this item were similar to those with other appetitive traits, if not stronger. Moreover, in the EDEN study, we found similar associations at 5 years, when appetite was assessed by a validated scale (20). As appetitive traits were reported by parental questionnaires and not assessed by a laboratory tool, this measurement could be influenced by social desirability. Finally, because of the originality and the uniqueness of the collected data, our results yet rely on data from only one cohort. Further studies conducted in other birth cohorts would be necessary to confirm and demonstrate the generalizability of these findings. Ideally, these studies would be designed with repeated use of validated tools throughout early childhood to assess eating behaviour or energy intake and conducted in populations with higher prevalence of overweight. As previous studies have shown that these BMI variants could to be relevant also to other non-Caucasian populations (31, 32), it would be interesting to conduct similar analyses in non-Caucasian populations.

287

288 **Conclusion**

289 The study provides further support from a prospective study for a partial mediation by  
290 appetitive traits on the relationship between genetic susceptibility and BMI. In particular, in  
291 the EDEN birth cohort, genetic susceptibility to childhood and adult obesity seems to play a  
292 very precocious role by influencing appetitive traits in infants and preschool children and  
293 subsequently promote faster growth. Despite the more typically recognised concerns  
294 regarding infant feeding difficulties, many parents might benefit from counselling and support  
295 to manage high appetitive traits in their infants and young children to prevent later obesity  
296 risk.

297



## 298 **Sources of support**

299 The EDEN study is supported by: Fondation pour la Recherche Médicale (FRM), French  
 300 Ministry of Research: Federative Research Institutes and Cohort Program, INSERM Human  
 301 Nutrition National Research Program, and Diabetes National Research Program (through a  
 302 collaboration with the French Association of Diabetic Patients (AFD)), French Ministry of  
 303 Health, French Agency for Environment Security (AFSSET), French National Institute for  
 304 Population Health Surveillance (InVS), Paris–Sud University, French National Institute for  
 305 Health Education (INPES), Nestlé, Mutuelle Générale de l'Éducation Nationale (MGEN),  
 306 French speaking association for the study of diabetes and metabolism (ALFEDIAM),  
 307 National Agency for Research (ANR non thematic program), National Institute for Research  
 308 in Public health (IRESP: TGIR 2008 cohort in health program). The genotyping was funded  
 309 by a Collaborative Research Grant from the European Society for Paediatric Endocrinology.  
 310 KKO is supported by the Medical Research Council (Unit program: MC\_UU\_12015/2).

## 311 **Acknowledgment section**

312 None of the authors have any financial relationships or conflict of interest to disclose.

313

314 BLG and BH designed the research and wrote the manuscript. BLG, YAK, JB, AF and SC  
 315 analysed the data. BH and MAC oversaw the EDEN study. BH, KC, VP, KKO and MAC  
 316 were responsible for data collection and genotyping in EDEN. All authors reviewed drafts,  
 317 provided critical feedback, approved the final manuscript and were responsible for the final  
 318 content of the paper. BH had full access to all the data in the study and takes responsibility for  
 319 the integrity of the data and the accuracy of the data analysis.


320

## **References**

1. Locke AE, Kahali B, Berndt SI, *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature* 2015; **518**, 197-206.
2. Speliotes EK, Willer CJ, Berndt SI, *et al.* Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet* 2010; **42**, 937-48.
3. Belsky DW, Moffitt TE, Houts R, *et al.* Polygenic risk, rapid childhood growth, and the development of obesity: evidence from a 4-decade longitudinal study. *Arch Pediatr Adolesc Med* 2012; **166**, 515-21.
4. Elks CE, Heude B, de Zegher F, *et al.* Associations between genetic obesity susceptibility and early postnatal fat and lean mass: an individual participant meta-analysis. *JAMA Pediatr* 2014; **168**, 1122-30.
5. Elks CE, Loos RJ, Hardy R, *et al.* Adult obesity susceptibility variants are associated with greater childhood weight gain and a faster tempo of growth: the 1946 British Birth Cohort Study. *Am J Clin Nutr* 2012; **95**, 1150-6.
6. Elks CE, Loos RJ, Sharp SJ, *et al.* Genetic markers of adult obesity risk are associated with greater early infancy weight gain and growth. *PLoS Med* 2010; **7**, e1000284.
7. Day FR and Loos RJ. Developments in obesity genetics in the era of genome-wide association studies. *J Nutrigenet Nutrigenomics* 2011; **4**, 222-38.
8. Elder SJ, Neale MC, Fuss PJ, *et al.* Genetic and Environmental Influences on Eating Behavior - A Study of Twin Pairs Reared Apart or Reared Together. *Open Nutr J* 2012; **6**, 59-70.
9. Llewellyn CH, van Jaarsveld CH, Boniface D, Carnell S and Wardle J. Eating rate is a heritable phenotype related to weight in children. *Am J Clin Nutr* 2008; **88**, 1560-6.
10. Llewellyn CH, van Jaarsveld CH, Plomin R, Fisher A and Wardle J. Inherited behavioral susceptibility to adiposity in infancy: a multivariate genetic analysis of appetite and weight in the Gemini birth cohort. *Am J Clin Nutr* 2012; **95**, 633-9.
11. van Jaarsveld CH, Llewellyn CH, Johnson L and Wardle J. Prospective associations between appetitive traits and weight gain in infancy. *Am J Clin Nutr* 2011; **94**, 1562-7.
12. Emond JA, Tovar A, Li Z, Lansigan RK and Gilbert-Diamond D. FTO genotype and weight status among preadolescents: Assessing the mediating effects of obesogenic appetitive traits. *Appetite* 2017; **117**, 321-329.
13. Wardle J, Carnell S, Haworth CM, Farooqi IS, O'Rahilly S and Plomin R. Obesity associated genetic variation in FTO is associated with diminished satiety. *J Clin Endocrinol Metab* 2008; **93**, 3640-3.
14. Llewellyn CH, Trzaskowski M, van Jaarsveld CH, Plomin R and Wardle J. Satiety mechanisms in genetic risk of obesity. *JAMA Pediatr* 2014; **168**, 338-44.
15. Steinsbekk S, Belsky D, Guzey IC, Wardle J and Wichstrom L. Polygenic Risk, Appetite Traits, and Weight Gain in Middle Childhood: A Longitudinal Study. *JAMA Pediatr* 2016; **170**, e154472.
16. Heude B, Forhan A, Slama R, *et al.* Cohort Profile: The EDEN mother-child cohort on the prenatal and early postnatal determinants of child health and development. *Int J Epidemiol* 2016; **45**, 353-63.
17. Regnault N, Botton J, Forhan A, *et al.* Determinants of early ponderal and statural growth in full-term infants in the EDEN mother-child cohort study. *Am J Clin Nutr* 2010; **92**, 594-602.


18. Betoko A, Charles MA, Hankard R, *et al.* Infant feeding patterns over the first year of life: influence of family characteristics. *Eur J Clin Nutr* 2013; **67**, 631-637.
19. Li S, Zhao JH, Luan J, *et al.* Cumulative effects and predictive value of common obesity-susceptibility variants identified by genome-wide association studies. *Am J Clin Nutr* 2010; **91**, 184-90.
20. Rigal N, Chabanet C, Issanchou S and Monnery-Patris S. Links between maternal feeding practices and children's eating difficulties. Validation of French tools. *Appetite* 2012; **58**, 629-37.
21. ANSES. French food composition table [internet] 2015, France: French Agency for Food, Environmental and Occupational Health & Safety (ANSES). **2015**.
22. Botton J, Scherdel P, Regnault N, Heude B, Charles MA and Eden Mother-Child Cohort Study Group. Postnatal weight and height growth modeling and prediction of body mass index as a function of time for the study of growth determinants. *Ann Nutr Metab* 2014; **65**, 156-66.
23. Mackinnon DP, Warsi G and Dwyer JH. A Simulation Study of Mediated Effect Measures. *Multivariate Behav Res* 1995; **30**, 41.
24. Sobel ME. Asymptotic intervals for indirect effect in structural equations models. In *Sociological methodology*, ed. Leinhardt S, 1982, pp 290-312 San Francisco: Jossey-Bass.
25. Jasti S, Dudley WN and Goldwater E. SAS macros for testing statistical mediation in data with binary mediators or outcomes. *Nurs Res* 2008; **57**, 118-22.
26. Cole TJ, Bellizzi MC, Flegal KM and Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000; **320**, 1240-3.
27. Quah PL, Chan YH, Aris IM, *et al.* Prospective associations of appetitive traits at 3 and 12 months of age with body mass index and weight gain in the first 2 years of life. *BMC Pediatr* 2015; **15**, 153.
28. de Barse LM, Tiemeier H, Leermakers ET, *et al.* Longitudinal association between preschool fussy eating and body composition at 6 years of age: The Generation R Study. *Int J Behav Nutr Phys Act* 2015; **12**, 153.
29. Mallan KM, Nambiar S, Magarey AM and Daniels LA. Satiety responsiveness in toddlerhood predicts energy intake and weight status at four years of age. *Appetite* 2014; **74**, 79-85.
30. Botton J, Heude B, Maccario J, Ducimetière P and Charles MA. Postnatal weight and height growth velocities at different ages between birth and 5 years and body composition in adolescent boys and girls. *Am J Clin Nutr* 2008; **87**, 1760-8.
31. Dorajoo R, Ong RT, Sim X, *et al.* The contribution of recently identified adult BMI risk loci to paediatric obesity in a Singaporean Chinese childhood dataset. *Pediatr Obes* 2017; **12**, e46-e50.
32. Zandoná MR, Sangalli CN, Campagnolo PdB, Vitolo MR, Almeida S and Mattevi VS. Validation of obesity susceptibility loci identified by genome-wide association studies in early childhood in South Brazilian children. *Pediatric Obesity* 2017; **12**, 85-92.

**Figure 1. Association between obesity risk-allele score and child's eating behaviour, up to 5 years**



*Figure legend.* Results are RR [95% CI], adjusted for recruitment's center, child's age and sex (n=1081 at 4mo, 1072 at 8 mo, 1024 at 1 year, 940 at 2 years, 892 at 3 years and 790 at 5 years). Energy intake was assessed only on infants who were not breastfed during the 3-d dietary records. (679 at 4 mo, 861 at 8 mo and 822 at 12 mo)

**Figure 2. Association between the obesity risk-allele score and child's WHO z-scores from 2 to 5 years**



*Figure legend.* ◆ Results are effect size [95% CI], adjusted for recruitment center and infant's sex (n=1142). ■ Results are effect size [95% CI], adjusted for recruitment center, infant's sex, child's age and high appetite at 2 years (n=940).

## SUPPORTING INFORMATION

**Table S1. BMI and eating behaviour traits of children at different assessment steps**

SNP	Locus	Effect allele	Other allele	SNP reported in GIANT	Evidence for association with childhood BMI	LD with GIANT SNP (r <sup>2</sup> )	EAF in EDEN
rs10146997	NRXN3	G	A	rs10150332	Reported in Speliotes et al.	1.0	0.18
rs13107325	SLC39A8	T	C	rs13107325	Reported in Elks et al.	-	0.08
rs1514175	TNNI3K	A	G	rs1514175	Reported in Speliotes et al.	-	0.38
rs1555543	PTBP2	C	A	rs1555543	Reported in Speliotes et al.	-	0.55
rs17782313	MC4R	C	T	rs571312	Reported in den Hoed et al.	0.96	0.21
rs2112347	FLJ35779	T	G	rs2112347	Reported in Elks et al.	-	0.61
rs2568958	NEGR1	A	G	rs2815752	Reported in den Hoed et al.	0.96	0.57
rs4836133	ZNF608	A	C/G	rs4836133	Reported in Elks et al.	-	0.40
rs4929949	RPL27A	C	T	rs4929949	Reported in Speliotes et al.	-	0.45
rs6548238	TMEM18	C	T	rs2867125	Reported in den Hoed et al.	1.0	0.81
rs713586	RBJ/POMC	C	T	rs713586	Reported in Speliotes et al.	-	0.44
rs7640855	CADM2	A	G	rs13078807	Reported in Speliotes et al.	1.0	0.17
rs7647305	TRA2B	C	T	rs9816226	Reported in den Hoed et al.	0.72	0.71
rs925946	BDNF	T	G	rs10767664	Reported in den Hoed et al.	1.0	0.25
rs987237	TFAP2B	G	A	rs987237	Reported in Elks et al.	-	0.15
rs9941349	FTO	T	C	rs1558902	Reported in den Hoed et al.	0.84	0.38

Effect allele refers to the allele associated with higher BMI in the Genetic Investigation of Anthropometric Traits (GIANT) consortium (Speliotes et al., Ref. 2).

EAF: Effect allele frequency.

LD: Linkage Disequilibrium; calculated based on 1000 Genomes pilot 1 data using SNAP (<https://www.broadinstitute.org/mpg/snap/>).

**References**

Speliotes EK, Willer CJ, Berndt SI, et al. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature Genetics*. 2010;42(11):937-948.

den Hoed M, Ekelund U, Brage S, et al. Genetic susceptibility to obesity and related traits in childhood and adolescence; influence of loci identified by genome-wide association studies. *Diabetes*. 2010;59(11):2980-2988.

Elks CE, Heude B, de Zegher F, et al. Associations between genetic obesity susceptibility and early postnatal fat and lean mass: an individual participant meta-analysis. *JAMA Pediatr* 2014; 168, 1122-30.

**Table S2. BMI and eating behaviour traits of children at different assessment steps**

	4 months	8 months	1 year	2 years	3 years	5 years
N	1081	1072	1024	940	892	790
Age (months)	4.3 (1.0)	8.3 (0.8)	12.1 (0.7)	24.4 (1.3)	37.9 (0.9)	67.7 (1.9)
Predicted BMI (kg/m <sup>2</sup> )	16.9 (1.3)	17.1 (1.3)	16.8 (1.2)	16.2 (1.2)	15.9 (1.2)	15.4 (1.3)
High appetite (always hungry) <sup>a</sup>	3.7% (40)	1.9% (20)	3.1% (32)	5.1% (48)	7.2% (64)	9.0% (71)
Energy intake (kcal)						
among children not receiving breastmilk during the 3-d record	n=680 589 (104)	n=861 716 (135)	n=822 809 (135)			

---

Values are means (sd) or % (n).

<sup>a</sup> at 5 years, high appetite was defined as a Low Appetite Score below the 10<sup>th</sup> percentile.



**Table S3. Association between eating behaviour and child's WHO z-scores after further adjustment on parental height, maternal smoking during pregnancy and breastfeeding duration**


	N	Age at WHO z-score assessment				
		1 year	2 years	3 years	4 years	5 years
<b>Outcome: Weight-for-age z-score</b>						
High energy intake (highest quintile)						
4 months	629	<b>0.39 [0.23 ; 0.55]</b>	<b>0.31 [0.15 ; 0.46]</b>	<b>0.25 [0.09 ; 0.41]</b>	<b>0.21 [0.05 ; 0.36]</b>	<b>0.18 [0.02 ; 0.33]</b>
8 months	791	<b>0.33 [0.19 ; 0.47]</b>	<b>0.29 [0.15 ; 0.43]</b>	<b>0.24 [0.10 ; 0.38]</b>	<b>0.19 [0.05 ; 0.33]</b>	<b>0.15 [0.02 ; 0.29]</b>
12 months	767	0.04 [-0.10 ; 0.19]	0.07 [-0.07 ; 0.22]	0.09 [-0.06 ; 0.23]	0.09 [-0.05 ; 0.23]	0.09 [-0.05 ; 0.23]
High appetite (always hungry)						
4 months	993	<b>0.32 [0.04 ; 0.59]</b>	0.21 [-0.07 ; 0.48]	0.14 [-0.14 ; 0.42]	0.09 [-0.18 ; 0.37]	0.06 [-0.21 ; 0.34]
8 months	987	0.28 [-0.10 ; 0.66]	0.23 [-0.14 ; 0.61]	0.17 [-0.21 ; 0.55]	0.11 [-0.26 ; 0.49]	0.08 [-0.3 ; 0.45]
1 year	945	0.21 [-0.10 ; 0.51]	0.23 [-0.08 ; 0.53]	0.20 [-0.11 ; 0.51]	0.16 [-0.14 ; 0.47]	0.14 [-0.17 ; 0.44]
2 years	868		<b>0.32 [0.08 ; 0.56]</b>	<b>0.28 [0.04 ; 0.53]</b>	<b>0.24 [0.00 ; 0.48]</b>	0.20 [-0.04 ; 0.44]
3 years	826			<b>0.28 [0.06 ; 0.50]</b>	<b>0.24 [0.02 ; 0.46]</b>	0.21 [-0.01 ; 0.43]
5 years	738					<b>0.41 [0.20 ; 0.61]</b>
<b>Outcome : Length/Height-for-age z-score</b>						
High energy intake (highest quintile)						
4 months	629	<b>0.25 [0.08 ; 0.42]</b>	0.13 [-0.04 ; 0.29]	0.11 [-0.06 ; 0.28]	0.10 [-0.06 ; 0.26]	0.11 [-0.05 ; 0.27]
8 months	791	<b>0.21 [0.06 ; 0.35]</b>	<b>0.20 [0.05 ; 0.34]</b>	<b>0.18 [0.03 ; 0.33]</b>	<b>0.13 [0.00 ; 0.27]</b>	0.11 [-0.03 ; 0.25]
12 months	767	0.11 [-0.05 ; 0.26]	0.09 [-0.05 ; 0.24]	0.10 [-0.06 ; 0.25]	0.09 [-0.06 ; 0.23]	0.08 [-0.06 ; 0.23]
High appetite (always hungry)						
4 months	993	0.07 [-0.23 ; 0.37]	0.03 [-0.26 ; 0.32]	0.04 [-0.26 ; 0.34]	0.05 [-0.22 ; 0.33]	0.07 [-0.21 ; 0.35]
8 months	987	0.15 [-0.25 ; 0.55]	0.11 [-0.28 ; 0.50]	0.08 [-0.33 ; 0.48]	0.05 [-0.33 ; 0.42]	0.02 [-0.36 ; 0.40]
1 year	945	-0.27 [-0.59 ; 0.06]	-0.22 [-0.53 ; 0.09]	-0.18 [-0.50 ; 0.15]	-0.12 [-0.42 ; 0.18]	-0.07 [-0.38 ; 0.23]
2 years	868		-0.20 [-0.45 ; 0.05]	-0.20 [-0.46 ; 0.07]	-0.17 [-0.41 ; 0.07]	-0.16 [-0.41 ; 0.08]
3 years	826			0.11 [-0.12 ; 0.34]	0.09 [-0.13 ; 0.30]	0.07 [-0.15 ; 0.29]

5 years	738					<b>0.30 [0.09 ; 0.50]</b>
<b>Outcome: BMI-for-age z-score</b>						
High energy intake (highest quintile)						
4 months	629	<b>0.35 [0.18 ; 0.52]</b>	<b>0.33 [0.16 ; 0.51]</b>	<b>0.28 [0.11 ; 0.45]</b>	<b>0.21 [0.04 ; 0.38]</b>	0.16 [-0.01 ; 0.33]
8 months	791	<b>0.29 [0.14 ; 0.44]</b>	<b>0.25 [0.10 ; 0.40]</b>	<b>0.21 [0.05 ; 0.36]</b>	<b>0.16 [0.02 ; 0.31]</b>	0.14 [-0.01 ; 0.28]
12 months	767	-0.02 [-0.18 ; 0.13]	0.03 [-0.13 ; 0.18]	0.05 [-0.11 ; 0.20]	0.06 [-0.10 ; 0.21]	0.06 [-0.09 ; 0.21]
High appetite (always hungry)						
4 months	993	<b>0.37 [0.08 ; 0.66]</b>	0.27 [-0.03 ; 0.57]	0.17 [-0.13 ; 0.47]	0.09 [-0.20 ; 0.38]	0.03 [-0.26 ; 0.33]
8 months	987	0.27 [-0.13 ; 0.67]	0.25 [-0.16 ; 0.66]	0.19 [-0.22 ; 0.60]	0.14 [-0.26 ; 0.54]	0.11 [-0.29 ; 0.51]
1 year	945	<b>0.49 [0.17 ; 0.81]</b>	<b>0.52 [0.20 ; 0.85]</b>	<b>0.45 [0.12 ; 0.78]</b>	<b>0.35 [0.02 ; 0.67]</b>	0.27 [-0.05 ; 0.59]
2 years	868		<b>0.64 [0.38 ; 0.90]</b>	<b>0.60 [0.34 ; 0.86]</b>	<b>0.52 [0.26 ; 0.78]</b>	<b>0.46 [0.20 ; 0.72]</b>
3 years	826			<b>0.32 [0.08 ; 0.56]</b>	<b>0.29 [0.05 ; 0.52]</b>	<b>0.27 [0.03 ; 0.50]</b>
5 years	738					<b>0.35 [0.13 ; 0.57]</b>

Results are effect size [95% CI] of high level of the considered appetitive trait on WHO z-score, adjusted for recruitment centre, infant's sex, age at eating behaviour assessment, parental height, maternal smoking during pregnancy and breastfeeding duration. Significant results are highlighted in bold.

<sup>a</sup> at 5 years, high appetite was defined as a Low Appetite Score below the 10<sup>th</sup> percentile.

**Figure S1. Figure depicting the mediation analyses.**



**Path a** shows the association between the BMI genetic risk score and eating behaviour

**Path b** shows the association between eating behaviour and WHO's z-score

**Path c** shows the relationship between the BMI genetic risk score and WHO's z-score

**Path c'** shows the relationship between the BMI genetic risk score and WHO's z-score, adjusted for the considered eating behaviour