



HAL
open science

Identification of a common transcriptional signature for regulatory B cells in Humans and Mice

Florian Dubois, Sophie Limou, Mélanie Chesneau, Sophie Brouard, Richard Danger

► **To cite this version:**

Florian Dubois, Sophie Limou, Mélanie Chesneau, Sophie Brouard, Richard Danger. Identification of a common transcriptional signature for regulatory B cells in Humans and Mice. FOCIS 2019, Jun 2019, Boston, United States. inserm-02160256

HAL Id: inserm-02160256

<https://inserm.hal.science/inserm-02160256v1>

Submitted on 19 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Identification of a common transcriptional signature for regulatory B cells in Humans and Mice

Florian Dubois^{1,2}, Sophie Limou^{1,2,3}, Mélanie Chesneau^{1,2}, Sophie Brouard^{1,2,4,*} and Richard Danger^{1,2,*}

¹CRTI UMR1064, INSERM, Université de Nantes, France; ²ITUN, CHU Nantes, Nantes, France; ³Ecole Centrale de Nantes, Nantes, France; ⁴Centre d'Investigation Clinique en Biothérapie, CRB, Labex IGO, Nantes, France, *Last authors equally contributed and are listed by alphabetical order.

Background


Regulatory B cells (Bregs) have been described in mice and humans for their ability to regulate inflammation through a variety of mechanisms in different pathological situations. Up to date, no consensual and common Breg phenotype has been described, and whether there is a Breg lineage commitment or if they acquire their function under certain environmental conditions remains unknown. To address these points, we performed a sample size weighted meta-analysis of publicly available transcriptomic data from 4 different Bregs studies in humans and 6 Bregs studies in mice.

Materials and Methods

In human we retrieved four datasets comparing Bregs and non-Bregs: Van de Ven et al., JACI 2013, Lin et al., Immunol 2014, Bigot et al., AJT 2016 and Chesneau et al., submitted. And 6 in mice: Shen et al., Nature 2014, Khan et al., Eur. J. Immunol 2015, Braza et al., Allergy 2015, Lino et al., Immunity 2018 and Ray et al., Nat com 2018. For each dataset, raw data were processed following a homogenous method and differentially expressed genes between Bregs and non-Bregs were identified using the limma R package with adjusted p-value <5% and fold change (FC) > 2 were considered as significantly differentially expressed. Finally, the sample size weighted meta-analysis (Stouffer's Z-score method) was performed using the METAL software (Willer et al., 2010).


Results

Core gene signature discriminating Bregs from non-Bregs in human



- 165 common genes were identified to be differentially expressed between Bregs and non-Bregs in the meta-analysis of the four studies.
- This human gene signature well discriminates Bregs and non-Bregs.


Core gene signature discriminating Bregs from non-Bregs in mice




- 93 common genes were identified with a great discrimination between Bregs and non-Bregs in mice datasets.

Predicted protein-protein interactions and potential enrichment of transcription factor targets in human Breg signature

String interaction among the Up-regulated differentially expressed genes




Enrichment of Transcription factors binding site




- The human Breg signature is composed of genes known as involved in immune regulation and cell cycle and related to 4 transcription factors regulating genes implicated in B cell biology.

Predicted protein-protein interactions and potential enrichment of transcription factor targets in mice Breg signature

String interaction among the Up-regulated differentially expressed genes



Enrichment of Transcription factors binding site



- Clear gene signature of mice Bregs with mostly up-regulated genes involved in regulation of immune response and an enrichment of transcription factor binding site in genes with an apoptotic function.

Conclusion

Our results highlight human Bregs with a proliferating state and provide new insights on possible mechanisms by which Bregs tightly regulate their homeostasis. In mice, Bregs exhibit a gene expression with function associated in the regulation of immune response. Moreover the absence of common signature between human and mice Bregs and the lack of a common transcription factor support the hypothesis of B cell acquiring their regulatory function under certain environmental conditions. Understanding what are the signal for B cells to become a Breg will help their study and understand their role in cancer, autoimmunity and in tolerance mechanisms.