



HAL
open science

Dopamine and addiction: what have we learned from 40 years of research.

Marcello Solinas, Pauline Belujon, Pierre Olivier Fernagut, Mohamed Jaber,
Nathalie Thiriet

► To cite this version:

Marcello Solinas, Pauline Belujon, Pierre Olivier Fernagut, Mohamed Jaber, Nathalie Thiriet. Dopamine and addiction: what have we learned from 40 years of research.. *Journal of Neural Transmission*, 2019, 126 (4), pp.481-516. 10.1007/s00702-018-1957-2 . inserm-02163255

HAL Id: inserm-02163255

<https://inserm.hal.science/inserm-02163255>

Submitted on 24 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Dopamine and Addiction: what have we learned from 40 years of research

Marcello Solinas¹, Pauline Belujon¹, Pierre Olivier Fernagut¹, Mohamed Jaber^{1,2},
Nathalie Thiriet¹

¹*Université de Poitiers, INSERM, Laboratoire de Neurosciences Expérimentales et Cliniques, Poitiers, France.*

²*CHU de Poitiers, Poitiers, France.*

Corresponding Author:

Marcello Solinas

e-mail: marcello.solinas@univ-poitiers.fr

Tel: +33-(0)- 549-366343

ABSTRACT

Among the neurotransmitters involved in addiction, dopamine (DA) is clearly the best known. The critical role of DA in addiction is supported by converging evidence that has been accumulated in the last forty years. In the present review, firstly we describe the dopaminergic system in terms of connectivity, functioning and involvement in reward processes. Secondly, we describe the functional, structural, and molecular changes induced by drugs within the DA system in terms of neuronal activity, synaptic plasticity and transcriptional and molecular adaptations. Thirdly, we describe how genetic mouse models have helped characterizing the role of DA in addiction. Fourthly, we describe the involvement of the DA system in the vulnerability to addiction and the interesting case of addiction DA replacement therapy in Parkinson's disease. Finally, we describe how the DA system has been targeted to treat patients suffering from addiction and the result obtained in clinical settings and we discuss how these different lines of evidence have been instrumental in shaping our understanding of the pathophysiology of drug addiction.

Keywords: substance abuse, reward, neuroplasticity, treatment, substantia nigra, VTA, striatum

Introduction	4
Addiction and the addiction cycle	4
The DA system.....	5
DA neurons, connectivity and activity	5
DA Receptor Subtypes and distribution.....	10
DAT expression and function	12
Dopaminoceptive neurons and their glutamatergic afferents	13
DA and reinforcement/reward processes	14
Drugs, DA and reinforcement.....	17
DA, drug withdrawal and negative affect.....	18
DA, craving and relapse.....	20
Addiction and DA brain imaging.....	21
Electrophysiological effects of drugs on the DA system	23
Effects of acute administration of drugs on the activity of DA neurons.....	23
Effects of chronic administration of drugs on the neuronal activity of DA neurons	27
Effects of chronic administration of drugs on the activity of NAc and DSt neurons.....	28
Addiction, DA and Synaptic plasticity	29
Addiction and DA structural plasticity	34
Drugs and transcriptional plasticity in DA regions.....	36
Drugs and molecular adaptations within the DA system	38
Genetically modified animals, DA and Addiction.....	40
Genetic manipulations of the DAT and responses to drugs of abuse	40
Genetic manipulation of DA receptors.....	46
Vulnerability to addiction	49
Environment, DA and vulnerability to addiction	50
Addiction to DA replacement therapy medications in Parkinson's disease.....	52
DA and the treatment of addiction	56
Conclusions.....	60

Introduction

The involvement of dopamine (DA) in addiction processes has been proposed more than 40 years ago (Dackis and Gold 1985; Yokel and Wise 1975). Indeed, several converging anatomical, neurochemical, functional and behavioral evidence suggested early on that DA plays an important role in addiction, especially to psychostimulants (Dackis and Gold 1985; Wise 1984; Wise and Bozarth 1987). Subsequently, an impressive amount of research has confirmed that the DA system is fundamental in the development, maintenance, withdrawal and relapse phases of addiction (Koob and Volkow 2010; Robinson and Berridge 2008; Di Chiara and Bassareo 2007; Hyman et al. 2006; Hyman 2005; Everitt and Robbins 2005; Schultz 2011). In this article, we summarize several lines of evidences that demonstrate the role of DA in addiction and we briefly discuss how this knowledge has translated into clinical therapy and the lessons that could be learned from this field of research.

Addiction and the addiction cycle

Drug addiction is a chronic, relapsing psychiatric disorder that represents a major public health problem that impacts society on multiple levels, including health care, loss of productivity, police and crime/accident costs. The diagnostic and statistical manual of the American Psychiatry Association describes eleven criteria/symptoms that can be used to diagnose addiction in humans (DSM-5 2013). These criteria itemize the compulsive seeking and taking of the drug despite negative consequences, craving and a negative state that appears when the drug is withdrawn which are considered the key elements of addiction.

Addiction is characterized by phases of active, excessive consumption of the drug, phases of more controlled use, phases of abstinence and episodes of relapse (Koob and Volkow 2010; Volkow et al. 2004). Indeed, one of the criteria of addiction is that

individuals make unsuccessful attempts to reduce or quit drug consumption. Therefore, the pathology of addiction is often described as a cycle with three stages that are associated with different behavioral and neurobiological mechanisms: binge/intoxication, withdrawal/negative affect and preoccupation/anticipation (Koob and Volkow 2010, 2016; Koob and Le Moal 2001). The binge/intoxication stage is associated with positive reinforcement and stimulation of the reward system in which DA plays a major role during initial use of drugs and, as drug use becomes more habitual with stimulation of dorsal striatal regions (Koob and Volkow 2010, 2016). Conversely, the withdrawal stage is associated with negative affect, hypoactivity of the DA system and hyperactivation of the stress system, which drives negative reinforcement, i.e. drugs are taken not for their positive effects but to avoid negative emotional states. Finally, the preoccupation/anticipation stage is associated with craving for the drug, activation of cortical areas and release of DA in areas associated with emotions and memory such as the amygdala and the hippocampus (Koob and Volkow 2010, 2016).

The DA system

DA neurons, connectivity and activity

DA neurons are located in three main groups in the midbrain: A8 in the retrorubral field, A9 in the substantia nigra pars compacta (SNc) and A10 in the ventral tegmental area (VTA). DA neurons from the VTA send projections to the nucleus accumbens (NAc) (meso-striatal system) and also provide inputs to other forebrain regions such as the prefrontal cortex (meso-cortical system) (Wise 2009; Haber and Knutson 2010; Haber et al. 1990). The SNc (nigrostriatal system) and the retrorubral area send projections to the dorsal striatum (DSt) (Wise 2009; Haber and Knutson

2010; Haber et al. 1990). Historically, the nigro-striatal pathway has been involved principally in the control of movement whereas the meso-cortico-striatal pathway has been involved in reward-learning and motivation. However, compelling evidence suggests that this clear-cut distinction is an oversimplification and that the DA system functions as a medial-to-lateral gradient of overlapping populations of DA neurons (Wise 2009).

DA neurons display multiple activity states: they can either be spontaneously active (firing) or silent (non-firing) (Figure 1, inset). The proportion of silent DA neurons appears to be maintained at a constant, hyperpolarized, inactive state by the ventral pallidum (VP) (Floresco et al. 2003). When DA neurons are spontaneously firing, they can fire with an irregular pattern (tonic activity) or displaying bursts of action potentials (phasic activity). Importantly, only spontaneously active DA neurons are able to fire in bursts (Lodge and Grace 2006) (Figure 1, inset). Thus, the proportion of neurons that fires spontaneously (“population activity”) supplies the stable baseline level of extrasynaptic DA in postsynaptic structures and has been termed the “tonic” DA state (Grace 2000). Changes in firing from irregular tonic firing to burst phasic firing has been shown to be involved in reinforcement learning. Indeed, DA neurons display phasic burst first during the primary presentation with an unpredicted reward (Schultz 1998; Zweifel et al. 2011), but then, after conditioning with repeated stimulus-reward pairing, burst firing switches to the reward-predicting stimulus alone (positive prediction error) (Lammel et al. 2011; Zweifel et al. 2011; Schultz 1997). When the reward is not produced anymore after the stimulus, a pause in DA neurons firing occurs (negative prediction error). Therefore, firing of DA neurons appears to encode the signaling of reward prediction error, i.e. the difference between reward prediction and actual received reward (see following sections for more details).

Spontaneous activity of DA neurons is generated by intrinsic pace-making membrane properties of DA neurons themselves as demonstrated by the fact that tonic spiking activity is preserved in brain slice preparation, where the synaptic inputs have been severed (Deister et al. 2009; Shepard and Bunney 1988). In contrast, burst spiking is lost in such preparations, and is dependent on glutamate receptors activation, since local application of glutamate receptor antagonists disrupts this pattern, suggesting the necessity of synaptic inputs for the generation of this pattern of activity (Grace and Bunney 1984; Smith and Grace 1992). Indeed, burst firing is triggered by the transient interaction between ionotropic glutamate receptors and voltage-gated ionic channels (Grace and Bunney 1984; Hyland et al. 2002; Smith and Grace 1992).

Changes in DA activity state in the SNc and the VTA have consequences on the levels of DA in the NAc and the DSt respectively. Tonic DA activity is responsible for the basal levels (in the range of nM; (Sharp et al. 1986)) of extracellular DA in the striatum. Although this level of extracellular DA is relatively low, it is sufficient to induce a partial activation of DA D2 auto-receptors (Farnebo and Hamberger 1971), which results in suppression of most of the spike-dependent DA release. Conversely, phasic DA release corresponds to the rapid release of a significant amount (in the range of μM) of DA in the synaptic cleft (May et al. 1988) resulting from an activation of DA neurons in response to behaviorally salient stimulus, and leading to the activation of receptors on postsynaptic neurons (Grace 2000). DA is then rapidly removed from the synapse, by the DA transporter (DAT), and the signal is terminated. This massive but transient DA release can be measured in behaving animals by continuous amperometry (Benoit-Marand et al. 2011; Benoit-Marand et al. 2001; Benoit-Marand et al. 2000a) or fast scan cyclic voltammetry (Phillips et al.

2003a; Robinson et al. 2003) but not by techniques such as microdialysis or classical voltammetry (May et al. 1988).

VTA and SNc DA neurons receive different inhibitory and excitatory inputs that modulate their firing (Figure 1). The control of tonic firing of VTA neurons involves the bed nucleus of the stria terminalis (Georges and Aston-Jones 2001) and the ventral pallidum (Mahler et al. 2014; Grace et al. 2007), whereas the control of phasic firing in the VTA involves the pedunculopontine tegmentum (PPTg) and the laterodorsal tegmentum (Floresco et al. 2003; Lodge and Grace 2006). Burst firing in the SNc is driven by direct STN projection and the PPTg (Chergui et al. 1994; Futami et al. 1995; Hammond et al. 1978) and tonic firing in the SNc is regulated by the VP. Whereas projections from the VP to the DA and non-DA neurons in the VTA is monosynaptic (Hjelmstad et al. 2013), modulation of DA activity in the SNc by the VP is likely due to innervation from the substantia nigra pars reticulata (SNr). Indeed, inhibitory neurons from the SNr densely innervate the SNc (Grace and Bunney 1979; Tepper and Lee 2007; Tepper et al. 1995) and the pathway from the VP to the SNr is stronger than to the SNc because GABA_A receptors are more sensitive in SNr GABAergic interneurons than SNc DA neurons (Celada et al. 1999; Grace and Bunney 1979; Tepper et al. 1995). Activity of DA neurons is also modulated by the lateral habenula (LHb) (Lecca et al. 2014). Thus, LHb stimulation has been shown to reduce firing of midbrain DA neurons (Ji and Shepard 2007; Christoph et al. 1986; Matsumoto and Hikosaka 2007). Since direct projection from the LHb to DA neurons is glutamatergic, this inhibition is suggested to be multisynaptic. A possible candidate is the rostromedial tegmental nucleus (RMTg, also called the tail of the VTA). Indeed, the RMTg receives strong innervation from the LHb (Kaufling et al. 2009) and sends strong GABAergic projections to DA neurons (Kaufling et al. 2009). It has been

shown that stimulation of the RMTg inhibits both the VTA and the SNc (Hong et al. 2011; Bourdy et al. 2014; Lecca et al. 2012), and RMTg inhibition increased DA neurons activity (Jalabert et al. 2011). Recently, Brown and collaborators have shown that RMTg neurons contribute directly to the LHb-induced inhibition of DA neurons activity (Brown et al. 2017). The RMTg receives afferents from a wide range of cerebral structures (for review see (Bourdy and Barrot 2012)) such as the cortex, the periaqueducal gray, the LDTg and PPTg, suggesting that the RMTg may integrate a variety of information that can modulate activity of both the VTA and SNc. It has been shown that a specific pathway that regulates the proportion of active DA neurons both in the SNc and the VTA is the ventral subiculum of the hippocampus (vSub)-NAc-VP-VTA/SNc pathway (Bortz and Grace 2018; Floresco et al. 2001c; Floresco et al. 2003). Two separate pathways from the NAc project to the VTA and the SNc: the NAc shell projects to the VTA and the NAc core projects to the SNc (Sesack and Grace 2010). Segregation is also likely to originate from the VP with the caudal VP projecting preferentially to the VTA and the rostral VP to the SNc (Mahler et al. 2014). Activity in the BLA decreases the population activity of DA neurons in the VTA, via a polysynaptic pathway, BLA-VP-VTA (Chang and Grace 2014) and this effect is under the control of the medial prefrontal cortex (mPFC) and, in particular, of the infralimbic cortex (Patton et al. 2013). There are major GABAergic feedback projections from the NAc/DSt and the VP into the VTA and the SNc (Kalivas 1993; Conrad and Pfaff 1976; Walaas and Fonnum 1980), forming parallel spiraling loops (Haber and Knutson 2010; Haber et al. 1990). Projections from the shell part of the NAc influence DA neurons in the VTA, which in turn projects to the core part of the NAc, which provides projections to the SNc, which sends its projection to the DSt (Haber and Knutson 2010; Haber et al. 1990). This is considered an anatomical base

for the hypothesis that transitions from goal-directed (voluntary) drug consumption to stimulus-response (habitual) drug use is dependent on this transition from medial/ventral neural circuits to more lateral/dorsal part of the VTA/SNc-NAc/DSt network (Everitt and Robbins 2005).

DA Receptor Subtypes and distribution

The diversity of DA receptors was revealed following the introduction of gene cloning procedures to the neurotransmitter receptor field (for a review see (Jaber et al., 1996)). DA receptors belong to the large family of G protein-coupled receptor (RCPG) and display considerable amino acid sequence conservation especially in their seven transmembrane domains. These receptors have been classified in 2 sub-families: D1 and D2. The D1/D2 classification concept developed in the late 1970s is still relevant even after the discovery of five distinct DA receptors. The D1 and D5 receptors are classified as "D1-like" because they share high sequence homology and stimulate the adenylate cyclase (AC) pathway and show classical pharmacological D1 responses (Jaber et al., 1996). The D2, D3, and D4 receptors are classified as "D2-like" and also share homologies. They mainly inhibit the AC pathway and show classical pharmacological D2 responses (Jaber et al., 1996). Members of the same family share common structural characteristics but have sequence variations that determine differences in their affinity for agonists and in the coupling to signal transduction pathways (Missale et al., 1998).

The D1 receptor (D1R) is the most widespread DA receptor and is expressed at a higher level than any other DA receptors (Dearry et al., 1990; Fremeau et al., 1991; Weiner et al., 1991). D1R mRNA is expressed in areas known to receive dopaminergic afferences such as the striatum, NAc and olfactory tubercle by

dopaminoceptive GABAergic neurons co-expressing dynorphin (Le Moine and Bloch, 1995; Le Moine et al., 1990) and to a lesser extent in the PFC. D1R mRNA is also found in the hypothalamus and thalamus. It is to note that in some brain areas D1R proteins are detected but not D1R mRNA suggesting that in these areas the D1 receptor is present in projections only.

The D2 receptor (D2R) gene contains exons and permits the generation of two splicing variants, named D2L and D2S (for long and short respectively) with the D2L mRNA being expressed at higher levels than the shorter variant (Jaber et al., 1996 for a review). It is mainly found in the olfactory tubercle and dorsal and ventral striatum where it is expressed by dopaminoceptive GABAergic neurons co-expressing enkephalins (Le Moine and Bloch, 1995; Le Moine et al., 1990). It is also found in the SNc and in the VTA, where it is expressed by dopaminergic neurons (Meador-Woodruff et al., 1992; Weiner et al., 1991). Thus, the D2R is expressed by both pre- and post-synaptic neurons.

The D3 receptor (D3R) has received considerable attention, mainly because it is specifically found in limbic areas such as the shell of the NAc, olfactory tubercle and islands of Calleja, and has low expression levels in the DSt (Bouthenet et al., 1991; Sokoloff et al., 1990).

Low levels of mRNA of the D4R are found in the basal ganglia whereas relatively higher levels in the frontal cortex, medulla, amygdala, hypothalamus and mesencephalon (O'Malley et al., 1992; Van Tol et al., 1991). However, the expression of the D4R remains weak when compared with expression of other DA receptors, including the D5R that is also notorious for its scarce expression. Indeed, the expression of D5R seems to be restricted to the hippocampus, the lateral mamillary nucleus and the parafascicular nucleus of the thalamus (Tiberi et al., 1991;

Meador-Woodruff et al., 1992) and the relative abundance of the DA receptors in the rat central nervous system would be D1> D2> D3> D5> D4 (Missale et al., 1998).

The co-localization of D1R and D2R within the same neuron has been the subject of debate for two decades in the literature. It seems now settled that the D1/D2 synergism is achieved not through co-localization of these specific receptors but through co-localization of receptors belonging the D1 and D2 families (Carter-Russel et al., 1995), i.e. the same neurons would express D1/D3/D4 or D2/D5 (Beaulieu and Gainetdinov, 2011).

DAT expression and function

The DAT belongs to the family of monoamine transporters that are Na⁺/Cl⁻-dependent plasma membrane proteins present at nerve endings. These transporters control the levels of monoamines by rapidly re-uptaking the released neurotransmitter and thus limiting the temporal persistence and the spatial spread of monoamines (Jaber 2006). As such, monoamine transporters constitute a major element in determining the intensity and duration of monoamine transmission. DAT mRNA was found only in DA synthesizing neurons and the corresponding protein is localized within dopaminergic innervation of several regions including ventral mesencephalon, medial forebrain bundle and dorsal and ventral striatum (Ciliax et al. 1995). At the protein level, the DAT is present perisynaptically rather than within the synaptic cleft thus promoting volume and somewhat paracrine transmission, rather than the classic synaptic transmission that is contained spatially to the targeted post-synaptic dendrites (Garris et al. 1994). Interestingly, at the sub-cellular levels, the DAT was found to be present in intracellular compartments as well as at the plasma membrane of dendrites, cell body and axon terminals. This suggests that the DAT

may be actually releasing DA at the dendritic level close to DA cell bodies (Hersch et al. 1997) and that its addressing to the membrane is regulated by the DA tone. The DAT is at the center of a great deal of attention given its key role in regulating DA transmission but also because it is targeted by psychostimulants such as cocaine and amphetamine (Jaber 2006) .

Dopaminoceptive neurons and their glutamatergic afferents

The striatum not only receives dense DA projections from the VTA and the SNc but also glutamatergic inputs from multiple brain structures and each of these glutamatergic inputs plays a role in different aspects of addictive behaviors (Sesack and Grace 2010). The NAc receives glutamatergic inputs from the PFC (the infralimbic part projecting preferentially to the NAc core and the prelimbic part to the NAc shell) (Groenewegen et al. 1999), the hippocampus and the amygdala (Britt et al. 2012; MacAskill et al. 2012). The DSt receives glutamatergic inputs from sensorimotor cortex (McGeorge and Faull 1989), and from the thalamus (Wall et al. 2013).

In the striatum, DA released in the extracellular space activates DA receptors that are located on medium spiny neurons (MSNs). MSNs in the DSt can be divided into two types based on their projection structures and peptide/DA receptors expression. D1-dynorphin-substance P-expressing MSNs project to the SNr/GPi (direct pathway to basal ganglia output structures) and D2-enkephalin-expressing MSNs project to the GPe (indirect pathway) (for review, see (Gerfen and Surmeier 2011)). In the NAc, D2R-expressing MSNs project to the VP and D1R-expressing MSNs project mainly to the VTA and SNr, but some collaterals to the VP have also been described (Chang and Kitai 1985; Berridge and Robinson 1998; Tripathi et al. 2010). However, recent studies suggest that this dichotomy is not absolute and that the direct and indirect

pathways may contain both D1- and D2-expressing MSNs (Kupchik et al. 2015). In addition, in the NAc, a subpopulation of MSNs expressing both D1R and D2R has also been described (Bertran-Gonzalez et al. 2008). DA release in the striatum promotes opposite changes in synaptic plasticity at glutamatergic synapses depending on the neuronal type. In fact, long-term potentiation occurs onto D1-expressing MSNs and long-term depression onto D2-expressing MSNs (for review see (Gerfen and Surmeier 2011)). It has been hypothesized that an imbalance between D1R- and D2R-expressing MSNs may contribute to addiction (Volkow et al. 2013; Lobo and Nestler 2011), and that each type of MSNs may play a crucial role in the regulation of addiction-like behaviors (for review see (Yager et al. 2015)), likely due to drug use-induced persistent changes in the afferent control of MSNs.

DA and reinforcement/reward processes

The first indirect demonstration of a role of the DA system in reinforcement and reward circuit was the discovery by Olds and colleagues that brain stimulation of specific brain areas was rewarding (Olds and Milner 1954). Although at the time DA was not considered a neurotransmitter, these experiments clearly demonstrated that a specific circuit was involved in reinforcement.

Importantly, when it was established that DA is a neurotransmitter (Carlsson et al. 1957; Carlsson and Lindqvist 1963), it was rapidly demonstrated that it played a major role in rewarding processes as lesions of this system, and notably the nigro-striatal pathway, by 6-hydroxy-DA causes severe adipsia and aphagia in animal models (Ungerstedt 1971; Olds and Milner 1954). In addition, it was found that dopaminergic neurons were concentrated in areas that maintained self-stimulation, these experiments set up the background for experiments that demonstrated that self-stimulation reward depends on DA neurotransmission (Fouriez et al. 1978).

In parallel to these experiments, Wise and colleagues started investigating the role of DA in the rewarding effects of more physiologically relevant stimuli such as food (Wise et al. 1978). They found that administration of the dopaminergic antagonist pimozide eliminated food reward. The effects of pimozide resembled those of extinction under condition of withdrawal of food reinforcement which lead Wise to suggest that pimozide prevented rats to experience the positive, rewarding consequence of consumption of food (Wise 1978). Several experiments have confirmed and extended this finding. Importantly, DA does not appear indispensable to drive feeding but it is necessary to motivate animals to exert an effort to obtain food (Salamone et al. 2007) and it appears to be sensitive to the novelty and the palatability of food (Di Chiara and Bassareo 2007). Finally, DA has been shown to be central for sexual reward and in particular for its appetitive aspects (Damsma et al. 1992; Everitt 1990; Fibiger et al. 1992; Micevych and Meisel 2017).

A critical contribution to the understanding of the role of the DA system in reward processes comes from the work of Wolfram Schultz (Schultz 2016; Schultz et al. 1997). Using electrophysiological recording of dopaminergic region (notably the SN) in behaving non-human primates, they found that DA neurons respond to novel or better than expected rewards but that, upon repeated exposure and learning, they cease to respond to reward itself and instead they respond to stimuli that are predictive of reward and that are unexpected because their occurrence is not signaled (Schultz 2016; Schultz et al. 1997). Moreover, they found that DA neurons are inhibited when a reward is withdrawn or is worse than expected leading to the postulate that DA represent a learning signal for reward prediction errors (Schultz 2016; Schultz et al. 1997). Thus, increases in DA activity signal that something positive has occurred and that the current knowledge about rewards needs to be

updated to take into consideration the new knowledge and increase the probability of repeating the actions that produce the reward (Schultz 2016; Schultz et al. 1997). Conversely, decreases in DA activity signal that the expected reward is no longer available (ex. depletion of food resources) and current knowledge needs to be updated to take into consideration this novel information and decrease the probability of repeating the actions that fail to produce the reward and to increase the probability of implementing new actions (Schultz 2016; Schultz et al. 1997). Finally, DA activity does not change when reward is obtained as expected and knowledge of the state of the world does not need to be changed or updated (Schultz 2016; Schultz et al. 1997). Other theoretical framework to explain the role of DA in reward processes have been proposed such as incentive motivation (Berridge and Robinson 1998; Robinson and Berridge 1993), habit formation (Graybiel 1998; Everitt and Robbins 2005), aberrant learning and memory processes (Di Chiara et al. 1999; Hyman 2005; Hyman et al. 2006) and effort-related aspects of motivation (Salamone et al. 2007).

In humans, brain-imaging studies strongly suggest a role for the DA system in non-drug rewards. Indeed, the dopaminergic system is activated by food consumption and food-related stimuli, especially in people with obesity and eating disorders (Volkow et al. 2017). In addition, the dopaminergic system appears involved in sexual attraction and arousal as well as in romantic love (Fisher et al. 2005; Georgiadis et al. 2012; Ruesink and Georgiadis 2017). Interestingly, in humans, monetary reward is also associated with the activation of the DA system indicating that this system is able to respond to abstract and cultural constructions of reward (Koepp et al. 1998; Zald et al. 2004).

Drugs, DA and reinforcement

One of the key discoveries in determining the role of DA in brain reward processes and in addiction was the demonstration that all drugs, regardless of their specific mechanism of action, increase the extracellular levels of DA in the striatum. Using transversal microdialysis, Di Chiara and collaborators were able to demonstrate that cocaine, amphetamine, morphine, nicotine and alcohol (Imperato and Di Chiara 1986; Di Chiara and Imperato 1988) increase DA levels preferentially in the ventral striatum. A decade later, using vertical microdialysis probes that allow more precise sampling of different subregions of NAc it was found that acute, passive injections of drugs increase DA levels preferentially in the Shell rather than the Core region of the NAc (Pontieri et al. 1995; Pontieri et al. 1996; Tanda et al. 1997). Importantly animals self-administering cocaine according to a fixed-ratio (FR1) schedule control the frequency of cocaine injections in order to titrate DA levels in order to maintain them above a certain threshold (Wise et al. 1995b). These effects have been confirmed and extended to other drugs such as heroin, amphetamine, nicotine and cannabinoids (Lecca et al. 2006a; Lecca et al. 2006b; Lecca et al. 2007; Ranaldi et al. 1999; Wise et al. 1995a). Whereas DA in the NAc is indispensable for the initial effects of drugs and in the development of addiction when behavior is goal-directed, when behavior becomes more habitual, the dorsal striatal regions take over and play a major role (Zapata et al. 2010). In addition, the involvement of the NAc Core and the more dorsal region effects may be particularly critical for conditioned responses. For example, Ito et al (Ito et al. 2004; Ito et al. 2002; Ito et al. 2000) found that in rats self-administering cocaine according to a second order schedule that is maintained by drug cues, DA elevations induced by conditioned stimuli are specific for the NAc core and that drug seeking maintained by these cues was specific of the DSt. In

addition, blockade of DA receptors in the DSt prevents cocaine seeking (Belin and Everitt 2008; Vanderschuren et al. 2005) and relapse after prolonged abstinence (See et al. 2007).

Fast scan voltammetry can measure changes in DA levels with sub-second resolution and allows investigating changes in phasic DA releases (Robinson et al. 2003; Phillips et al. 2003a). This technique has also been instrumental to characterize the role of DA in addiction. In particular, it has been shown that injections of cocaine increase DA levels in the NAc in rats that actively self-administer this drug (Phillips et al. 2003b) but also that, in trained animal, increases of DA in the NAc levels occur before animals actually perform the action (lever press) that leads to cocaine delivery, further supporting the role of DA in driving drug-seeking behavior (Phillips et al. 2003b). Finally, it has been shown that phasic DA release decreases in the NAc, whereas it increases in the DSt after several weeks of access to short sessions of cocaine self-administration (Willuhn et al. 2012). However, under conditions of extended access to cocaine, the increase in the DSt appeared transient and after 3 weeks of cocaine escalation, no increase in DA levels was found following self-injections of cocaine (Willuhn et al. 2014). These results suggest that DA elevations occur in the initial stages of addiction but not after repeated excessive intake of drugs which may instead be associated with a sort of DA depletion as supported by the fact that L-dopa treatment restored DA signaling and blocked escalation of cocaine intake (Willuhn et al. 2014).

DA, drug withdrawal and negative affect

If drugs increase DA levels in the NAc when administered acutely or repeatedly, what happens when drug administration is discontinued? Homeostatic principles and the “opponent processes” theory (Solomon and Corbit 1974), predict that the body

should adapt to drug effects and produce adaptations that oppose and counteract these effects and therefore, DA levels in the NAc are expected to drop below basal levels, which would result in aversion and low mood. This concept and its role in addiction have been further developed by Koob and Le Moal in their allostatic theory of addiction (Koob and Le Moal 2001), which postulates that upon repeated cycles of intoxication and withdrawal the brain would be overwhelmed by drug-induced neuroadaptations and would not be able to restore normal activity levels of the reward system. Thus, individuals would find themselves locked in a low emotional state that would make them particularly vulnerable to the motivational effects of drug-related stimuli and stressors leading to accrued risks of relapse. Indeed, basal DA levels have been shown to be lower than normal during acute withdrawal from cocaine (Parsons et al. 1991; Weiss et al. 1992; Rossetti et al. 1992), morphine (Pothos et al. 1991; Rossetti et al. 1992; Crippens and Robinson 1994; Acquas et al. 1991), amphetamine (Rossetti et al. 1992; Crippens and Robinson 1994), alcohol (Rossetti et al. 1992; Weiss et al. 1996), nicotine (Hildebrand et al. 1998) and THC (Tanda et al. 1999). Conversely, activity of VTA DA neurons measured by in vivo electrophysiology has been shown to be altered with decreases found during withdrawal from alcohol (Diana et al. 1992b), morphine (Diana et al. 1995), THC (Diana et al. 1998) and increases found after cocaine (Ackerman and White 1990; Zhang et al. 1997), amphetamine (Zhang et al. 1997) and nicotine (Rasmussen and Czachura 1995).

In parallel, intracranial self-stimulation, a measure of the sensitivity of the reward system, has demonstrated that rats have deficits in self-stimulation reward after withdrawal from cocaine (Markou and Koob 1991), amphetamine (Paterson et al. 2000; Wise and Munn 1995), morphine (Schulteis et al. 1994), alcohol (Schulteis et

al. 1995) and nicotine (Epping-Jordan et al. 1998; Kenny and Markou 2001). Altogether, these findings suggest that withdrawal from drugs is associated with decrease activity of the dopaminergic system that may participate to a negative state and drive negative reinforcement. However, it should be noticed that most if not all of these effects have been found after hours or days of withdrawal and most of these dysregulations recover upon protracted abstinence (Koob 2006, 2008). Therefore, it is likely that neurotransmitter and neuroendocrine systems other than the DA systems are responsible for the persistent negative effect that may be associated with long-term abstinence (Koob 2006, 2008).

DA, craving and relapse

In addition to the phases of intoxication and withdrawal, DA plays a key role in craving and relapse to drugs. First of all, administration of low doses of the drug (i.e. drug priming), that induce DA elevations in the NAc, is a strong trigger of reinstatement (Shaham et al. 2003; Shalev et al. 2002) and these effects appear at least in part to be mediated by DA because DA receptor antagonists reduce drug seeking behavior (Shaham et al. 2003; Shalev et al. 2002; Andreoli et al. 2003; Higley et al. 2011; Norman et al. 1999; Liu and Weiss 2002). In addition, direct injections of DA agonists in the NAc trigger relapse to cocaine (Bachtell et al. 2005; Schmidt et al. 2006). Cue-induced and stress-induced reinstatement appears also to depend, at least in part, on DA neurotransmission (Mantsch et al. 2016; Crombag et al. 2008; Feltenstein and See 2008). However, when it comes to relapse, dopaminergic regions other than the NAc such as the DSt, the amygdala, the PFC and the hippocampus appear to play a major role in this phenomenon (Feltenstein and See 2008).

Addiction and DA brain imaging

In the last thirty years, brain-imaging techniques have provided considerable insights into the mechanisms underlying addiction (Volkow et al. 2009; Volkow et al. 2004). These non-invasive approaches have allowed for the first time to investigate the functional and structural alterations that are associated with addiction in humans allowing for the investigation of the changes that occur at different stages of addiction. Altogether, these studies have confirmed that drugs of abuse activate DA systems and that major modifications occur in the DA system in individuals suffering from addiction. For example, DA release induced by drugs has been estimated by PET brain imaging quantifying the displacement of DA receptor ligands (Volkow et al. 2009). It has been shown that acute administration of drugs of abuse such as amphetamine (Drevets et al. 2001; Leyton et al. 2002; Martinez et al. 2005), alcohol (Boileau et al. 2003; Urban et al. 2010), nicotine (Barrett et al. 2004; Brody et al. 2004), delta-9-tetrahydrocannabinol (Bossong et al. 2009) increase DA levels in the striatum, and increases in the NAc correlate with their euphoric effects (Volkow et al. 2009). However, addiction is generally associated with reduction in DA D2R availability in the striatum and these effects appear to persist even after several months of discontinuation of drug consumption (Volkow et al. 2009; Volkow et al. 2011). More importantly, release of DA by drugs in addicted individuals is blunted compared to drug-naïve controls which suggests that a sort of tolerance, rather than sensitization, to the reinforcing effects of drugs (Volkow et al. 2009; Volkow et al. 2011). Finally, similarly to what found in animals, drug cues increased DA release in the DSt but not the NAc (Volkow et al. 2006; Wong et al. 2006) and in the PFC (Milella et al. 2016) in people with cocaine use disorder but also in recreational cocaine users (Cox et al. 2017). Although functional magnetic resonance imaging

(fMRI) approaches cannot discriminate the effects of specific neurotransmitters, they also show that drugs and drug-related stimuli affect the activity of dopaminergic areas (Suckling and Nestor 2017; Zilverstand et al. 2018).

One of the most replicable alterations found in people suffering from addiction compared to controls, is that the levels of dopaminergic D2R in the striatum are decreased (Volkow et al. 2009; Volkow et al. 2004). Conversely, high levels of D2R reduce the positive effects of psychostimulants (Volkow et al. 2002; Volkow et al. 1999) and therefore, low levels of D2R could represent a risk factor for the development of addiction to psychostimulants as suggested by studies in animal models (Dalley et al. 2007; Morgan et al. 2002; Nader et al. 2006).

The development of brain imaging techniques for laboratory animals have further expanded the ability of these approaches and will allow to provide more mechanistic insights into addiction processes (Caprioli et al. 2013; Jonckers et al. 2015; Gould et al. 2014). Brain imaging studies in animals have mostly focused on cocaine, because this drug produces the most reliable self-administration behavior and allows producing addiction-like phenotypes in rats (Ahmed and Koob 1998; Deroche-Gamonet et al. 2004; Vanderschuren and Everitt 2004). In the last fifteen years, studies in non-human primates have allowed better understanding of addiction processes and its evolution over time (Gould et al. 2014). For example, they have demonstrated that over the course of chronic self-administration the reduction in brain metabolic activity induced by cocaine shifts from the ventral to the DSt (Porrino et al. 2004). In rats with a history of escalation of cocaine self-administration reduction in brain metabolic activity of the ventral striatum has been reported after short periods of withdrawal (Calipari et al. 2013; Gozzi et al. 2011; Nicolas et al.

2017). However, after longer periods of abstinence, the reduction in brain metabolic activity was found in dorsal striatal regions (Nicolas et al. 2017).

Electrophysiological effects of drugs on the DA system

Effects of acute administration of drugs on the activity of DA neurons

The population activity (i.e. the proportion of DA neurons that are active at a given time), the firing rate and the firing pattern of SNc and VTA dopaminergic neurons are major determinants of the levels of DA released from terminals, to their targets, the DSt and the NAc respectively (Hedreen and DeLong 1991; Lynd-Balta and Haber 1994; Szabo 1979). We will describe findings that mostly refer to VTA DA neurons, which have been more widely studied in the field of addiction than SNc DA neurons.

Psychostimulants (such as cocaine and amphetamine) decrease the firing and bursting rate of DA neurons recorded from the VTA and the SNc. Cocaine blocks the reuptake of DA, and amphetamine inverts DA transport in the presynaptic neurons thereby increasing DA release, ultimately inducing the activation of negative feedback mechanism from forebrain structures. Psychostimulants also increase DA levels in the VTA, which can activate D2R present on the soma and dendrites of DA neurons, as well as D2-autoreceptors located on terminals, and further inhibits dopaminergic activity (French 1997; Lodge and Grace 2005; Belujon et al. 2016; Bunney et al. 1973; Einhorn et al. 1988; Groves et al. 1975). Notwithstanding this reduction in VTA activity, the net effect of psychostimulant administration is an increase in DA levels in the NAc and the VTA, which suggests that the effect of cocaine on DAT is sufficient to elevate extracellular DA levels (Chen and Reith 1994). Acute injections of amphetamine have also been shown to decrease burst firing and population activity of DA neurons in the VTA (Belujon et al. 2016; Lodge

and Grace 2008). Most of these changes after acute drug exposure are transient, but are likely to play a role in the development of addictive behavior (for review, see (White and Wolf 1991)).

For other substances of abuse (such as ethanol, morphine, nicotine and cannabinoids), acute drug exposure increases the spontaneous activity, in terms of firing rate and bursting activity of VTA and SNc DA neurons (Diana 1998; Gessa et al. 1985; Grenhoff et al. 1986; French 1997; Budygin et al. 2004). Opiates act on mu-opioid receptors and disinhibit DA neurons in the VTA through indirect inhibition of GABAergic inputs (Johnson and North 1992; Luscher and Ungless 2006). Systemic and intra-VTA infusion of morphine has also been shown to increase the firing rate and bursting activity of DA VTA (Jalabert et al. 2011). Although the mechanism of action of alcohol has not yet been fully understood, it is established that glutamate and GABA receptors play an important role in alcohol reinforcement (Luscher and Ungless 2006). Intra-VTA administration of ethanol produces an increase of DA levels in the NAc, suggesting that one the main sites for ethanol reinforcement is the VTA and the consequent activation of DA neurons (Beveridge et al. 2009). Consistent with this idea, it has been shown that ethanol is voluntarily self-administered in the VTA (McBride et al. 1999). *In vivo* (Foddai et al. 2004; Mereu and Gessa 1984) and *in vitro* (Brodie et al. 1999) studies have shown that ethanol stimulates the firing activity of both VTA and SNc DA neurons, mediated via the inhibition of potassium M-current (Koyama et al. 2007). This increase in spontaneous activity of DA neurons is associated with an increase in DA release in the NAc and DSt (Fadda et al. 2005; Zhang et al. 2009).

Acute administration of psychostimulants has been shown to suppress NAc neurons firing by multiple mechanisms (for review see (White and Kalivas 1998)). In fact, using *in vivo* preparations, it has been shown that psychostimulants elicit either an inhibitory or excitatory response in NAc MSNs, depending on the depolarized (up) or hyperpolarized (down) state, where inputs from the hippocampus can gate cortical inputs by driving NAc neurons in the up states (O'Donnell and Grace 1995). Indeed, suppression of hippocampal afferents results in the elimination of the “up” state of NAc MSNs, and the failure to evoke spike firing after activation of prefrontal inputs to the NAc (O'Donnell and Grace 1995). Acute injections of methamphetamine induce a loss of the up and down states fluctuations in the NAc, with a resting membrane potential more depolarized than baseline (Brady et al. 2005). It should be noted that the information flow from principal glutamatergic inputs to NAc MSNs arising from the prefrontal cortex, hippocampus and amygdala (Groenewegen et al. 1999) depends on DA (Floresco et al. 2001a; Belujon and Grace 2014; Floresco et al. 2001b). Indeed, changes in DA release in the NAc have been shown to have an effect on DA inputs as well as glutamatergic inputs onto MSNs.

Other drugs, such as ethanol, nicotine or opioids affect MSNs. For example, acute ethanol has been shown to inhibit MSNs in the DSt via changes in cholinergic tone (Blomeley et al. 2011). Indeed, ethanol inhibits cholinergic and low-threshold interneurons, while it excites fast-spiking ones. The reduction of acetylcholine induces a large hyperpolarization of MSNs, which leads to downregulation of GABA and glutamate receptors in MSNs (Blomeley et al. 2011). Conversely, acute opioids in the DSt appear to inhibit cortico-striatal synapses at the presynaptic level. Indeed, it has been shown that *in vitro* opioids agonists selective for mu- and delta- receptors presynaptically inhibit excitatory synaptic potentials in MSNs, whereas only delta-

selective agonists decrease inhibitory synaptic potentials (Jiang and North 1992). Acute opioids have also been shown to produce long-term depression (LTD) in the DSt. Thus, *in vitro*, application of an opioid receptor agonist induced a long-lasting reduction of electrically evoked EPSCs amplitude in the DSt (Atwood et al. 2014). In addition, a robust inhibition of glutamatergic synaptic inputs to MSNs neurons by application of a mu-agonist has been reported also *in vitro* in the NAc (Hoffman and Lupica 2001). The same study reported no effect of mu-agonists of GABAergic synaptic responses in the NAc. On the other hand, glutamatergic inputs are only marginally inhibited by cannabinoids, whereas GABAergic inputs are consistently inhibited (Hoffman and Lupica 2001). It has been suggested that this presynaptic inhibition is due to activation of CB1 receptors located on the terminals of a small population of interneurons in the NAc, playing a critical role in feed-forward inhibition of MSNs outputs (Hoffman and Lupica 2001). Finally, acute nicotine has been shown to facilitate endocannabinoid (eCB)-long-term depression in the DSt (Adermark et al. 2018). This facilitation is occluded by a D2 agonist, as well as a muscarinic acetylcholine antagonist, suggesting that nicotine affects DA and cholinergic neurotransmission in the DSt (Adermark et al. 2018). Indeed, it has been shown behaviorally that nicotine releases endogenous cannabinoid that can produce THC-like discriminative effects (Solinas et al. 2007) and that this effect resembles that of DA agonists (Solinas et al. 2010a), suggesting that the three systems convey converging information (for review see (Gamaledin et al. 2015)).

Therefore, the actions of different drugs of abuse on synaptic processes in the striatum participate in the modulation of activity of MSNs and likely contribute to addictive properties of these drugs.

Effects of chronic administration of drugs on the neuronal activity of DA neurons

Whereas acute administration of psychostimulants induces a DA-mediated feedback inhibition of DA neurons activity, after a chronic regimen, and eventually the discontinuation of the drug administration, different and sometimes opposite effects can be found. For example, seven days after withdrawal from chronic cocaine, VTA DA neurons population activity is increased (Lee et al. 1999; Gao et al. 1998). This effect is reversed by blockade and desensitization of D2R, suggesting changes in the function of D2R involved in feedback inhibition in the VTA-NAc pathway (Lee et al. 1999; Gao et al. 1998). Wolf and collaborators in 1993 have also shown in chronically amphetamine-treated rats that VTA DA neurons were sub-sensitive to the D2 agonist quinpirole after short-term withdrawal but not after longer period of withdrawal (Wolf et al. 1993). Repeated cocaine administration induced an increase in the firing rate and the number of spontaneously active DA neurons in the VTA (Henry et al. 1989).

Chronic treatment with cannabinoids has been shown to not alter spontaneous activity of VTA and SNc DA neurons *in vivo* 24h after the last injection of cannabinoid (Wu and French 2000). In contrast, chronic morphine (Diana et al. 1995) and nicotine (Rasmussen and Czachura 1995) induce a significant decrease in the firing rate and the bursting rate of VTA neurons and chronic nicotine had no effect on SNc DA neurons (Rasmussen and Czachura 1995). *In vivo*, chronic ethanol has been shown to induce an increased in the spontaneous activity of VTA DA neurons (Diana et al. 1992a) whereas *in vitro* studies found no differences between ethanol and naïve-treated rats (Brodie 2002). However, in this study, a sensitization to ethanol

excitation of DA VTA neurons has been observed, suggesting that alcohol induced adaptations of DA neurons. Indeed, DA neurons undergo adaptive changes that might be unmasked during withdrawal from drugs of abuse (Brodie 2002).

Effects of chronic administration of drugs on the activity of NAc and DSt neurons

Several studies have suggested that the NAc and the DSt are subjected to neurobiological changes as drug use develops from acute to chronic, both in animal models (Macey et al. 2004; Nader et al. 2002) and in humans (Hanlon et al. 2009). For example, during chronic self-administration of cocaine, MSNs recorded from the NAc core show a strong phasic inhibition during the beginning of the operant training, that become blunted across sessions (Coffey et al. 2015). However, the activity of MSNs recorded from the NAc shell, as well as MSNs from the DSt remained stable throughout several sessions of self-administration (Coffey et al. 2015). These results suggest that different parts of the striatum have some distinct functional neuroadaptations during chronic drug use that may participate differently in pathological processing of drug-related information.

Chronic methamphetamine reduces the up-state duration in NAc MSNs implying that MSNs will be less responsive to afferent inputs, the balance among excitatory inputs is disrupted, suggesting an inappropriate processing of irrelevant information (Brady et al. 2005). Indeed, DA in the NAc is thought to play a role in mediating the selection of appropriate input processing by NAc MSNs (O'Donnell 2003; Floresco et al. 2003). Therefore, disruption in the gating function in the NAc after chronic methamphetamine implies a disruption in the filtering of irrelevant information (Brady et al. 2005), which could lead to an increased level of salience for drug-associated

cues. After protracted periods of withdrawal from chronic amphetamine (>20 days), alterations in the cortico-accumbens circuit have also been observed, i.e. increased bistable membrane activity, and increased intercellular coupling in the PFC and NAc (Onn and Grace 2000). Interestingly, these changes were not observed during the chronic administration, suggesting that withdrawal from amphetamine unmasked alterations in the cortico-accumbens circuit (Onn and Grace 2000).

Addiction, DA and Synaptic plasticity

Addiction has long been associated with pathological neuroplasticity in the mesolimbic system, as reviewed thoroughly previously (Luscher and Malenka 2011; Gipson et al. 2014; Jones and Bonci 2005; Kauer and Malenka 2007; Hyman et al. 2006; Nestler 2001; Wolf and Ferrario 2010; Russo et al. 2010). Here we review the main findings in different regions of the DA system.

VTA

Synaptic plasticity in the VTA is modulated by different drugs of abuse and is thought to contribute to different and specific aspects of addiction. Both excitatory and inhibitory inputs onto DA neurons in the VTA undergo long-term changes after acute and/or chronic exposure to drugs (Luscher and Malenka 2011). Many studies have demonstrated an enhancement of glutamatergic synaptic transmission in the VTA induced by acute (Ungless et al. 2001; Wanat and Bonci 2008) and chronic (Borgland et al. 2004) cocaine. A seminal study, published in 2001 has shown that an acute injection of cocaine was sufficient to induce long-term potentiation of AMPA-mediated currents at excitatory synapses onto DA VTA neurons, that persists until 5 days after the injection; an occluded induced-LTP as well as an enhanced LTD has also been described (Borgland et al. 2004) suggesting that this potentiation might be

involved in the early stages of the development of addiction. For both acute and chronic studies, this potentiation was transient and lasted less than 10 days (Ungless et al. 2001; Wanat and Bonci 2008; Borgland et al. 2004), and might contribute to the increase in DA neurons firing observed after a 24h withdrawal from chronic cocaine (Henry et al. 1989). A single exposure to ethanol has also been shown to induce synaptic plasticity changes. Thus, acute ethanol abolishes completely the long-term depression induced by high frequency stimulation (HFS) of GABAergic inputs to the VTA (Guan and Ye 2010), and it reduces the expression of LTP at glutamatergic synapses onto VTA neurons (Wanat et al. 2009). Another study has shown that self-administration of cocaine, but not passive cocaine infusions, produced persistent potentiation of VTA excitatory synapses, still present after 3 months of abstinence and resistant to behavioral extinction, whereas sucrose or food self-administration induced only a transient potentiation of VTA glutamatergic signaling (Chen et al. 2008). These data suggest that the pharmacological effect of cocaine itself is not sufficient to potentiate VTA glutamatergic signaling, and that the drug-context pairing is necessary to induce persistent changes in synaptic plasticity. Similarly, amphetamine enhances AMPA/NMDA ratio at excitatory synapses onto VTA DA neurons (Faleiro et al. 2004), although the specific mechanism seems different from cocaine (Wolf and Tseng 2012).

Chronic ethanol induced an increase in AMPA receptor functions in the VTA (Stuber et al. 2008), which is similar to LTP-processes (Bonci and Malenka 1999), and may facilitate the drive to consume ethanol. Acute morphine induced a loss of potentiation at GABAergic synapses onto VTA DA neurons and enhanced glutamatergic synapses 24h after exposure, as described with nicotine. Thus, all addictive drugs tested *in vitro* induce a potentiation of AMPA/NMDA ratio in VTA DA neurons after *in*

vivo administration of a single dose (Borgland et al. 2004). Moreover, addictive drugs (such as cocaine, nicotine and morphine) have been shown to induce the insertion of GluA2-lacking calcium-permeable AMPAR (CP-AMPA) into the synapses in the VTA (Argilli et al. 2008; Bellone and Luscher 2006; Brown et al. 2010)

SNc

A single exposure to drugs and in particular psychostimulants, also alter the activity of SNc DA neurons and play a role in the increased expression of behavior mediated by DA release in the DSt (Zahniser et al. 1998; Wolf 1998). A recent study, using optogenetics, has highlighted changes in specific glutamatergic afferents onto SNc neurons following a single cocaine exposure (Beaudoin et al. 2018). Whereas previous studies showed that a single exposure to cocaine increases EPSCs in the VTA but failed to find an effect in the SNc (Ungless et al. 2001; Lammel et al. 2011), this could be due to the use of non-specific electrical stimulation of SNc DA neurons excitatory inputs. Thus, cocaine-induced plasticity appears to be dependent on the types of afferents involved, specifically the STN-SNc and PPTg-SNc (Beaudoin et al. 2018). DA neurons from the SNc can selectively modulate the information processed by different input nuclei. For example, the AMPA/NMDA ratio is decreased at the PPTg-inputs to SNc, whereas it is unchanged in the STN-inputs to SNc (Beaudoin et al. 2018).

NAc

Several studies have found that cocaine exposure, whether acute or chronic, alters the capacity of NAc glutamatergic synapses to undergo synaptic plasticity (Fourgeaud et al. 2004; Goto and Grace 2005; Yao et al. 2004; Martin et al. 2006).

Synaptic plasticity is often studied by examining excitatory synaptic transmission using the ratio AMPA/NMDA-mediated excitatory postsynaptic currents (EPSCs). Unlike in the VTA, a single cocaine injection is not sufficient to cause by itself changes in the synaptic strength in the NAc, when measured 24h later (Kourrich et al. 2007; Thomas et al. 2001), but it can induce changes in the induction of low-frequency induced-LTD in indirect pathway MSNs (Grueter et al. 2010). However, chronic cocaine reduces AMPA receptors in NAc shell, not core, MSNs, and decreases synaptic strength at excitatory cortical synapses (Kourrich et al. 2007; Thomas et al. 2001), which is similar to what has been described after chronic amphetamine, i.e. a decrease in AMPA receptor levels in the NAc (Lu and Wolf 1999). These synaptic adaptations in the NAc have been shown to follow and to depend on changes in the synaptic function described in the VTA, i.e. potentiation of excitatory inputs (Mameli et al. 2009). Moreover, some studies have shown an increase in AMPA receptors EPSC after cocaine treatment followed by protracted abstinence (14 days), suggesting that cocaine produces a potentiation of excitatory inputs that develops or is unmasked during withdrawal (Kourrich et al. 2007; Ortinski et al. 2012; Boudreau and Wolf 2005).

In the last 15 years, particular attention has been given to changes in plasticity in the NAc which are associated with the phenomenon of incubation of cocaine craving (Grimm et al. 2001). A critical mechanism, thoroughly reviewed by Wolf and collaborators (Wolf 2016; Dong et al. 2017) is the synaptic accumulation of CP-AMPA at excitatory synapses on MSNs in the NAc. Indeed, whereas GluA2-containing calcium-impermeable AMPA (CI-AMPA) receptors are responsible for the majority of excitatory synaptic transmission in drug-naïve rats or during short withdrawal from chronic cocaine, during protracted abstinence, the contribution of

CP-AMPA receptors increases, which leads to an increased responsiveness of MSNs to glutamate (Purgianto et al. 2013). Other studies highlighted the role of silent synapses in the incubation of craving (Lee et al. 2013; Ma et al. 2014). First, it has been shown that silent synapses (expressing stable NMDA receptors but absent or labile AMPA receptors) become unsilenced as the withdrawal period progresses, a process that involves CP-AMPA receptors. Recruitment of CP-AMPA receptors has been shown in amygdala-NAc synapses (Lee et al. 2013) as well as in infralimbic PFC-NAc synapses (Ma et al. 2014). Interestingly, optogenetic stimulation induces downregulation of CP-AMPA at these two different synapses but reduces incubation of craving only in amygdala-NAc synapses, whereas downregulation of CP-AMPA receptors in infralimbic PFC-NAc synapses increases incubation of craving (Lee et al. 2013; Ma et al. 2014), suggesting that reorganization of specific circuits is critical for relapse.

Dorsal striatum

Whereas drug-induced deregulation of synaptic plasticity has been well studied in the NAc, limited evidence exists regarding drug-induced synaptic plasticity in DSt. A study in primates chronically exposed to ethanol has shown that MSNs of the putamen exhibited enhanced glutamatergic transmission as well as an increased intrinsic excitability (Cuzon Carlson et al. 2011).

Chronic nicotine induced a change in the regulation of cortico-striatal synaptic plasticity, in particular in the indirect pathway (D2 MSNs). Indeed, whereas in control animals, application of HFS normally induced an LTD, in animals chronically treated with nicotine, HFS induced an LTP, therefore inverting the direction of plasticity (Xia et al. 2017), an effect likely mediated by disruption of DA signaling (Koranda et al.

2014). Metabotropic glutamate receptors of subtype 5 (mGluR5) in the striatum are necessary for the induction of LTD (Sung et al. 2001) and mGluR5 function appears to be altered following abstinence from cocaine self-administration (Knackstedt et al. 2014). In particular, mGluR5-dependent LTD in DSt slices could not be induced following 3 weeks of abstinence from self-administration which was restored by the application of a positive allosteric modulator of mGluR5 (Knackstedt et al. 2014). This led the authors to suggest that pharmacological enhancement of GluR5 function could be a novel target to restore normal plasticity in the DSt and reduce the risk of relapse.

Addiction and DA structural plasticity

Drugs of abuse produce structural changes within the brain, and in particular within the mesolimbic dopaminergic system. The term “structural plasticity” refers to the reorganization of synaptic connections in brain circuits (Robinson and Kolb 2004). The interaction between neurons can be physically increased by the formation of new synaptic “boutons” and/or of new dendritic spines or reduced by their elimination. Dendritic spines are highly dynamic and the morphology of dendritic arborization can evolve rapidly to modulate the synaptic contacts between neurons (Berry and Nedivi 2017). The first demonstration that drugs of abuse can increase the dendritic arborization of neurons was provided by Kunz and collaborators that showed an increased spine density in the hippocampus in response to alcohol (Riley and Walker 1978; Kunz et al. 1976). Later on, Kolb and Robinson showed that chronic administrations of cocaine and amphetamine produce an increase in spine density on MSNs in the NAc shell and core (Lee et al. 2006; Maze et al. 2010; Norrholm et al. 2003; Robinson et al. 2001; Robinson and Kolb 1997, 1999). Chronic cocaine administration also increased spine density in the PFC, a brain area also targeted by

DA neurons from the VTA (Robinson and Kolb 2004). Similar increases have also been described in response to nicotine (Brown and Kolb 2001), MDMA (Ball et al. 2009), methamphetamine (Jedynak et al. 2007) and THC (Kolb et al. 2018). On the contrary, chronic alcohol (Zhou et al. 2007) and morphine (Robinson and Kolb 1999), decrease the spine density in the NAc. These structural changes, initially described in response to repeated administration of drugs, have also been found after a single administration of cocaine (Dos Santos et al. 2017). Interestingly, abstinence from drug has also been shown to produce alterations in dendritic spines that may participate in the withdrawal symptoms and the negative affect state that characterize this phase of addiction (Spiga et al. 2014a). Thus, acute or chronic administration of drugs produce long-lasting structural changes in the brain and particularly in the connectivity between DA neurons and striatal neurons.

Striatal structural changes in MSNs induced by drugs are mediated, at least in part, by DA. Indeed, structural changes found in culture of MSNs or *in vivo* after administration of cocaine depend on increases in the extracellular DA concentration (Fasano et al. 2013; Martin et al. 2011) and the stimulation of DA receptors (Dos Santos et al. 2017; Ren et al. 2010; Zhang et al. 2012). The increase in the density of dendritic spines observed after cocaine exposure was initially reported in both D1R- and D2R-expressing DA neurons but appears to be more durable in D1R-expressing neurons (Lee et al. 2006). More recent reports also indicate that this increase mainly occurs in D1R-expressing MSNs (Gipson et al. 2014; Gipson and Olive 2017; Kim et al. 2011; Russo et al. 2010). Interestingly, it was recently shown that this increased number of spines of MSNs is not associated with an increased number of

glutamatergic boutons (Heck et al. 2015), but instead with an increased number of dopaminergic boutons in the NAc Core (Dos Santos et al. 2018).

These drug-induced neuroplastic structural adaptations might modify communications within the dopaminergic system and could be part of the “drug memory” responsible for the high risk of relapse to drug addiction. In fact, NAc MSNs play a central role in the integration of cortical and mesencephalic afferents in distal spines of dendrites. This striatal “microcircuit” is thought to allow DA to influence cortical glutamatergic axons (Spiga et al. 2014b). Consequently, changes in dendritic spines by drugs might participate in synaptic plasticity described in response to drugs and in the behavioral consequences of their administration. For example, blockade of the Mitogen Activated Protein kinase ERK pathway prevents formation of new synapses in response to cocaine *in vitro* and *in vivo* (Dos Santos et al. 2017) as well as the behavioral effects of cocaine (Valjent et al. 2010; Valjent et al. 2006). Therefore, the long-lasting structural remodeling induced by acute or repeated drugs might play a role in the development and persistence of drug addiction (Brown et al. 2011; Robinson and Kolb 2004).

Drugs and transcriptional plasticity in DA regions

The consequence of drug-induced increases in DA transmission in the NAc is an over-stimulation of post-synaptic MSNs by activation of dopaminergic receptors, mainly D1R and D2R, that modulate the cyclic AMP pathway (Jaber et al. 1996). In addition, dopaminergic receptors interact with pathways activated by glutamatergic transmission and particularly the pathway involving ERK (Girault et al. 2007; Valjent et al. 2000). These rapid and transient effects of cocaine play a major role in triggering long-term molecular adaptations within dopaminergic regions, and particularly in the ventral and dorsal striatum and PFC (Pierce and Kalivas 1997;

Pierce and Wolf 2013; Wolf 2010, 2016). In fact, drugs activate constitutive transcription factors, such as the cAMP responsive element binding (CREB), that produce persistent changes in neuronal functioning that are believed to be central for addiction (Briand and Blendy 2010; Carlezon et al. 2005; Nestler 2004). Interestingly, CREB activation has also been shown to participate to the formation of new dendritic spines (Brown et al. 2011). CREB targets several genes including those of the immediate early gene *fos* or *egr* family (Beckmann and Wilce 1997; Janknecht 1995), which encode inducible transcription factors, and as a consequence, produce durable protein expression changes. Acute administration of cocaine (Hope et al. 1992; Thiriet et al. 2001a), amphetamines (Moratalla et al. 1992; Thiriet et al. 2001b; Graybiel et al. 1990), nicotine (Kiba and Jayaraman 1994; Ren and Sagar 1992), opiates (Bontempi and Sharp 1997; Garcia et al. 1995; Liu et al. 1994), THC (Mailleux et al. 1994; Valjent et al. 2002) and alcohol at high doses (Thiriet et al. 2000) have been shown to trigger the expression of immediate early genes of the *fos* and *egr* families in the striatum and the PFC. Usually, the expression of transcription factors of the Fos protein family is rather transient and tolerance to this response has been described with chronic passive administration of cocaine (Alibhai et al. 2007; Hope et al. 1992). In contrast, chronic administration of drugs was shown to produce an accumulation of Delta-Fos B (DFosB) in several brain regions, involved in addiction, namely the NAc, the DSt, the PFC, the hippocampus and the amygdala (Hope et al. 1994; McClung et al. 2004; Nestler et al. 2001; Perrotti et al. 2008) DFosB is a truncated form of the FosB protein, and present a much higher stability than FosB (Chen et al. 1997). Because of this stability, DFosB accumulates in neurons even after drug intake discontinuation and it is thought to be one of the main molecular adaptations involved in the long-term effects of drugs and to participate in

the switch to addiction (Nestler 2001). As for structural plasticity changes, accumulation of DFosB following drug administration takes place mostly in the D1R-expressing MSNs (Colby et al. 2003; Kelz et al. 1999; Lafragette et al. 2017; Lobo et al. 2010). By acting on this panoply of transcription factors in brain regions targeted by the dopaminergic system, drugs of abuse produce molecular adaptations and change the expression of proteins of different classes, which will have consequences both structurally and functionally within this system.

Drugs and molecular adaptations within the DA system

There are many discrepancies concerning the effects of drugs of abuse on elements of the DA system (i.e. the DA transporters and receptors) probably because of the different protocols used for drug administration (acute vs chronic, short vs long, passive vs active, etc) and the time from the last administration (from a few hours to several days) (McGinty et al. 2008). For example, some studies found a decrease in the levels of mRNA of the DAT immediately after chronic exposure to cocaine in the SN and VTA (Letchworth et al. 1997; Xia et al. 1992) whereas other studies found no effect (Arroyo et al. 2000; Maggos et al. 1997). In contrast, after 10 days withdrawal of cocaine self-administration an increase in DAT levels was found in the VTA but not in the SN (Arroyo et al. 2000), but not after passive administration (Cerruti et al. 1994). In rats, an increase in DAT binding in the NAc was found 1h after the last administration of moderate doses of cocaine (Letchworth et al. 1997) and a decrease in DAT binding after 10 days of withdrawal (Cerruti et al. 1994). However, withdrawal from cocaine self-administration results in an increase in DAT binding in rats with short access to cocaine, whereas rats that have extended access to cocaine and show escalation of self-administration do not show any difference compared to controls (Ben-Shahar et al. 2006). Finally, Letchworth et al. found a decrease in the

levels of DAT in the DSt after initial stages of self-administration but then, as self-administration becomes more chronic, they found an increase in the NAc (Letchworth et al. 2001). Importantly, these levels remained elevated after 30 days of withdrawal and returned to control levels only after 90 days of abstinence (Beveridge et al. 2009). The effects of other drugs of abuse on DAT have been less studied but no change in DAT levels was found after methamphetamine (Shepard et al. 2006) and heroin (Pattison et al. 2014) self-administration whereas alcohol increases DAT levels in the NAc of Wistar–Kyoto rats (Jiao et al. 2006).

Concerning DA receptors, controversial results have also been obtained. Increases, decreases, or no changes in the levels of D1R and D2R have been reported depending on the protocol of administration, time from the last injection and the technique of detection used (binding vs mRNA levels) for cocaine (Unterwald et al. 1994; Kunko et al. 1998; Sousa et al. 1999; Laurier et al. 1994; Ben-Shahar et al. 2007), amphetamines (Stefanski et al. 1999; Stefanski et al. 2002), alcohol (Cowen and Lawrence 2001; Hamdi and Prasad 1993; Tajuddin and Druse 1996; Eravci et al. 1997) and morphine (Georges et al. 1999). Interestingly, changes in receptor-mediated signal transduction have been shown for cocaine and morphine at the D1R leading to increased stimulation of the cAMP/ cAMP-dependent protein kinase (PKA) pathway (Hummel and Unterwald 2002; Terwilliger et al. 1991; Unterwald et al. 1996). In addition, changes at the levels of G-protein, especially to lower levels of the $G\alpha_i$ protein, which has an inhibitory function on the cAMP/PKA pathway have also been observed after drug administration (Nestler 1996; Terwilliger et al. 1991).

Although these preclinical results highlight the complexity of the neuroadaptations induced by drugs on the dopaminergic systems, as detailed earlier, studies in humans using PET tracers for DA receptors have been much more consistent and

have shown that people suffering from addiction to different drugs have lower levels of D2R in the striatum (Volkow et al. 2009; Volkow et al. 1993). In addition, methamphetamine users show also reduced striatal levels of the DAT (Chang et al. 2007).

Genetically modified animals, DA and Addiction

In the last 25 years, development of mice with mutations targeting members of the DA system has provided a wealth of information regarding the mechanisms of action of drugs of abuse and the synaptic plasticity occurring following these manipulations. Mice with a specific mutation of DAT or one of the five DA receptors have also been used to determine the specificity of action of a given drug, mainly, but not only, psychostimulants. Most studies have used classic knock-out (KO) procedures and some have implemented knock-down, knock-in or overexpression of some of these elements of the DA system. The genetic manipulation that has probably most impacted this research field may well be the knocking down of the DAT and its consequences on drug action, use, abuse and addiction. In this section, we will summarize main findings following various forms of genetic manipulation of elements of the DA system and further discuss their relevance in the behavioral and cellular responses to various drugs of abuse.

Genetic manipulations of the DAT and responses to drugs of abuse

a. DAT knock-out animals

DAT-KO mice have been generated over 20 years ago and were the first KO of any neurotransmitter transporter (Gainetdinov 2008; Giros et al. 1996; Jaber et al. 1997). Since that time, these mice have provided a wealth of information as to the action of

drugs of abuse on the DA system. Using these mice, it was shown that the DAT is involved in the lifetime control of extracellular DA (Benoit-Marand et al. 2000b; Giros et al. 1996; Jones et al. 1998), maintenance of pre-synaptic DA functions such as autoreceptor regulation (Benoit-Marand et al. 2000b; Jones et al. 1999), DA synthesis (Jaber et al. 1999) and storage mechanisms (Jones et al. 1998), as well as postsynaptic regulation such as receptor endocytosis (Dumartin et al., 2000) and gene expression (Fauchey et al. 2000a; Fauchey et al. 2000b).

DAT^{-/-} mice present a hyperactive DA phenotype that is reflected behaviorally by several abnormalities due to a dysfunction within (i) the dopaminergic mesolimbic system (locomotor hyperactivity, lack of habituation to novelty and increased response to environmental stimuli) (Spielewoy et al. 2000), (ii) the dopaminergic nigrostriatal system (motor and sensorimotor integration deficits) (Fernagut et al. 2003; Fernagut et al. 2002) and (iii) the hypothalamo-infundibular system (decreased growth hormone and prolactin secretion) (Bosse et al. 1997). An obvious phenotype observed in mice lacking DAT is the remarkable increase in their spontaneous locomotor activity during both day and night cycles (Giros et al. 1996). This increase is of the same magnitude as in normal mice treated with very high doses of amphetamine and cocaine, which are known to produce this locomotor effect by increasing the amount of striatal DA (Giros et al. 1996).

Besides providing strong direct evidence for a role of the DAT in the tight control of DA homeostasis and transmission, DAT-KO mice have provided key elements in understanding the way psychostimulants act on the DA system.

Behaviorally, DAT-KO mice do not respond to cocaine and amphetamine, when these drugs are given in their home cages, at doses that produce marked increases in locomotor activity and stereotyped behaviors in normal animals (Giros et al. 1996).

These findings are consistent with cyclic voltammetry data which shows that, in the absence of the DAT, amphetamine is completely ineffective in promoting release of DA despite being able to abolish intracellular stores of the neurotransmitter demonstrating the necessity of the DAT for the DA-releasing action of amphetamine (Giros et al. 1996). These conclusions have also been corroborated by results obtained from studies using short and long-term unilateral DAT antisense treatments in the SN of rats (Silvia et al. 1997). Indeed, the rotational behavior obtained in this model demonstrates that cocaine acts as a DAT blocker, whereas amphetamine acts as a DA releaser and this release is DAT-dependent.

Interestingly, DAT-KO mice were still able to express conditioned place preference (CPP) to cocaine (Sora et al. 1998), long term behavioral sensitization (Morice et al. 2010) and the ability to self-administer the drug, although less than control (Rocha et al. 1998). While DAT-KO still showed rewarding properties to amphetamine (Budygin et al., 2004), this psychostimulant was also able to significantly reduce the locomotor activity of the DAT-KO mice when given in a novel environment. This was also the case with methylphenidate (Gainetdinov et al. 1999). These findings lead to the conclusion that, in the absence of DAT, the rewarding properties of psychostimulants are maintained through their action on the noradrenaline transporter, which is present in the ventral parts of the NAc and that has in fact higher affinity to DA than the DAT itself (Carboni et al. 2001; Moron et al. 2002). These findings have led to the proposal that DAT inhibition is dispensable for cocaine reward, at least in DAT-KO mice, and that redundant monoamine systems might mediate responses to cocaine.

The effects of drugs of abuse other than psychostimulants were also tested on DAT-KO animals. Alcohol for instance had similar effects in DAT-KO animals and control increasing extracellular level of DA by 80% thus indicating that the DAT is not

mandatory for the effects of alcohol on DA (Mathews et al. 2006). Morphine induced higher CPP, DA levels and accumbal c-fos levels in DAT-KO mice than control littermates further indicating that rewarding responses were still present and effective in these animal models (Mathews et al. 2006). In support of this are the findings with natural rewards demonstrating that the rewarding properties of a sucrose solution are higher in DAT-KO mice than controls (Rossi and Yin 2015).

Because of continuously increasing evidence for a role of endocannabinoids in reward processes (Solinas et al. 2008), endocannabinoid signaling was assessed in DAT-KO mice and the ability of endocannabinoid ligands to normalize behavioral deficits was evaluated, namely spontaneous hyperlocomotion (Tzavara et al. 2006). DAT-KO mice had reduced striatal anandamide levels and in these mice, anandamide reuptake inhibitors attenuated spontaneous hyperlocomotion. These results suggest that maintaining an endocannabinoid homeostasis might constitute an interesting avenue to explore to counter psychostimulant-induced hyperdopaminergia and potentially their behavioral consequences.

b. Manipulation DAT expression levels and drugs of abuse

Hyperdopaminergic mutant mice were also generated by reducing expression of the DAT to 10% of wild-type levels and are referred to as DAT-knockdown animals (Zhuang et al. 2001). These animals display hyperactivity, increased grooming, stereotypy and impaired response habituation in novel environments (Zhuang et al. 2001). Additionally, these mice show higher food intake and enhanced motivation to rewarding stimuli (Cagniard et al. 2006; Pecina et al. 2003) suggesting that motivation to work for a food reward may be under the critical influence of tonic dopaminergic activity.

Another model useful for determining how inhibiting the DAT may contribute to the effects of cocaine was generated with mice that express a cocaine-insensitive DAT (DAT-CI mice). These mice bear a DAT gene that has no binding site for cocaine but can still reuptake DA (Chen et al. 2006). In these mice, cocaine suppresses locomotor activity, does not elevate extracellular DA in the NAc, and does not produce reward as measured by CPP (Chen et al. 2006). In addition, these mice do not show any cocaine-induced spine density within the NAc (Martin et al. 2011). These results suggest that blockade of DAT is necessary for cocaine reward in mice with a functional DAT. However, these mice still maintain motivation to food reward and self-administer amphetamine (Thomsen et al. 2009). Viral-mediated re-expression of wild-type DAT in the rostromedial striatum restored cocaine-induced locomotor stimulation and sensitization in DAT-CI mice (O'Neill et al. 2014). In contrast, the expression of wild-type DAT in the DSt, or in the medial NAc, did not restore cocaine-induced locomotor stimulation. This indicates that DAT-inhibition in the rostromedial striatum alone is sufficient for cocaine's locomotor stimulating effect. A recent study has used mice with different DAT expression levels: constitutive DAT knockdown (with ~5% of wild-type DAT expression levels), DAT knockdown heterozygotes (with ~50% of wild-type DAT expression levels) and inducible DAT knockdown mice, all compared to wild-type animals (Cagniard et al. 2014). The authors tested whether the effects of cocaine and amphetamine, because of their different mechanisms of actions, would be affected differently by changes in DAT expression levels. The results obtained clearly indicated that there are both presynaptic and postsynaptic mechanisms involved in differential responses to cocaine and amphetamine in relation with DAT expression levels (Cagniard et al. 2014). In other words, responses to cocaine and amphetamine seem to depend both

on psychostimulant drug doses and DAT expression levels. For instance, mice with low DAT expression levels were more sensitive to further DAT blockade by small doses of cocaine. In contrast, reduced DAT level reduced amphetamine-induced DA release through reversal of DAT, and thus reduce sensitivity to this drug (Cagniard et al. 2014).

Researchers have also generated transgenic mice overexpressing DAT (DAT-tg) which have a 3-fold increase in DAT protein levels resulting in a 40% reduction of extracellular DA concentration in the striatum (Ghisi et al. 2009). As expected, these mice have increased levels of striatal D1 (30%) and D2 (~60%) DA receptors levels and 300% increase in the basal levels of phospho-Akt in the striatum, indicative of the reduced extracellular DA tone in these animals. Behaviorally, DAT-tg animals showed increased climbing behavior after stimulation with either apomorphine or a co-administration of selective D1- and D2-receptor agonists. They also showed markedly increased locomotor responses to amphetamine accompanied by a 3-fold greater increase in the amount of DA released by amphetamine (Salahpour et al. 2008). In contrast with mice with no or low levels of DAT, overexpression of DAT resulted in reduced operant responding for natural rewards (Salahpour et al. 2008).

The val 559 mutation is a nonsynonymous single nucleotide polymorphism (SNP), converting Ala559 to Val (A559V) in the coding region of the DAT and that was found to be associated in humans with bipolar disorder and ADHD (Grunhage et al. 2000). Cells carrying this mutation show normal DAT levels, normal presence of the DAT at the membrane and normal reuptake of DA (Mazei-Robison et al. 2005). Mice bearing this mutation show elevated extracellular DA levels and less behavioral response to amphetamine with no changes in spontaneous locomotor activity (Mergy et al. 2014).

DAT contains a C-terminal PDZ (PSD-95/Discs-large/ZO-1) domain-binding sequence believed to bind synaptic scaffolding proteins (Torres et al. 2001). Disruption of PDZ domains in mice has been performed and result in almost complete loss of DAT expression mirrored by a spontaneous hyperactivity and reduced response to amphetamine (Rickhag et al. 2013). These results clearly support the proposal that PDZ domain interactions are critical for synaptic distribution of DAT *in vivo* and thereby for proper maintenance of DA homeostasis and responses to psychostimulants.

Genetic manipulation of DA receptors

The ability to make genetic changes in a predetermined way has provided an invaluable new tool to study individual players of dopaminergic transmission (Le Foll et al. 2009). Pharmacological manipulations using the psychostimulants cocaine and amphetamine have suggested that it is mainly the D1R that mediates the molecular and behavioral effects of these drugs (Kalivas and Stewart 1991). This has been confirmed using D1R-KO mice that show no further increase in their locomotor response to high doses of cocaine (Xu et al. 1994). In fact their locomotor response is decreased following psychostimulant treatment and this may be due to the contribution of the serotonergic pathway in regulating locomotor activity (Xu et al. 1994). Additionally, D1R-KO animals show no deficits in acquisition of cocaine-induced CPP (Miner et al. 1995) but do not self-administer cocaine further indicating that the D1R is critical for the reinforcing effects of cocaine (Caine et al. 2007). Interestingly, food reinforcement is maintained at normal levels in these animals (Caine et al. 2007).

The role of the D1R subtype in alcohol-seeking behaviors was studied in D1R-KO mice (El-Ghundi et al., 1998). Voluntary ethanol consumption and preference over water are markedly reduced in these animals as compared to heterozygous and wild-type controls (El-Ghundi et al. 1998). This is also the case when transgenic animals are offered a single drinking tube containing alcohol as their only source of fluid. D1R blockade with SCH-23390 effectively reduced alcohol intake in control mice to the level seen in untreated D1-KO mice. D2R blockade with sulpiride caused a small but significant reduction in alcohol intake and preference in wild type mice and only slightly attenuated residual alcohol drinking in D1R-KO mice (El-Ghundi et al., 1998). This demonstrate that the D1R has a key role in the motivation for alcohol consumption

Interestingly, inactivation of the D2 gene produced almost the opposite phenotype in the mutant mice. D2R-KO mice are akinetic and bradykinetic with significantly reduced spontaneous movement, a phenotype that resembles that observed with specific D2 antagonists as well as in extrapyramidal syndromes of Parkinson's disease (Baik et al. 1995). In a cocaine self-administration paradigm, when high doses of cocaine on the descending part of the cocaine dose-effect function were available, D2R-KO mice self-administered at higher rates than their heterozygous or wild-type littermates, but the ascending limb of the cocaine dose-effect function did not differ between genotypes (Caine et al. 2002). In addition, D2-like antagonist eticlopride increased rates of high doses of cocaine self-administration in wild-type but not in D2R-KO mice (Caine et al. 2002). Collectively, these findings suggest that although D2R is not necessary for cocaine self-administration, it is involved in some mechanism that seems to limit the rates of self-administration of high-dose of cocaine. This is in line with findings showing that the DA-releasing effects of

psychostimulants seem to be higher in these KO mice, suggesting that lower doses may produce higher effects (Rouge-Pont et al. 2002).

In an operant ethanol self-administration using a continuous access procedure, D2R-KO mice display lower ethanol-lever responding compared to wild-type mice (Risinger et al. 2000). Food lever responding and water intake were the same in both genotypes (Risinger et al. 2000). Under conditions that produced reliable place preference to ethanol in both wild-type and heterozygous (HET) mice, D2R-KO mice show no evidence of place conditioning (Cunningham et al., 2000) supporting the hypothesis that DA D2 receptors is involved in ethanol reward in mice.

The rewarding properties of morphine in the CPP paradigm are completely absent in D2R-KO mice but morphine withdrawal is unchanged in these mice (Maldonado et al. 1997). Food-induced CPP was still expressed in these animals (Maldonado et al. 1997). The ability of intravenously delivered morphine to maintain lever pressing in these mice was studied under fixed ratio 4 (FR4) schedule and a progressive ratio (PR) schedules (Elmer et al. 2002). D2R-KO mice did not respond more to morphine than to saline and did not respond more when increased ratios were required by the PR schedule. Thus, morphine failed to serve as a positive reinforcer in the D2R-KO mice indicating that the rewarding effects of morphine appear to depend critically on an intact D2R system.

D3R-KO mice showed higher basal locomotor activity levels in comparison with wild-type mice and ethanol produced similar magnitudes of CPP in both genotypes (Boyce-Rustay and Risinger 2003; McQuade et al. 2003). In a two-bottle drinking procedure, no difference was found between D3R-KO and wild-type mice in either alcohol consumption or preference (Boyce-Rustay and Risinger 2003; McQuade et

al. 2003). In an operant self-administration procedure D3-KO and wild-type mice had similar response rates to ethanol or food as well as similar water intakes (Boyce-Rustay and Risinger 2003; McQuade et al. 2003). Overall, these results indicate that D3R is not involved in either ethanol reward or intake. In contrast, D4R-KO mice displayed supersensitivity to ethanol (Rubinstein et al. 1997) as well as to psychostimulants (Katz et al. 2003). Indeed, cocaine was a more potent stimulant of locomotor activity in D4R-KO mice compared to wild-type littermate. In addition, cocaine was more potent in producing discriminative-stimulus effects in D4R-KO mice further suggesting a role of the D4R in vulnerability to stimulant abuse (Katz et al. 2003).

Finally, D5R-KO mice show normal cocaine discrimination and normal dose response activity (Elliot et al. 2003). However, cocaine dose-dependently stimulated activity at higher levels in D5R-KO mice compared to control indicating an involvement of D5R in the locomotor stimulant effects of cocaine but limited involvement in the discriminative-stimulus effects of this drug (Elliot et al. 2003).

Vulnerability to addiction

It is evident that people are not equal in relation to the risks of becoming addicts. Indeed, whereas most people try and consume drugs sporadically, only 10-30% develop an addiction over time (Everitt et al. 2008; Kreek et al. 2005; Marinelli and Piazza 2002; Piazza and Le Moal 1996; Deroche-Gamonet and Piazza 2014; Solinas et al. 2010b). This differential vulnerability is due to complex interactions between genetic factors, the effects of the drug and the environment (Deroche-Gamonet and Piazza 2014; Solinas et al. 2010b; Kreek et al. 2005; Renthal and Nestler 2008). The term environment here actually includes a great number of factors from socio-economical factors, to family and peer relations, life events, exposure to pollutants

and medications. These factors participate in shaping the personality and characters of individuals, in determining the socio-cultural context that favors or not drug taking but they also influence the effects of drugs and their ability to interact with neuronal circuits and to produce neuroadaptations. Since evidence that genetic manipulations of the dopaminergic system influence the response to addictive drugs has been reviewed above, in the following section we will focus on the effects on addiction of environmental manipulations.

Environment, DA and vulnerability to addiction

Epidemiological data and experimental data clearly demonstrate that environment plays a major role in the vulnerability to addiction (Renthal and Nestler 2008; Solinas et al. 2010b; Kreek et al. 2005; Stairs and Bardo 2009; Sinha 2007, 2008; Koob 2008). Indeed, people with genetic predisposition will not necessarily develop addiction if they are exposed to positive environments and, in contrast, people with genetic resistance to drugs may develop addiction if they are exposed to negative environments.

In humans, the definition of positive or negative environment can be established “*a posteriori*”. All environmental conditions that increase the risk of addiction are considered negative and, conversely, all environmental conditions that decrease the risk of addiction are considered positive (Jessor and Jessor 1980; Kodjo and Klein 2002). One of the most common factors in negative environmental conditions is the exposure to life stress and trauma that greatly increase the risk to develop addiction (Koob 2008; Piazza and Le Moal 1996; Sinha 2001; De Bellis 2002). In contrast, good family and peer relationships, a stable, understanding and loving entourage but

also physical and mental stimulation decrease the risk of addiction (Jessor and Jessor 1980; Kodjo and Klein 2002; Solinas et al. 2010b; Pang et al. 2018).

In animal models, negative environment is always mimicked by exposure to stress, which may be more or less intense, more or less “early” and more or less chronic (Goeders 2002; Koob 2008; Piazza and Le Moal 1996). In contrast, positive environment is mostly mimicked by environmental enrichment, which is an environment design to facilitate social stimulation, curiosity, cognition and physical exercise (Pang et al. 2018; Solinas et al. 2010b; Stairs and Bardo 2009). Using these two types of manipulation a considerable amount of literature has established that stress indeed increases whereas environmental enrichment reduces the risk to develop and maintain an addiction (Renthal and Nestler 2008; Solinas et al. 2010b; Kreek et al. 2005; Stairs and Bardo 2009; Sinha 2007, 2008; Koob 2008).

Several lines of evidence suggest that the DA system is involved in environment-mediated alteration of vulnerability to addiction (Nader et al. 2008; Goeders 2002; Piazza and Le Moal 1996; Solinas et al. 2010b). For example, a seminal study by the group of Michael Nader showed that, in non-human primates, changes in social ranking influence levels of DA D2Rs in the striatum with dominant individuals showing less D2R than subordinates and self-administering less cocaine (Morgan et al. 2002). In addition, it has been clearly established that stress produces alterations in the DA system that increase the reinforcing effects of drugs (Goeders 2002; Koob 2008; Piazza and Le Moal 1996; Ungless et al. 2010). Conversely, exposure to environmental enrichment produces changes in the DA system that may be related to decreased vulnerability to addiction (Bezard et al. 2003; El Rawas et al. 2009; Kim et al. 2016; Solinas et al. 2010b; Solinas et al. 2009; Thiriet et al. 2008; Lafragette et al. 2017). However, this effect may be mostly related to post-synaptic changes because

drug-induced release of DA in the ventral striatum is not affected by environmental enrichment (El Rawas et al. 2009; Solinas et al. 2009).

Addiction to DA replacement therapy medications in Parkinson's disease

A particular phenomenon that provides support for the role of DA in addictive behavior is the development of addiction to dopaminergic medications in people suffering from Parkinson's disease (PD). PD is characterized by the degeneration of several neuronal ensembles including mesencephalic dopaminergic neurons in the SN and to a lesser extent in the VTA (Chinaglia et al. 1992). DA loss leads to basal ganglia dysfunction and the expression of a parkinsonian syndrome characterized by the variable association of several motor symptoms (akinesia/bradykinesia, rigidity, tremor and postural instability). Management of motor symptoms in PD is achieved with DA replacement therapy (DRT), which includes the DA precursor L-dopa and DA agonists. Chronic intake of DRT can lead to a number of motor and non-motor side effects with severe consequences on the quality of life of the patients and their caregivers (Bastide et al. 2015). Motor side-effects are characterized by several types of motor fluctuations ranging from erratic changes in mobility to abnormal involuntary movements defined as L-dopa induced dyskinesia and that manifest with chorea, dystonia and ballism (for review, see (Bastide et al. 2015)). Non-motor side effects of DRT encompass various abnormal behaviors such as punding or hobbyism (stereotyped/repetitive behaviors), impulse control disorders (behavioral addictions such as pathological gambling, binge eating, hypersexuality or compulsive shopping) as well as the DA dysregulation syndrome (DDS), which is characterized by the compulsive use of DRT (for reviews, see (Voon et al. 2009; Voon et al. 2017)). Compulsive use of DRT is characterized by excessive intake of dopaminergic

medications by the patient, which in turn leads to the emergence or worsening of L-dopa induced dyskinesia, impulse control disorders, as well as hypomania and psychosis (Giovannoni et al. 2000).

Compulsive use of DRT had been occasionally described since the introduction of L-dopa (Nausieda 1985), but more recently it has become the subject of specific attention to precisely define the clinical syndrome, its prevalence, and the profile of affected PD patients (Giovannoni et al. 2000; Pezzella et al. 2005). Giovannoni and colleagues provided a detailed investigation of patients presenting with compulsive use of DRT, and reported that males with a younger age at disease onset were more likely to suffer from DDS (Giovannoni et al. 2000). DDS has an estimated prevalence of 3-4% of PD patients (Giovannoni et al. 2000; Pezzella et al. 2005). Specific diagnosis criteria were also defined based on DSM criteria for substance dependence with additional items specific to PD (Giovannoni et al. 2000). DDS is mainly associated with L-dopa or some DA agonists (essentially apomorphine and bromocriptine). Patients start to self-medicate and rapidly increase their intake of DRT in excess of what would be sufficient to relieve parkinsonism. Patients suffering from DDS display an altered perception of their motor status, feeling relieved from motor symptoms only when they are highly dyskinetic (Giovannoni et al. 2000; Lawrence et al. 2003). Over 90% of PD patients suffering from DDS indeed display drug seeking behavior and escalation of L-dopa intake (Warren et al. 2017). They rapidly consume the amount of prescribed DRT, and then they engage in pathological drug seeking and hoarding to maintain adequate drug supply (Warren et al. 2017). Such massive intake of DRT then leads to the emergence of several behavioral disorders. These include an occurrence of severe dyskinesia, as well as behaviors such as punding (stereotyped, repetitive behavior initially described in

methamphetamine abusers), hobbyism, walkabout, hypersexuality, pathological gambling or shopping (Giovannoni et al. 2000).

Because addiction to DRT shares several features with psychostimulant addiction, this syndrome was discussed in relation with several theories/models of psychostimulant addiction (Lawrence et al. 2003). DDS was initially described as a type of hedonistic homeostatic dysregulation (Giovannoni et al. 2000), with reference to the model proposed by Koob and Le Moal (Koob and Le Moal 1997), where three components (anticipation/preoccupation of drug intake, binge intoxication and negative affect occurring during drug withdrawal) of a spiraling cycle become progressively dysregulated due to sensitization and counter-adaptation processes. In the hedonistic homeostatic dysregulation model, the negative affective state occurring during withdrawal is an important contributor to the addiction cycle and tapering or cessation of L-dopa administration can lead to a negative affective state in over 60% of patients (Warren et al. 2017). Importantly, in the specific context of PD, this negative state is due not only to the symptoms of drug withdrawal but also to the resurgence of the parkinsonian symptoms as the treatment is stopped. However, a withdrawal syndrome has also been described in the tapering of the treatment with DA agonists in PD patients displaying impulse control disorders (Rabinak and Nirenberg 2010), even though these patients do not display compulsive drug intake. DDS can also be viewed as a form of incentive sensitization, according to the theory developed by Robinson and Berridge (Robinson and Berridge 1993) that posits that sensitization of neural circuits involved in the processing of incentive salience results in an increased valence of drug-associated cues and stimuli. In this model, sensitization leads to an increased motivation (wanting) for the drug but not to increased pleasurable effects of the drug (liking) (Robinson and Berridge 1993).

Clinical and neuroimaging evidence suggest that incentive sensitization is involved in DDS since patients report increased “wanting”, but not “liking” the drug, and this “wanting” is correlated with increased DA neurotransmission in the NAc (Evans et al. 2006).

Unlike psychostimulant addiction, which results from the initially voluntary intake of drug in neurologically intact subjects, DDS involves the medical prescription of drugs that act on a brain already displaying severe alterations of DA homeostasis (and other neurotransmitters) due to the degenerative process of PD. How the degenerative process, DRT and specific vulnerability trait interacts in the specific context of DDS remains largely unknown. With regards to the rewarding effects of DRT, L-dopa and DA agonists seem to possess different properties. Administration of L-dopa to healthy subjects is not associated with pleasurable or psychostimulant effects (Liggins et al. 2012), while PD patients (particularly those suffering from DDS) report feelings of pleasure, well-being and a psychostimulant effect when on L-dopa (Castrito et al. 2013), suggesting that dopaminergic neurodegeneration may be required for the expression of psychostimulant-like properties of L-dopa. This hypothesis was confirmed in an animal model of PD where L-dopa displayed rewarding properties and decreased the palatability of a non-drug reward only in rats with a bilateral lesion of the SN (Engeln et al. 2013). Conversely, DA agonists such as apomorphine, bromocriptine or pramipexole display rewarding properties in normal rats (Hoffman et al. 1988; Riddle et al. 2012; van der Kooy et al. 1983). However, the rewarding properties of apomorphine and pramipexole are increased in animals with a lesion of the SN (Campbell et al. 2014; Riddle et al. 2012). Collectively, these studies indicate that the different classes of DRT display specific rewarding properties that are increased following nigral dopaminergic degeneration.

Several neuroadaptations consecutive to the dopaminergic denervation may be involved. These include the supersensitivity of post-synaptic DAR, desensitisation of D2 autoreceptors, as well as overstimulation due to DRT in conjunction with the remaining endogenous dopamine in the brain regions that display low levels of dopaminergic denervation, an effect named “DA overdose”.

Neuroimaging studies provide some support to the DA overdose theory (Cools et al. 2001; Swainson et al. 2000). Using displacement of raclopride binding as an index of L-dopa-induced DA release, PD patients suffering from DDS were found to display an enhanced L-Dopa-induced reduction of raclopride binding in the ventral striatum, thereby suggesting a sensitization of DA transmission in this striatal subregion (Evans et al. 2006). Interestingly, a similar sensitization was also found in the putamen of PD patients displaying L-dopa induced dyskinesia (de la Fuente-Fernandez et al. 2000; de la Fuente-Fernandez et al. 2004), suggesting that both motor and non-motor side effects of L-dopa may share some common neurobiological mechanisms. Whether the altered synaptic plasticity and associated dysregulation of intracellular signaling occurring in L-dopa induced dyskinesia are recapitulated in DDS remains to be determined.

DA and the treatment of addiction

Given the vast literature about the critical role of DA in drug addiction, as it should be expected, extensive research has been conducted on the use of dopaminergic compounds for the treatment of drug addiction. For example, blocking DA neurotransmission was expected to block the reinforcing effects of drugs and therefore, to reduce addiction-related behavior. However, so far accumulating evidence suggests that dopaminergic antagonists are not effective to treat drug addiction (Alvarez et al. 2013; Verrico et al. 2013). The reasons for this partial failure

are multiple. First of all, although DA appears to play an important role in addiction to all drugs, it is clear that its role is much more critical for psychostimulants whereas the role in opiate, nicotine, alcohol and cannabis abuse is much less central (Pierce and Kumaresan 2006; Nutt et al. 2015). A second aspect that should be taken into consideration is that pharmacological blockade could be overwhelmed by increasing the amount of drug consumed. Indeed, studies in animals and humans show that, under most conditions, blockade of DAR increases drug self-administration as individuals try to compensate for the reduction of the effects of unitary doses (Arnold and Roberts 1997; Koob et al. 1997; Mello and Negus 1996) and similar results have been found in humans (Haney et al. 2001). Thus, in naturalistic settings, humans would have the possibility to counteract the effects of DA blocked by increasing the dose of the drug. Thirdly, whereas DA system is arguably the major actor in reward process, reward processes only constitute a part of the addiction pathology (Koob and Volkow 2010, 2016; Volkow et al. 2016). Indeed, as discussed earlier positive reinforcement is critical for the development of addiction but upon repeated administration, conditioned effects of drugs and habits start playing a major role (Everitt and Robbins 2005; Koob and Volkow 2010, 2016; Volkow et al. 2016). Although DA also plays role in conditioned effects and habits, these phenomena appear to be much more resistant to DA blockade than simple reinforcement (Everitt et al. 2008; Everitt and Robbins 2005). Fourthly, the cyclic nature of addiction pathology implies that phase of active self-administration and elevated DA levels are alternated with phase of abstinence and lower levels of DA that may be associated with anhedonia and craving (Koob and Volkow 2010, 2016; Volkow et al. 2016). Therefore, DA antagonist may well exacerbate negative affective states and increase negative reinforcement, which could lead to low tolerance and compliance for the

medication, cessation of the treatment and finally relapse (Dackis and Gold 1985; Koob and Volkow 2010, 2016; Kuhar and Pilotte 1996). Indeed, most medication with selective blockade of DA receptors used in clinical settings, which are mostly D2 antagonists, are known to produce aversive psychological effects that reduce their compliance (Alvarez et al. 2013). With this in mind, antipsychotic medications with mixed pharmacological profile comprising both D2 blockade and effects on other targets such as serotonin 5-HT₂ receptors (Alvarez et al. 2013). However, a meta-analysis of the studies using these compounds has also concluded that their effectiveness in the treatment of addiction is very limited (Alvarez et al. 2013). Importantly, the use of selective D1-receptor antagonists for clinical applications has been hampered by issues with bioavailability, binding affinity, pharmacological kinetics, and side effects but new pharmacological compounds may provide novel tools for the pharmacotherapy of addiction (Kim et al. 2015). The use of the selective D1 antagonist, ecopipam, has been tested on self-administration of cocaine, in humans but it was found that its administration increased self-administration of cocaine while decreasing cocaine craving (but in the same patients craving for alcohol and tobacco smoking increased) suggesting that it is unlikely to be an effective treatment for cocaine addiction (Haney et al. 2001). Some authors have proposed that signaling of D1R may be modulated by interacting targeting second messengers, allosteric modulators, or by making targeted modifications to D1R machinery (Kim et al. 2015). However, lessons from the use of naltrexone for opiate addiction suggest that therapy with antagonists may be effective only in a limited number of circumstances and mostly in very motivated people receiving long periods of high level of counseling (Blanco-Gandia and Rodriguez-Arias 2018). It should be noted that because of their specific neurobiological characteristics, several authors

have proposed targeting the D3R with antagonists and partial to better management of addiction (Newman et al. 2012; Le Foll et al. 2014). However, no conclusive evidence has been provided by clinical trials (Le Foll et al. 2014)

The case for the use of dopaminergic agonists in the management of addiction to psychostimulants is much stronger (Herin et al. 2010) and it is supported by parallels in opiate and nicotine addiction in which substitution therapy with opioid agonists such as methadone and buprenorphine (Dematteis et al. 2017) and nicotine replacement therapy (Haustein 2000; Etter and Stapleton 2006) have been proven to be relatively effective. Direct agonist of dopaminergic receptors do not appear to present the best pharmacological profile, do not appear to stimulate the reward system strongly enough and may produce unwanted effects such as impulsivity (Kim et al. 2015). Thus, most interest has been given to the investigation of indirect DA agonists that act on dopaminergic (and other monoaminergic) transporters. Among these drugs modafinil, bupropion, methylphenidate and slow release formulations of amphetamine and methamphetamine, that are used in humans for the treatment of other psychiatric disorders, have been tested in humans and although some studies have found some beneficial effects of these compounds the overall effects have been limited (Herin et al. 2010; Verrico et al. 2013). New avenues for this therapeutic strategy may arise from the development of new synthetic blockers of DAT that bind to the transporter without producing the psychotropic and reinforcing of cocaine and amphetamine (Rothman et al. 2008).

It should be finally mentioned that other indirect strategies to boosting dopaminergic activities such as decrease metabolism by the MAO antagonist selegiline or increase

in DA precursors such as administration of DOPA and carbi-DOPA have been tested but have shown limited success (Herin et al. 2010; Verrico et al. 2013).

Altogether these studies show that notwithstanding the high expectations on the possibility to target the DA system for the treatment of addiction, until now the results have not met these expectations (Nutt et al. 2015). However, the quest for new effective DA medications continues by refining of the elements in the DA system to target and the specific conditions in which these medications may be effective.

Conclusions

We know a lot, arguably more than any other neurotransmitter, about the role of DA in addiction. Although this knowledge has not translated in medications as it could have been hoped (Nutt et al. 2015), the experience with DA may be an exemplary showcase of the role of research (preclinical and clinical) in shedding light into brain functioning. In the attempt to understand drug addiction, not only we have produced considerable advances in the understanding of how brain DA functions in ways that are helpful for different neurological and psychiatric disorders, but this knowledge has also been used to predict vulnerability, effectiveness of the treatment and recovery from addiction. More importantly, the failure of some of the predictions of the original DA-driven hypotheses have allowed the field to re-conceptualize addiction as a complex disorder with different aspects and different phases (Koob and Volkow 2010), to consider that different drugs may be associated with different addiction-related behaviors (Badiani et al. 2011) and to question the validity of established animal models addiction and their ability to predict effectiveness of new medications in humans and to develop new ones (Pierce et al. 2012). Finally, the humbling reality of the difficulties in translating the best preclinical and clinical knowledge into new effective treatments should teach researchers and the general public to accept that

the path from research to real-life applications is arduous, unpredictable and slow and that even the most promising discoveries may need considerable time and energy to lead to meaningful clinical results.

REFERENCES

- Ackerman JM, White FJ (1990) A10 somatodendritic dopamine autoreceptor sensitivity following withdrawal from repeated cocaine treatment. *Neurosci Lett* 117 (1-2):181-187
- Acquas E, Carboni E, Di Chiara G (1991) Profound depression of mesolimbic dopamine release after morphine withdrawal in dependent rats. *Eur J Pharmacol* 193 (1):133-134
- Adermark L, Morud J, Lotfi A, Ericson M, Soderpalm B (2018) Acute and chronic modulation of striatal endocannabinoid-mediated plasticity by nicotine. *Addict Biol*. doi:10.1111/adb.12598
- Ahmed SH, Koob GF (1998) Transition from moderate to excessive drug intake: change in hedonic set point. *Science* 282 (5387):298-300
- Alibhai IN, Green TA, Potashkin JA, Nestler EJ (2007) Regulation of fosB and DeltafosB mRNA expression: in vivo and in vitro studies. *Brain Res* 1143:22-33. doi:10.1016/j.brainres.2007.01.069
- Alvarez Y, Perez-Mana C, Torrens M, Farre M (2013) Antipsychotic drugs in cocaine dependence: a systematic review and meta-analysis. *J Subst Abuse Treat* 45 (1):1-10. doi:10.1016/j.jsat.2012.12.013
- Andreoli M, Tessari M, Pilla M, Valerio E, Hagan JJ, Heidbreder CA (2003) Selective antagonism at dopamine D3 receptors prevents nicotine-triggered relapse to nicotine-seeking behavior. *Neuropsychopharmacology* 28 (7):1272-1280. doi:10.1038/sj.npp.1300183
- Argilli E, Sibley DR, Malenka RC, England PM, Bonci A (2008) Mechanism and time course of cocaine-induced long-term potentiation in the ventral tegmental area. *J Neurosci* 28 (37):9092-9100. doi:10.1523/JNEUROSCI.1001-08.2008
- Arnold JM, Roberts DC (1997) A critique of fixed and progressive ratio schedules used to examine the neural substrates of drug reinforcement. *Pharmacol Biochem Behav* 57 (3):441-447
- Arroyo M, Baker WA, Everitt BJ (2000) Cocaine self-administration in rats differentially alters mRNA levels of the monoamine transporters and striatal neuropeptides. *Brain Res Mol Brain Res* 83 (1-2):107-120
- Atwood BK, Kupferschmidt DA, Lovinger DM (2014) Opioids induce dissociable forms of long-term depression of excitatory inputs to the dorsal striatum. *Nat Neurosci* 17 (4):540-548. doi:10.1038/nn.3652
- Bachtell RK, Whisler K, Karanian D, Self DW (2005) Effects of intra-nucleus accumbens shell administration of dopamine agonists and antagonists on cocaine-taking and cocaine-seeking behaviors in the rat. *Psychopharmacology (Berl)* 183 (1):41-53. doi:10.1007/s00213-005-0133-1
- Badiani A, Belin D, Epstein D, Calu D, Shaham Y (2011) Opiate versus psychostimulant addiction: the differences do matter. *Nat Rev Neurosci* 12 (11):685-700. doi:10.1038/nrn3104
- Baik JH, Picetti R, Saiardi A, Thiriet G, Dierich A, Depaulis A, Le Meur M, Borrelli E (1995) Parkinsonian-like locomotor impairment in mice lacking dopamine D2 receptors. *Nature* 377 (6548):424-428
- Ball KT, Wellman CL, Fortenberry E, Rebec GV (2009) Sensitizing regimens of (+/-)-3, 4-methylenedioxymethamphetamine (ecstasy) elicit enduring and differential structural alterations in the brain motive circuit of the rat. *Neuroscience* 160 (2):264-274. doi:10.1016/j.neuroscience.2009.02.025

- Barrett SP, Boileau I, Okker J, Pihl RO, Dagher A (2004) The hedonic response to cigarette smoking is proportional to dopamine release in the human striatum as measured by positron emission tomography and [¹¹C]raclopride. *Synapse* 54 (2):65-71. doi:10.1002/syn.20066
- Bastide MF, Meissner WG, Picconi B, Fasano S, Fernagut PO, Feyder M, Francardo V, Alcacer C, Ding Y, Brambilla R, Fisone G, Jon Stoessl A, Bourdenx M, Engeln M, Navailles S, De Deurwaerdere P, Ko WK, Simola N, Morelli M, Groc L, Rodriguez MC, Gurevich EV, Quik M, Morari M, Mellone M, Gardoni F, Tronci E, Guehl D, Tison F, Crossman AR, Kang UJ, Steece-Collier K, Fox S, Carta M, Angela Cenci M, Bezard E (2015) Pathophysiology of L-dopa-induced motor and non-motor complications in Parkinson's disease. *Prog Neurobiol* 132:96-168. doi:10.1016/j.pneurobio.2015.07.002
- Beaudoin GMJ, 3rd, Gomez JA, Perkins J, Bland JL, Petko AK, Paladini CA (2018) Cocaine Selectively Reorganizes Excitatory Inputs to Substantia Nigra Pars Compacta Dopamine Neurons. *J Neurosci* 38 (5):1151-1159. doi:10.1523/JNEUROSCI.1975-17.2017
- Beckmann AM, Wilce PA (1997) Egr transcription factors in the nervous system. *Neurochem Int* 31 (4):477-510; discussion 517-476
- Belin D, Everitt BJ (2008) Cocaine seeking habits depend upon dopamine-dependent serial connectivity linking the ventral with the dorsal striatum. *Neuron* 57 (3):432-441. doi:10.1016/j.neuron.2007.12.019
- Bellone C, Luscher C (2006) Cocaine triggered AMPA receptor redistribution is reversed in vivo by mGluR-dependent long-term depression. *Nat Neurosci* 9 (5):636-641. doi:10.1038/nn1682
- Belujon P, Grace AA (2014) Restoring mood balance in depression: ketamine reverses deficit in dopamine-dependent synaptic plasticity. *Biol Psychiatry* 76 (12):927-936. doi:10.1016/j.biopsych.2014.04.014
- Belujon P, Jakobowski NL, Dollish HK, Grace AA (2016) Withdrawal from Acute Amphetamine Induces an Amygdala-Driven Attenuation of Dopamine Neuron Activity: Reversal by Ketamine. *Neuropsychopharmacology* 41 (2):619-627. doi:10.1038/npp.2015.191
- Ben-Shahar O, Keeley P, Cook M, Brake W, Joyce M, Nyffeler M, Heston R, Ettenberg A (2007) Changes in levels of D1, D2, or NMDA receptors during withdrawal from brief or extended daily access to IV cocaine. *Brain Res* 1131 (1):220-228. doi:10.1016/j.brainres.2006.10.069
- Ben-Shahar O, Moscarello JM, Ettenberg A (2006) One hour, but not six hours, of daily access to self-administered cocaine results in elevated levels of the dopamine transporter. *Brain Res* 1095 (1):148-153. doi:10.1016/j.brainres.2006.04.002
- Benoit-Marand M, Ballion B, Borrelli E, Boraud T, Gonon F (2011) Inhibition of dopamine uptake by D2 antagonists: an in vivo study. *J Neurochem* 116 (3):449-458. doi:10.1111/j.1471-4159.2010.07125.x
- Benoit-Marand M, Borrelli E, Gonon F (2001) Inhibition of dopamine release via presynaptic D2 receptors: time course and functional characteristics in vivo. *J Neurosci* 21 (23):9134-9141
- Benoit-Marand M, Jaber M, Gonon F (2000a) Release and elimination of dopamine in vivo in mice lacking the dopamine transporter: functional consequences. *Eur J Neurosci* 12 (8):2985-2992

- Benoit-Marand M, Jaber M, Gonon F (2000b) Release and elimination of dopamine in vivo in mice lacking the dopamine transporter: functional consequences. *Eur J Neurosci* 12 (8):2985-2992
- Berridge KC, Robinson TE (1998) What is the role of dopamine in reward: hedonic impact, reward learning, or incentive salience? *Brain Res Brain Res Rev* 28 (3):309-369
- Berry KP, Nedivi E (2017) Spine Dynamics: Are They All the Same? *Neuron* 96 (1):43-55. doi:10.1016/j.neuron.2017.08.008
- Bertran-Gonzalez J, Bosch C, Maroteaux M, Matamales M, Herve D, Valjent E, Girault JA (2008) Opposing patterns of signaling activation in dopamine D1 and D2 receptor-expressing striatal neurons in response to cocaine and haloperidol. *J Neurosci* 28 (22):5671-5685. doi:10.1523/JNEUROSCI.1039-08.2008
- Beveridge TJ, Smith HR, Nader MA, Porrino LJ (2009) Abstinence from chronic cocaine self-administration alters striatal dopamine systems in rhesus monkeys. *Neuropsychopharmacology* 34 (5):1162-1171. doi:10.1038/npp.2008.135
- Bezard E, Dovero S, Belin D, Duconger S, Jackson-Lewis V, Przedborski S, Piazza PV, Gross CE, Jaber M (2003) Enriched environment confers resistance to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine and cocaine: involvement of dopamine transporter and trophic factors. *J Neurosci* 23 (35):10999-11007
- Blanco-Gandia MC, Rodriguez-Arias M (2018) Pharmacological treatments for opiate and alcohol addiction: A historical perspective of the last 50 years. *Eur J Pharmacol* 836:89-101. doi:10.1016/j.ejphar.2018.08.007
- Blomeley CP, Cains S, Smith R, Bracci E (2011) Ethanol affects striatal interneurons directly and projection neurons through a reduction in cholinergic tone. *Neuropsychopharmacology* 36 (5):1033-1046. doi:10.1038/npp.2010.241
- Boileau I, Assaad JM, Pihl RO, Benkelfat C, Leyton M, Diksic M, Tremblay RE, Dagher A (2003) Alcohol promotes dopamine release in the human nucleus accumbens. *Synapse* 49 (4):226-231. doi:10.1002/syn.10226
- Bonci A, Malenka RC (1999) Properties and plasticity of excitatory synapses on dopaminergic and GABAergic cells in the ventral tegmental area. *J Neurosci* 19 (10):3723-3730
- Bontempi B, Sharp FR (1997) Systemic morphine-induced Fos protein in the rat striatum and nucleus accumbens is regulated by mu opioid receptors in the substantia nigra and ventral tegmental area. *J Neurosci* 17 (21):8596-8612
- Borgland SL, Malenka RC, Bonci A (2004) Acute and chronic cocaine-induced potentiation of synaptic strength in the ventral tegmental area: electrophysiological and behavioral correlates in individual rats. *J Neurosci* 24 (34):7482-7490. doi:10.1523/JNEUROSCI.1312-04.2004
- Bortz DM, Grace AA (2018) Medial septum differentially regulates dopamine neuron activity in the rat ventral tegmental area and substantia nigra via distinct pathways. *Neuropsychopharmacology* 43 (10):2093-2100. doi:10.1038/s41386-018-0048-2
- Bosse R, Fumagalli F, Jaber M, Giros B, Gainetdinov RR, Wetsel WC, Missale C, Caron MG (1997) Anterior pituitary hypoplasia and dwarfism in mice lacking the dopamine transporter. *Neuron* 19 (1):127-138
- Bossong MG, van Berckel BN, Boellaard R, Zuurman L, Schuit RC, Windhorst AD, van Gerven JM, Ramsey NF, Lammertsma AA, Kahn RS (2009) Delta 9-tetrahydrocannabinol induces dopamine release in the human striatum. *Neuropsychopharmacology* 34 (3):759-766. doi:10.1038/npp.2008.138

- Boudreau AC, Wolf ME (2005) Behavioral sensitization to cocaine is associated with increased AMPA receptor surface expression in the nucleus accumbens. *J Neurosci* 25 (40):9144-9151. doi:10.1523/JNEUROSCI.2252-05.2005
- Bourdy R, Barrot M (2012) A new control center for dopaminergic systems: pulling the VTA by the tail. *Trends Neurosci* 35 (11):681-690. doi:10.1016/j.tins.2012.06.007
- Bourdy R, Sanchez-Catalan MJ, Kaufling J, Balcita-Pedicino JJ, Freund-Mercier MJ, Veinante P, Sesack SR, Georges F, Barrot M (2014) Control of the nigrostriatal dopamine neuron activity and motor function by the tail of the ventral tegmental area. *Neuropsychopharmacology* 39 (12):2788-2798. doi:10.1038/npp.2014.129
- Boyce-Rustay JM, Risinger FO (2003) Dopamine D3 receptor knockout mice and the motivational effects of ethanol. *Pharmacol Biochem Behav* 75 (2):373-379
- Brady AM, Glick SD, O'Donnell P (2005) Selective disruption of nucleus accumbens gating mechanisms in rats behaviorally sensitized to methamphetamine. *J Neurosci* 25 (28):6687-6695. doi:10.1523/JNEUROSCI.0643-05.2005
- Briand LA, Blendy JA (2010) Molecular and genetic substrates linking stress and addiction. *Brain Res* 1314:219-234. doi:10.1016/j.brainres.2009.11.002
- Britt JP, Benaliouad F, McDevitt RA, Stuber GD, Wise RA, Bonci A (2012) Synaptic and behavioral profile of multiple glutamatergic inputs to the nucleus accumbens. *Neuron* 76 (4):790-803. doi:10.1016/j.neuron.2012.09.040
- Brodie MS (2002) Increased ethanol excitation of dopaminergic neurons of the ventral tegmental area after chronic ethanol treatment. *Alcohol Clin Exp Res* 26 (7):1024-1030. doi:10.1097/01.ALC.0000021336.33310.6B
- Brodie MS, Pesold C, Appel SB (1999) Ethanol directly excites dopaminergic ventral tegmental area reward neurons. *Alcohol Clin Exp Res* 23 (11):1848-1852
- Brody AL, Olmstead RE, London ED, Farahi J, Meyer JH, Grossman P, Lee GS, Huang J, Hahn EL, Mandelkern MA (2004) Smoking-induced ventral striatum dopamine release. *Am J Psychiatry* 161 (7):1211-1218. doi:10.1176/appi.ajp.161.7.1211
- Brown MT, Bellone C, Mameli M, Labouebe G, Bocklisch C, Balland B, Dahan L, Lujan R, Deisseroth K, Luscher C (2010) Drug-driven AMPA receptor redistribution mimicked by selective dopamine neuron stimulation. *PloS one* 5 (12):e15870. doi:10.1371/journal.pone.0015870
- Brown PL, Palacorolla H, Brady D, Riegger K, Elmer GI, Shepard PD (2017) Habenula-Induced Inhibition of Midbrain Dopamine Neurons Is Diminished by Lesions of the Rostromedial Tegmental Nucleus. *J Neurosci* 37 (1):217-225. doi:10.1523/JNEUROSCI.1353-16.2016
- Brown RW, Kolb B (2001) Nicotine sensitization increases dendritic length and spine density in the nucleus accumbens and cingulate cortex. *Brain Res* 899 (1-2):94-100
- Brown TE, Lee BR, Mu P, Ferguson D, Dietz D, Ohnishi YN, Lin Y, Suska A, Ishikawa M, Huang YH, Shen H, Kalivas PW, Sorg BA, Zukin RS, Nestler EJ, Dong Y, Schluter OM (2011) A silent synapse-based mechanism for cocaine-induced locomotor sensitization. *J Neurosci* 31 (22):8163-8174. doi:10.1523/JNEUROSCI.0016-11.2011
- Budygin EA, Brodie MS, Sotnikova TD, Mateo Y, John CE, Cyr M, Gainetdinov RR, Jones SR (2004) Dissociation of rewarding and dopamine transporter-mediated properties of amphetamine. *Proc Natl Acad Sci U S A* 101 (20):7781-7786. doi:10.1073/pnas.0401418101

- Bunney BS, Aghajanian GK, Roth RH (1973) Comparison of effects of L-dopa, amphetamine and apomorphine on firing rate of rat dopaminergic neurones. *Nat New Biol* 245 (143):123-125
- Cagniard B, Balsam PD, Brunner D, Zhuang X (2006) Mice with chronically elevated dopamine exhibit enhanced motivation, but not learning, for a food reward. *Neuropsychopharmacology* 31 (7):1362-1370. doi:10.1038/sj.npp.1300966
- Cagniard B, Sotnikova TD, Gainetdinov RR, Zhuang X (2014) The dopamine transporter expression level differentially affects responses to cocaine and amphetamine. *J Neurogenet* 28 (1-2):112-121. doi:10.3109/01677063.2014.908191
- Caine SB, Negus SS, Mello NK, Patel S, Bristow L, Kulagowski J, Vallone D, Saiardi A, Borrelli E (2002) Role of dopamine D2-like receptors in cocaine self-administration: studies with D2 receptor mutant mice and novel D2 receptor antagonists. *J Neurosci* 22 (7):2977-2988. doi:20026264
- Caine SB, Thomsen M, Gabriel KI, Berkowitz JS, Gold LH, Koob GF, Tonegawa S, Zhang J, Xu M (2007) Lack of self-administration of cocaine in dopamine D1 receptor knock-out mice. *J Neurosci* 27 (48):13140-13150. doi:10.1523/JNEUROSCI.2284-07.2007
- Calipari ES, Beveridge TJ, Jones SR, Porrino LJ (2013) Withdrawal from extended-access cocaine self-administration results in dysregulated functional activity and altered locomotor activity in rats. *Eur J Neurosci* 38 (12):3749-3757. doi:10.1111/ejn.12381
- Campbell JC, Jeyamohan SB, De La Cruz P, Chen N, Shin D, Pilitsis JG (2014) Place conditioning to apomorphine in rat models of Parkinson's disease: differences by dose and side-effect expression. *Behav Brain Res* 275:114-119. doi:10.1016/j.bbr.2014.09.002
- Caprioli D, Fryer TD, Sawiak SJ, Aigbirhio FI, Dalley JW (2013) Translating positron emission tomography studies in animals to stimulant addiction: promises and pitfalls. *Curr Opin Neurobiol* 23 (4):597-606. doi:10.1016/j.conb.2013.04.003
- Carboni E, Spielesoy C, Vacca C, Nosten-Bertrand M, Giros B, Di Chiara G (2001) Cocaine and amphetamine increase extracellular dopamine in the nucleus accumbens of mice lacking the dopamine transporter gene. *J Neurosci* 21 (9):RC141: 141-144
- Carlezon WA, Jr., Duman RS, Nestler EJ (2005) The many faces of CREB. *Trends Neurosci* 28 (8):436-445. doi:10.1016/j.tins.2005.06.005
- Carlsson A, Lindqvist M (1963) Effect of Chlorpromazine or Haloperidol on Formation of 3-methoxytyramine and Normetanephrine in Mouse Brain. *Acta Pharmacol Toxicol (Copenh)* 20:140-144
- Carlsson A, Lindqvist M, Magnusson T (1957) 3,4-Dihydroxyphenylalanine and 5-hydroxytryptophan as reserpine antagonists. *Nature* 180 (4596):1200
- Castrioto A, Kistner A, Klinger H, Lhomme E, Schmitt E, Fraix V, Chabardes S, Mertens P, Quesada JL, Broussolle E, Pollak P, Thobois SC, Krack P (2013) Psychostimulant effect of levodopa: reversing sensitisation is possible. *Journal of neurology, neurosurgery, and psychiatry* 84 (1):18-22. doi:10.1136/jnnp-2012-302444
- Celada P, Paladini CA, Tepper JM (1999) GABAergic control of rat substantia nigra dopaminergic neurons: role of globus pallidus and substantia nigra pars reticulata. *Neuroscience* 89 (3):813-825
- Cerruti C, Pilote NS, Uhl G, Kuhar MJ (1994) Reduction in dopamine transporter mRNA after cessation of repeated cocaine administration. *Brain Res Mol Brain Res* 22 (1-4):132-138

- Chang CH, Grace AA (2014) Amygdala-ventral pallidum pathway decreases dopamine activity after chronic mild stress in rats. *Biol Psychiatry* 76 (3):223-230. doi:10.1016/j.biopsych.2013.09.020
- Chang HT, Kitai ST (1985) Projection neurons of the nucleus accumbens: an intracellular labeling study. *Brain Res* 347 (1):112-116
- Chang L, Alicata D, Ernst T, Volkow N (2007) Structural and metabolic brain changes in the striatum associated with methamphetamine abuse. *Addiction* 102 Suppl 1:16-32. doi:10.1111/j.1360-0443.2006.01782.x
- Chen BT, Bowers MS, Martin M, Hopf FW, Guillory AM, Carelli RM, Chou JK, Bonci A (2008) Cocaine but not natural reward self-administration nor passive cocaine infusion produces persistent LTP in the VTA. *Neuron* 59 (2):288-297
- Chen J, Kelz MB, Hope BT, Nakabeppu Y, Nestler EJ (1997) Chronic Fos-related antigens: stable variants of deltaFosB induced in brain by chronic treatments. *J Neurosci* 17 (13):4933-4941
- Chen NH, Reith ME (1994) Autoregulation and monoamine interactions in the ventral tegmental area in the absence and presence of cocaine: a microdialysis study in freely moving rats. *J Pharmacol Exp Ther* 271 (3):1597-1610
- Chen R, Tilley MR, Wei H, Zhou F, Zhou FM, Ching S, Quan N, Stephens RL, Hill ER, Nottoli T, Han DD, Gu HH (2006) Abolished cocaine reward in mice with a cocaine-insensitive dopamine transporter. *Proc Natl Acad Sci U S A* 103 (24):9333-9338. doi:10.1073/pnas.0600905103
- Chergui K, Akaoka H, Charley PJ, Saunier CF, Buda M, Chouvet G (1994) Subthalamic nucleus modulates burst firing of nigral dopamine neurones via NMDA receptors. *Neuroreport* 5 (10):1185-1188
- Chinaglia G, Alvarez FJ, Probst A, Palacios JM (1992) Mesostriatal and mesolimbic dopamine uptake binding sites are reduced in Parkinson's disease and progressive supranuclear palsy: a quantitative autoradiographic study using [³H]mazindol. *Neuroscience* 49 (2):317-327
- Christoph GR, Leonzio RJ, Wilcox KS (1986) Stimulation of the lateral habenula inhibits dopamine-containing neurons in the substantia nigra and ventral tegmental area of the rat. *J Neurosci* 6 (3):613-619
- Ciliax BJ, Heilman C, Demchyshyn LL, Pristupa ZB, Ince E, Hersch SM, Niznik HB, Levey AI (1995) The dopamine transporter: immunochemical characterization and localization in brain. *J Neurosci* 15 (3 Pt 1):1714-1723
- Coffey KR, Barker DJ, Gayliard N, Kulik JM, Pawlak AP, Stamos JP, West MO (2015) Electrophysiological evidence of alterations to the nucleus accumbens and dorsolateral striatum during chronic cocaine self-administration. *Eur J Neurosci* 41 (12):1538-1552. doi:10.1111/ejn.12904
- Colby CR, Whisler K, Steffen C, Nestler EJ, Self DW (2003) Striatal cell type-specific overexpression of DeltaFosB enhances incentive for cocaine. *J Neurosci* 23 (6):2488-2493
- Conrad LC, Pfaff DW (1976) Autoradiographic tracing of nucleus accumbens efferents in the rat. *Brain Res* 113 (3):589-596
- Cools R, Barker RA, Sahakian BJ, Robbins TW (2001) Enhanced or impaired cognitive function in Parkinson's disease as a function of dopaminergic medication and task demands. *Cereb Cortex* 11 (12):1136-1143
- Cowen MS, Lawrence AJ (2001) Alterations in central preproenkephalin mRNA expression after chronic free-choice ethanol consumption by fawn-hooded rats. *Alcohol Clin Exp Res* 25 (8):1126-1133

- Cox SML, Yau Y, Larcher K, Durand F, Kolivakis T, Delaney JS, Dagher A, Benkelfat C, Leyton M (2017) Cocaine Cue-Induced Dopamine Release in Recreational Cocaine Users. *Sci Rep* 7:46665. doi:10.1038/srep46665
- Crippens D, Robinson TE (1994) Withdrawal from morphine or amphetamine: different effects on dopamine in the ventral-medial striatum studied with microdialysis. *Brain Res* 650 (1):56-62
- Crombag HS, Bossert JM, Koya E, Shaham Y (2008) Review. Context-induced relapse to drug seeking: a review. *Philos Trans R Soc Lond B Biol Sci* 363 (1507):3233-3243. doi:10.1098/rstb.2008.0090
- Cuzon Carlson VC, Seabold GK, Helms CM, Garg N, Odagiri M, Rau AR, Daunais J, Alvarez VA, Lovinger DM, Grant KA (2011) Synaptic and morphological neuroadaptations in the putamen associated with long-term, relapsing alcohol drinking in primates. *Neuropsychopharmacology* 36 (12):2513-2528. doi:10.1038/npp.2011.140
- Dackis CA, Gold MS (1985) New concepts in cocaine addiction: the dopamine depletion hypothesis. *Neurosci Biobehav Rev* 9 (3):469-477
- Dalley JW, Fryer TD, Brichard L, Robinson ES, Theobald DE, Laane K, Pena Y, Murphy ER, Shah Y, Probst K, Abakumova I, Aigbirhio FI, Richards HK, Hong Y, Baron JC, Everitt BJ, Robbins TW (2007) Nucleus accumbens D2/3 receptors predict trait impulsivity and cocaine reinforcement. *Science* 315 (5816):1267-1270. doi:10.1126/science.1137073
- Damsma G, Pfaus JG, Wenkstern D, Phillips AG, Fibiger HC (1992) Sexual behavior increases dopamine transmission in the nucleus accumbens and striatum of male rats: comparison with novelty and locomotion. *Behav Neurosci* 106 (1):181-191
- De Bellis MD (2002) Developmental traumatology: a contributory mechanism for alcohol and substance use disorders. *Psychoneuroendocrinology* 27 (1-2):155-170
- de la Fuente-Fernandez R, Pal PK, Vingerhoets FJ, Kishore A, Schulzer M, Mak EK, Ruth TJ, Snow BJ, Calne DB, Stoessl AJ (2000) Evidence for impaired presynaptic dopamine function in parkinsonian patients with motor fluctuations. *J Neural Transm* 107 (1):49-57
- de la Fuente-Fernandez R, Sossi V, Huang Z, Furtado S, Lu JQ, Calne DB, Ruth TJ, Stoessl AJ (2004) Levodopa-induced changes in synaptic dopamine levels increase with progression of Parkinson's disease: implications for dyskinesias. *Brain* 127 (Pt 12):2747-2754
- Deister CA, Teagarden MA, Wilson CJ, Paladini CA (2009) An intrinsic neuronal oscillator underlies dopaminergic neuron bursting. *J Neurosci* 29 (50):15888-15897. doi:10.1523/JNEUROSCI.4053-09.2009
- Dematteis M, Auriacombe M, D'Agnone O, Somaini L, Szerman N, Littlewood R, Alam F, Alho H, Benyamina A, Bobes J, Daulouede JP, Leonardi C, Maremmanni I, Torrens M, Walcher S, Soyka M (2017) Recommendations for buprenorphine and methadone therapy in opioid use disorder: a European consensus. *Expert Opin Pharmacother* 18 (18):1987-1999. doi:10.1080/14656566.2017.1409722
- Deroche-Gamonet V, Belin D, Piazza PV (2004) Evidence for addiction-like behavior in the rat. *Science* 305 (5686):1014-1017
- Deroche-Gamonet V, Piazza PV (2014) Psychobiology of cocaine addiction: Contribution of a multi-symptomatic animal model of loss of control. *Neuropharmacology* 76 Pt B:437-449. doi:10.1016/j.neuropharm.2013.07.014
- Di Chiara G, Bassareo V (2007) Reward system and addiction: what dopamine does and doesn't do. *Curr Opin Pharmacol* 7 (1):69-76. doi:10.1016/j.coph.2006.11.003

- Di Chiara G, Imperato A (1988) Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci U S A* 85 (14):5274-5278
- Di Chiara G, Tanda G, Bassareo V, Pontieri F, Acquas E, Fenu S, Cadoni C, Carboni E (1999) Drug addiction as a disorder of associative learning. Role of nucleus accumbens shell/extended amygdala dopamine. *Ann N Y Acad Sci* 877:461-485
- Diana M (1998) Drugs of abuse and dopamine cell activity. *Adv Pharmacol* 42:998-1001
- Diana M, Gessa GL, Rossetti ZL (1992a) Lack of tolerance to ethanol-induced stimulation of mesolimbic dopamine system. *Alcohol Alcohol* 27 (4):329-333
- Diana M, Melis M, Muntoni AL, Gessa GL (1998) Mesolimbic dopaminergic decline after cannabinoid withdrawal. *Proc Natl Acad Sci U S A* 95 (17):10269-10273
- Diana M, Pistis M, Muntoni A, Gessa G (1995) Profound decrease of mesolimbic dopaminergic neuronal activity in morphine withdrawn rats. *J Pharmacol Exp Ther* 272 (2):781-785
- Diana M, Pistis M, Muntoni A, Rossetti ZL, Gessa G (1992b) Marked decrease of A10 dopamine neuronal firing during ethanol withdrawal syndrome in rats. *Eur J Pharmacol* 221 (2-3):403-404
- Dong Y, Taylor JR, Wolf ME, Shaham Y (2017) Circuit and Synaptic Plasticity Mechanisms of Drug Relapse. *J Neurosci* 37 (45):10867-10876. doi:10.1523/JNEUROSCI.1821-17.2017
- Dos Santos M, Cahill EN, Bo GD, Vanhoutte P, Caboche J, Giros B, Heck N (2018) Cocaine increases dopaminergic connectivity in the nucleus accumbens. *Brain Struct Funct* 223 (2):913-923. doi:10.1007/s00429-017-1532-x
- Dos Santos M, Salery M, Forget B, Garcia Perez MA, Betuing S, Boudier T, Vanhoutte P, Caboche J, Heck N (2017) Rapid Synaptogenesis in the Nucleus Accumbens Is Induced by a Single Cocaine Administration and Stabilized by Mitogen-Activated Protein Kinase Interacting Kinase-1 Activity. *Biol Psychiatry* 82 (11):806-818. doi:10.1016/j.biopsych.2017.03.014
- Drevets WC, Gautier C, Price JC, Kupfer DJ, Kinahan PE, Grace AA, Price JL, Mathis CA (2001) Amphetamine-induced dopamine release in human ventral striatum correlates with euphoria. *Biol Psychiatry* 49 (2):81-96
- DSM-5 (2013) Diagnostic and statistical manual of mental disorders: DSM-5. . American Psychiatric Association,
- Einhorn LC, Johansen PA, White FJ (1988) Electrophysiological effects of cocaine in the mesoaccumbens dopamine system: studies in the ventral tegmental area. *J Neurosci* 8 (1):100-112
- El Rawas R, Thiriet N, Lardeux V, Jaber M, Solinas M (2009) Environmental enrichment decreases the rewarding but not the activating effects of heroin. *Psychopharmacology (Berl)* 203 (3):561-570. doi:10.1007/s00213-008-1402-6
- El-Ghundi M, George SR, Drago J, Fletcher PJ, Fan T, Nguyen T, Liu C, Sibley DR, Westphal H, O'Dowd BF (1998) Disruption of dopamine D1 receptor gene expression attenuates alcohol-seeking behavior. *Eur J Pharmacol* 353 (2-3):149-158
- Elliot EE, Sibley DR, Katz JL (2003) Locomotor and discriminative-stimulus effects of cocaine in dopamine D5 receptor knockout mice. *Psychopharmacology (Berl)* 169 (2):161-168. doi:10.1007/s00213-003-1494-y
- Elmer GI, Pieper JO, Rubinstein M, Low MJ, Grandy DK, Wise RA (2002) Failure of intravenous morphine to serve as an effective instrumental reinforcer in dopamine D2 receptor knock-out mice. *J Neurosci* 22 (10):RC224. doi:20026412

- Engeln M, Fasano S, Ahmed SH, Cador M, Baekelandt V, Bezard E, Fernagut PO (2013) Levodopa gains psychostimulant-like properties after nigral dopaminergic loss. *Annals of neurology* 74 (1):140-144. doi:10.1002/ana.23881
- Epping-Jordan MP, Watkins SS, Koob GF, Markou A (1998) Dramatic decreases in brain reward function during nicotine withdrawal. *Nature* 393 (6680):76-79. doi:10.1038/30001
- Eravci M, Grosspietsch T, Pinna G, Schulz O, Kley S, Bachmann M, Wolffgramm J, Gotz E, Heyne A, Meinhold H, Baumgartner A (1997) Dopamine receptor gene expression in an animal model of 'behavioral dependence' on ethanol. *Brain Res Mol Brain Res* 50 (1-2):221-229
- Etter JF, Stapleton JA (2006) Nicotine replacement therapy for long-term smoking cessation: a meta-analysis. *Tob Control* 15 (4):280-285. doi:10.1136/tc.2005.015487
- Evans AH, Pavese N, Lawrence AD, Tai YF, Appel S, Doder M, Brooks DJ, Lees AJ, Piccini P (2006) Compulsive drug use linked to sensitized ventral striatal dopamine transmission. *Annals of neurology* 59 (5):852-858
- Everitt BJ (1990) Sexual motivation: a neural and behavioural analysis of the mechanisms underlying appetitive and copulatory responses of male rats. *Neurosci Biobehav Rev* 14 (2):217-232
- Everitt BJ, Belin D, Economidou D, Pelloux Y, Dalley JW, Robbins TW (2008) Review. Neural mechanisms underlying the vulnerability to develop compulsive drug-seeking habits and addiction. *Philos Trans R Soc Lond B Biol Sci* 363 (1507):3125-3135
- Everitt BJ, Robbins TW (2005) Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat Neurosci* 8 (11):1481-1489
- Fadda P, Scherma M, Fresu A, Collu M, Fratta W (2005) Dopamine and serotonin release in dorsal striatum and nucleus accumbens is differentially modulated by morphine in DBA/2J and C57BL/6J mice. *Synapse* 56 (1):29-38. doi:10.1002/syn.20122
- Faleiro LJ, Jones S, Kauer JA (2004) Rapid synaptic plasticity of glutamatergic synapses on dopamine neurons in the ventral tegmental area in response to acute amphetamine injection. *Neuropsychopharmacology* 29 (12):2115-2125. doi:10.1038/sj.npp.1300495
- Farnebo LO, Hamberger B (1971) Drug-induced changes in the release of 3 H-monoamines from field stimulated rat brain slices. *Acta Physiol Scand Suppl* 371:35-44
- Fasano C, Bourque MJ, Lapointe G, Leo D, Thibault D, Haber M, Kortleven C, Desgroseillers L, Murai KK, Trudeau LE (2013) Dopamine facilitates dendritic spine formation by cultured striatal medium spiny neurons through both D1 and D2 dopamine receptors. *Neuropharmacology* 67:432-443. doi:10.1016/j.neuropharm.2012.11.030
- Fauchey V, Jaber M, Bloch B, Le Moine C (2000a) Dopamine control of striatal gene expression during development: relevance to knockout mice for the dopamine transporter. *Eur J Neurosci* 12 (9):3415-3425
- Fauchey V, Jaber M, Caron MG, Bloch B, Le Moine C (2000b) Differential regulation of the dopamine D1, D2 and D3 receptor gene expression and changes in the phenotype of the striatal neurons in mice lacking the dopamine transporter. *Eur J Neurosci* 12 (1):19-26

- Feltenstein MW, See RE (2008) The neurocircuitry of addiction: an overview. *Br J Pharmacol* 154 (2):261-274. doi:10.1038/bjp.2008.51
- Fernagut PO, Chalon S, Diguët E, Guilloteau D, Tison F, Jaber M (2003) Motor behaviour deficits and their histopathological and functional correlates in the nigrostriatal system of dopamine transporter knockout mice. *Neuroscience* 116 (4):1123-1130
- Fernagut PO, Diguët E, Jaber M, Bioulac B, Tison F (2002) Dopamine transporter knockout mice are hypersensitive to 3-nitropropionic acid-induced striatal damage. *Eur J Neurosci* 15 (12):2053-2056
- Fibiger HC, Nomikos GG, Pfau JG, Damsma G (1992) Sexual behavior, eating and mesolimbic dopamine. *Clin Neuropharmacol* 15 Suppl 1 Pt A:566A-567A
- Fisher H, Aron A, Brown LL (2005) Romantic love: an fMRI study of a neural mechanism for mate choice. *J Comp Neurol* 493 (1):58-62. doi:10.1002/cne.20772
- Floresco SB, Blaha CD, Yang CR, Phillips AG (2001a) Dopamine D1 and NMDA receptors mediate potentiation of basolateral amygdala-evoked firing of nucleus accumbens neurons. *J Neurosci* 21 (16):6370-6376
- Floresco SB, Blaha CD, Yang CR, Phillips AG (2001b) Modulation of hippocampal and amygdalar-evoked activity of nucleus accumbens neurons by dopamine: cellular mechanisms of input selection. *J Neurosci* 21 (8):2851-2860
- Floresco SB, Todd CL, Grace AA (2001c) Glutamatergic afferents from the hippocampus to the nucleus accumbens regulate activity of ventral tegmental area dopamine neurons. *J Neurosci* 21 (13):4915-4922
- Floresco SB, West AR, Ash B, Moore H, Grace AA (2003) Afferent modulation of dopamine neuron firing differentially regulates tonic and phasic dopamine transmission. *Nat Neurosci* 6 (9):968-973
- Foddai M, Dosia G, Spiga S, Diana M (2004) Acetaldehyde increases dopaminergic neuronal activity in the VTA. *Neuropsychopharmacology* 29 (3):530-536. doi:10.1038/sj.npp.1300326
- Fourgeaud L, Mato S, Bouchet D, Hemar A, Worley PF, Manzoni OJ (2004) A single in vivo exposure to cocaine abolishes endocannabinoid-mediated long-term depression in the nucleus accumbens. *J Neurosci* 24 (31):6939-6945. doi:10.1523/JNEUROSCI.0671-04.2004
- Fouriez G, Hansson P, Wise RA (1978) Neuroleptic-induced attenuation of brain stimulation reward in rats. *J Comp Physiol Psychol* 92 (4):661-671
- French ED (1997) delta9-Tetrahydrocannabinol excites rat VTA dopamine neurons through activation of cannabinoid CB1 but not opioid receptors. *Neurosci Lett* 226 (3):159-162
- Futami T, Takakusaki K, Kitai ST (1995) Glutamatergic and cholinergic inputs from the pedunculopontine tegmental nucleus to dopamine neurons in the substantia nigra pars compacta. *Neurosci Res* 21 (4):331-342
- Gainetdinov RR (2008) Dopamine transporter mutant mice in experimental neuropharmacology. *Naunyn Schmiedebergs Arch Pharmacol* 377 (4-6):301-313. doi:10.1007/s00210-007-0216-0
- Gainetdinov RR, Wetsel WC, Jones SR, Levin ED, Jaber M, Caron MG (1999) Role of serotonin in the paradoxical calming effect of psychostimulants on hyperactivity. *Science* 283 (5400):397-401
- Gamaleddin IH, Trigo JM, Gueye AB, Zvonok A, Makriyannis A, Goldberg SR, Le Foll B (2015) Role of the endogenous cannabinoid system in nicotine addiction: novel insights. *Front Psychiatry* 6:41. doi:10.3389/fpsy.2015.00041

- Gao WY, Lee TH, King GR, Ellinwood EH (1998) Alterations in baseline activity and quinpirole sensitivity in putative dopamine neurons in the substantia nigra and ventral tegmental area after withdrawal from cocaine pretreatment. *Neuropsychopharmacology* 18 (3):222-232. doi:10.1016/S0893-133X(97)00132-2
- Garcia MM, Brown HE, Harlan RE (1995) Alterations in immediate-early gene proteins in the rat forebrain induced by acute morphine injection. *Brain Res* 692 (1-2):23-40
- Garris PA, Ciolkowski EL, Pastore P, Wightman RM (1994) Efflux of dopamine from the synaptic cleft in the nucleus accumbens of the rat brain. *J Neurosci* 14 (10):6084-6093
- Georges F, Aston-Jones G (2001) Potent regulation of midbrain dopamine neurons by the bed nucleus of the stria terminalis. *J Neurosci* 21 (16):RC160
- Georges F, Stinus L, Bloch B, Le Moine C (1999) Chronic morphine exposure and spontaneous withdrawal are associated with modifications of dopamine receptor and neuropeptide gene expression in the rat striatum. *Eur J Neurosci* 11 (2):481-490
- Georgiadis JR, Kringelbach ML, Pfaus JG (2012) Sex for fun: a synthesis of human and animal neurobiology. *Nat Rev Urol* 9 (9):486-498. doi:10.1038/nrurol.2012.151
- Gerfen CR, Surmeier DJ (2011) Modulation of striatal projection systems by dopamine. *Annu Rev Neurosci* 34:441-466. doi:10.1146/annurev-neuro-061010-113641
- Gessa GL, Muntoni F, Collu M, Vargiu L, Mereu G (1985) Low doses of ethanol activate dopaminergic neurons in the ventral tegmental area. *Brain Res* 348 (1):201-203
- Ghisi V, Ramsey AJ, Masri B, Gainetdinov RR, Caron MG, Salahpour A (2009) Reduced D2-mediated signaling activity and trans-synaptic upregulation of D1 and D2 dopamine receptors in mice overexpressing the dopamine transporter. *Cell Signal* 21 (1):87-94. doi:10.1016/j.cellsig.2008.09.011
- Giovannoni G, O'Sullivan JD, Turner K, Manson AJ, Lees AJ (2000) Hedonistic homeostatic dysregulation in patients with Parkinson's disease on dopamine replacement therapies. *Journal of neurology, neurosurgery, and psychiatry* 68 (4):423-428
- Gipson CD, Kupchik YM, Kalivas PW (2014) Rapid, transient synaptic plasticity in addiction. *Neuropharmacology* 76 Pt B:276-286. doi:10.1016/j.neuropharm.2013.04.032
- Gipson CD, Olive MF (2017) Structural and functional plasticity of dendritic spines - root or result of behavior? *Genes Brain Behav* 16 (1):101-117. doi:10.1111/gbb.12324
- Girault JA, Valjent E, Caboche J, Herve D (2007) ERK2: a logical AND gate critical for drug-induced plasticity? *Curr Opin Pharmacol* 7 (1):77-85. doi:10.1016/j.coph.2006.08.012
- Giros B, Jaber M, Jones SR, Wightman RM, Caron MG (1996) Hyperlocomotion and indifference to cocaine and amphetamine in mice lacking the dopamine transporter. *Nature* 379 (6566):606-612. doi:10.1038/379606a0
- Goeders NE (2002) Stress and cocaine addiction. *J Pharmacol Exp Ther* 301 (3):785-789
- Goto Y, Grace AA (2005) Dopamine-dependent interactions between limbic and prefrontal cortical plasticity in the nucleus accumbens: disruption by cocaine sensitization. *Neuron* 47 (2):255-266
- Gould RW, Duke AN, Nader MA (2014) PET studies in nonhuman primate models of cocaine abuse: translational research related to vulnerability and

- neuroadaptations. *Neuropharmacology* 84:138-151.
doi:10.1016/j.neuropharm.2013.02.004
- Gozzi A, Tessari M, Dacome L, Agosta F, Lepore S, Lanzoni A, Cristofori P, Pich EM, Corsi M, Bifone A (2011) Neuroimaging evidence of altered fronto-cortical and striatal function after prolonged cocaine self-administration in the rat. *Neuropsychopharmacology* 36 (12):2431-2440. doi:10.1038/npp.2011.129
- Grace AA (2000) The tonic/phasic model of dopamine system regulation and its implications for understanding alcohol and psychostimulant craving. *Addiction* 95 Suppl 2:S119-128
- Grace AA, Bunney BS (1979) Paradoxical GABA excitation of nigral dopaminergic cells: indirect mediation through reticulata inhibitory neurons. *Eur J Pharmacol* 59 (3-4):211-218
- Grace AA, Bunney BS (1984) The control of firing pattern in nigral dopamine neurons: burst firing. *J Neurosci* 4 (11):2877-2890
- Grace AA, Floresco SB, Goto Y, Lodge DJ (2007) Regulation of firing of dopaminergic neurons and control of goal-directed behaviors. *Trends Neurosci* 30 (5):220-227. doi:10.1016/j.tins.2007.03.003
- Graybiel AM (1998) The basal ganglia and chunking of action repertoires. *Neurobiol Learn Mem* 70 (1-2):119-136. doi:10.1006/nlme.1998.3843
- Graybiel AM, Moratalla R, Robertson HA (1990) Amphetamine and cocaine induce drug-specific activation of the c-fos gene in striosome-matrix compartments and limbic subdivisions of the striatum. *Proc Natl Acad Sci U S A* 87 (17):6912-6916
- Grenhoff J, Aston-Jones G, Svensson TH (1986) Nicotinic effects on the firing pattern of midbrain dopamine neurons. *Acta Physiol Scand* 128 (3):351-358. doi:10.1111/j.1748-1716.1986.tb07988.x
- Grimm JW, Hope BT, Wise RA, Shaham Y (2001) Neuroadaptation. Incubation of cocaine craving after withdrawal. *Nature* 412 (6843):141-142
- Groenewegen HJ, Wright CI, Beijer AV, Voorn P (1999) Convergence and segregation of ventral striatal inputs and outputs. *Ann N Y Acad Sci* 877:49-63
- Groves PM, Wilson CJ, Young SJ, Rebec GV (1975) Self-inhibition by dopaminergic neurons. *Science* 190 (4214):522-528
- Grueter BA, Brasnjo G, Malenka RC (2010) Postsynaptic TRPV1 triggers cell type-specific long-term depression in the nucleus accumbens. *Nat Neurosci* 13 (12):1519-1525. doi:10.1038/nn.2685
- Grunhage F, Schulze TG, Muller DJ, Lanczik M, Franzek E, Albus M, Borrmann-Hassenbach M, Knapp M, Cichon S, Maier W, Rietschel M, Propping P, Nothen MM (2000) Systematic screening for DNA sequence variation in the coding region of the human dopamine transporter gene (DAT1). *Mol Psychiatry* 5 (3):275-282
- Haber SN, Knutson B (2010) The reward circuit: linking primate anatomy and human imaging. *Neuropsychopharmacology* 35 (1):4-26. doi:10.1038/npp.2009.129
- Haber SN, Lynd E, Klein C, Groenewegen HJ (1990) Topographic organization of the ventral striatal efferent projections in the rhesus monkey: an anterograde tracing study. *J Comp Neurol* 293 (2):282-298
- Hamdi A, Prasad C (1993) Bidirectional changes in striatal D1-dopamine receptor density during chronic ethanol intake. *Life Sci* 52 (3):251-257
- Hammond C, Deniau JM, Rizk A, Feger J (1978) Electrophysiological demonstration of an excitatory subthalamonigral pathway in the rat. *Brain Res* 151 (2):235-244

- Haney M, Ward AS, Foltin RW, Fischman MW (2001) Effects of ecopipam, a selective dopamine D1 antagonist, on smoked cocaine self-administration by humans. *Psychopharmacology (Berl)* 155 (4):330-337
- Hanlon CA, Wesley MJ, Porrino LJ (2009) Loss of functional specificity in the dorsal striatum of chronic cocaine users. *Drug Alcohol Depend* 102 (1-3):88-94. doi:10.1016/j.drugalcdep.2009.01.005
- Haustein KO (2000) Pharmacotherapy of nicotine dependence. *Int J Clin Pharmacol Ther* 38 (6):273-290
- Heck N, Dos Santos M, Amairi B, Salery M, Besnard A, Herzog E, Boudier T, Vanhoutte P, Caboche J (2015) A new automated 3D detection of synaptic contacts reveals the formation of cortico-striatal synapses upon cocaine treatment in vivo. *Brain Struct Funct* 220 (5):2953-2966. doi:10.1007/s00429-014-0837-2
- Hedreen JC, DeLong MR (1991) Organization of striatopallidal, striatonigral, and nigrostriatal projections in the macaque. *J Comp Neurol* 304 (4):569-595. doi:10.1002/cne.903040406
- Henry DJ, Greene MA, White FJ (1989) Electrophysiological effects of cocaine in the mesoaccumbens dopamine system: repeated administration. *J Pharmacol Exp Ther* 251 (3):833-839
- Herin DV, Rush CR, Grabowski J (2010) Agonist-like pharmacotherapy for stimulant dependence: preclinical, human laboratory, and clinical studies. *Ann N Y Acad Sci* 1187:76-100. doi:10.1111/j.1749-6632.2009.05145.x
- Hersch SM, Yi H, Heilman CJ, Edwards RH, Levey AI (1997) Subcellular localization and molecular topology of the dopamine transporter in the striatum and substantia nigra. *J Comp Neurol* 388 (2):211-227
- Higley AE, Kiefer SW, Li X, Gaal J, Xi ZX, Gardner EL (2011) Dopamine D(3) receptor antagonist SB-277011A inhibits methamphetamine self-administration and methamphetamine-induced reinstatement of drug-seeking in rats. *Eur J Pharmacol* 659 (2-3):187-192. doi:10.1016/j.ejphar.2011.02.046
- Hildebrand BE, Nomikos GG, Hertel P, Schilstrom B, Svensson TH (1998) Reduced dopamine output in the nucleus accumbens but not in the medial prefrontal cortex in rats displaying a mecamylamine-precipitated nicotine withdrawal syndrome. *Brain Res* 779 (1-2):214-225
- Hjelmstad GO, Xia Y, Margolis EB, Fields HL (2013) Opioid modulation of ventral pallidal afferents to ventral tegmental area neurons. *J Neurosci* 33 (15):6454-6459. doi:10.1523/JNEUROSCI.0178-13.2013
- Hoffman AF, Lupica CR (2001) Direct actions of cannabinoids on synaptic transmission in the nucleus accumbens: a comparison with opioids. *J Neurophysiol* 85 (1):72-83. doi:10.1152/jn.2001.85.1.72
- Hoffman DC, Dickson PR, Beninger RJ (1988) The dopamine D2 receptor agonists, quinpirole and bromocriptine produce conditioned place preferences. *Prog Neuropsychopharmacol Biol Psychiatry* 12 (2-3):315-322
- Hong S, Jhou TC, Smith M, Saleem KS, Hikosaka O (2011) Negative reward signals from the lateral habenula to dopamine neurons are mediated by rostromedial tegmental nucleus in primates. *J Neurosci* 31 (32):11457-11471. doi:10.1523/JNEUROSCI.1384-11.2011
- Hope B, Kosofsky B, Hyman SE, Nestler EJ (1992) Regulation of immediate early gene expression and AP-1 binding in the rat nucleus accumbens by chronic cocaine. *Proc Natl Acad Sci U S A* 89 (13):5764-5768

- Hope BT, Nye HE, Kelz MB, Self DW, Iadarola MJ, Nakabeppu Y, Duman RS, Nestler EJ (1994) Induction of a long-lasting AP-1 complex composed of altered Fos-like proteins in brain by chronic cocaine and other chronic treatments. *Neuron* 13 (5):1235-1244
- Hummel M, Unterwald EM (2002) D1 dopamine receptor: a putative neurochemical and behavioral link to cocaine action. *J Cell Physiol* 191 (1):17-27. doi:10.1002/jcp.10078
- Hyland BI, Reynolds JN, Hay J, Perk CG, Miller R (2002) Firing modes of midbrain dopamine cells in the freely moving rat. *Neuroscience* 114 (2):475-492
- Hyman SE (2005) Addiction: a disease of learning and memory. *Am J Psychiatry* 162 (8):1414-1422
- Hyman SE, Malenka RC, Nestler EJ (2006) Neural mechanisms of addiction: the role of reward-related learning and memory. *Annu Rev Neurosci* 29:565-598. doi:10.1146/annurev.neuro.29.051605.113009
- Imperato A, Di Chiara G (1986) Preferential stimulation of dopamine release in the nucleus accumbens of freely moving rats by ethanol. *J Pharmacol Exp Ther* 239 (1):219-228
- Ito R, Dalley JW, Howes SR, Robbins TW, Everitt BJ (2000) Dissociation in conditioned dopamine release in the nucleus accumbens core and shell in response to cocaine cues and during cocaine-seeking behavior in rats. *J Neurosci* 20 (19):7489-7495
- Ito R, Dalley JW, Robbins TW, Everitt BJ (2002) Dopamine release in the dorsal striatum during cocaine-seeking behavior under the control of a drug-associated cue. *J Neurosci* 22 (14):6247-6253. doi:20026606
- Ito R, Robbins TW, Everitt BJ (2004) Differential control over cocaine-seeking behavior by nucleus accumbens core and shell. *Nat Neurosci* 7 (4):389-397. doi:10.1038/nn1217
- Jaber M (2006) Monoamine Transporters. . In: Gorwood P, Hamon M (eds) Klüwer Publ.
- Jaber M, Dumartin B, Sagne C, Haycock JW, Roubert C, Giros B, Bloch B, Caron MG (1999) Differential regulation of tyrosine hydroxylase in the basal ganglia of mice lacking the dopamine transporter. *Eur J Neurosci* 11 (10):3499-3511
- Jaber M, Jones S, Giros B, Caron MG (1997) The dopamine transporter: a crucial component regulating dopamine transmission. *Mov Disord* 12 (5):629-633. doi:10.1002/mds.870120502
- Jaber M, Robinson SW, Missale C, Caron MG (1996) Dopamine receptors and brain function. *Neuropharmacology* 35 (11):1503-1519
- Jalabert M, Bourdy R, Courtin J, Veinante P, Manzoni OJ, Barrot M, Georges F (2011) Neuronal circuits underlying acute morphine action on dopamine neurons. *Proc Natl Acad Sci U S A* 108 (39):16446-16450. doi:10.1073/pnas.1105418108
- Janknecht R (1995) Regulation of the c-fos promoter. *Immunobiology* 193 (2-4):137-142. doi:10.1016/S0171-2985(11)80536-X
- Jedynak JP, Uslaner JM, Esteban JA, Robinson TE (2007) Methamphetamine-induced structural plasticity in the dorsal striatum. *Eur J Neurosci* 25 (3):847-853. doi:10.1111/j.1460-9568.2007.05316.x
- Jessor R, Jessor S (1980) A social-psychological framework for studying drug use. *NIDA Res Monogr* 30:102-109
- Ji H, Shepard PD (2007) Lateral habenula stimulation inhibits rat midbrain dopamine neurons through a GABA(A) receptor-mediated mechanism. *J Neurosci* 27 (26):6923-6930. doi:10.1523/JNEUROSCI.0958-07.2007

- Jiang ZG, North RA (1992) Pre- and postsynaptic inhibition by opioids in rat striatum. *J Neurosci* 12 (1):356-361
- Jiao X, Pare WP, Tejani-Butt SM (2006) Alcohol consumption alters dopamine transporter sites in Wistar-Kyoto rat brain. *Brain Res* 1073-1074:175-182. doi:10.1016/j.brainres.2005.12.009
- Johnson SW, North RA (1992) Opioids excite dopamine neurons by hyperpolarization of local interneurons. *J Neurosci* 12 (2):483-488
- Jonckers E, Shah D, Hamaide J, Verhoye M, Van der Linden A (2015) The power of using functional fMRI on small rodents to study brain pharmacology and disease. *Front Pharmacol* 6:231. doi:10.3389/fphar.2015.00231
- Jones S, Bonci A (2005) Synaptic plasticity and drug addiction. *Curr Opin Pharmacol* 5 (1):20-25
- Jones SR, Gainetdinov RR, Hu XT, Cooper DC, Wightman RM, White FJ, Caron MG (1999) Loss of autoreceptor functions in mice lacking the dopamine transporter. *Nat Neurosci* 2 (7):649-655. doi:10.1038/10204
- Jones SR, Gainetdinov RR, Jaber M, Giros B, Wightman RM, Caron MG (1998) Profound neuronal plasticity in response to inactivation of the dopamine transporter. *Proc Natl Acad Sci U S A* 95 (7):4029-4034
- Kalivas PW (1993) Neurotransmitter regulation of dopamine neurons in the ventral tegmental area. *Brain Res Brain Res Rev* 18 (1):75-113
- Kalivas PW, Stewart J (1991) Dopamine transmission in the initiation and expression of drug- and stress-induced sensitization of motor activity. *Brain Res Brain Res Rev* 16 (3):223-244
- Katz JL, Chausmer AL, Elmer GI, Rubinstein M, Low MJ, Grandy DK (2003) Cocaine-induced locomotor activity and cocaine discrimination in dopamine D4 receptor mutant mice. *Psychopharmacology (Berl)* 170 (1):108-114. doi:10.1007/s00213-003-1513-z
- Kauer JA, Malenka RC (2007) Synaptic plasticity and addiction. *Nat Rev Neurosci* 8 (11):844-858. doi:10.1038/nrn2234
- Kaufling J, Veinante P, Pawlowski SA, Freund-Mercier MJ, Barrot M (2009) Afferents to the GABAergic tail of the ventral tegmental area in the rat. *J Comp Neurol* 513 (6):597-621. doi:10.1002/cne.21983
- Kelz MB, Chen J, Carlezon WA, Jr., Whisler K, Gilden L, Beckmann AM, Steffen C, Zhang YJ, Marotti L, Self DW, Tkatch T, Baranauskas G, Surmeier DJ, Neve RL, Duman RS, Picciotto MR, Nestler EJ (1999) Expression of the transcription factor deltaFosB in the brain controls sensitivity to cocaine. *Nature* 401 (6750):272-276. doi:10.1038/45790
- Kenny PJ, Markou A (2001) Neurobiology of the nicotine withdrawal syndrome. *Pharmacol Biochem Behav* 70 (4):531-549
- Kiba H, Jayaraman A (1994) Nicotine induced c-fos expression in the striatum is mediated mostly by dopamine D1 receptor and is dependent on NMDA stimulation. *Brain Res Mol Brain Res* 23 (1-2):1-13
- Kim J, Park BH, Lee JH, Park SK, Kim JH (2011) Cell type-specific alterations in the nucleus accumbens by repeated exposures to cocaine. *Biol Psychiatry* 69 (11):1026-1034. doi:10.1016/j.biopsych.2011.01.013
- Kim MS, Yu JH, Kim CH, Choi JY, Seo JH, Lee MY, Yi CH, Choi TH, Ryu YH, Lee JE, Lee BH, Kim H, Cho SR (2016) Environmental enrichment enhances synaptic plasticity by internalization of striatal dopamine transporters. *J Cereb Blood Flow Metab* 36 (12):2122-2133. doi:10.1177/0271678X15613525

- Kim YC, Alberico SL, Emmons E, Narayanan NS (2015) New therapeutic strategies targeting D1-type dopamine receptors for neuropsychiatric disease. *Front Biol (Beijing)* 10 (3):230-238. doi:10.1007/s11515-015-1360-4
- Knackstedt LA, Trantham-Davidson HL, Schwendt M (2014) The role of ventral and dorsal striatum mGluR5 in relapse to cocaine-seeking and extinction learning. *Addict Biol* 19 (1):87-101. doi:10.1111/adb.12061
- Kodjo CM, Klein JD (2002) Prevention and risk of adolescent substance abuse. The role of adolescents, families, and communities. *Pediatr Clin North Am* 49 (2):257-268
- Koepp MJ, Gunn RN, Lawrence AD, Cunningham VJ, Dagher A, Jones T, Brooks DJ, Bench CJ, Grasby PM (1998) Evidence for striatal dopamine release during a video game. *Nature* 393 (6682):266-268. doi:10.1038/30498
- Kolb B, Li Y, Robinson T, Parker LA (2018) THC alters morphology of neurons in medial prefrontal cortex, orbital prefrontal cortex, and nucleus accumbens and alters the ability of later experience to promote structural plasticity. *Synapse* 72 (3). doi:10.1002/syn.22020
- Koob GF (2006) The neurobiology of addiction: a neuroadaptational view relevant for diagnosis. *Addiction* 101 Suppl 1:23-30
- Koob GF (2008) A role for brain stress systems in addiction. *Neuron* 59 (1):11-34. doi:S0896-6273(08)00530-8 [pii] 10.1016/j.neuron.2008.06.012
- Koob GF, Caine SB, Parsons L, Markou A, Weiss F (1997) Opponent process model and psychostimulant addiction. *Pharmacol Biochem Behav* 57 (3):513-521
- Koob GF, Le Moal M (1997) Drug abuse: hedonic homeostatic dysregulation. *Science* 278 (5335):52-58
- Koob GF, Le Moal M (2001) Drug addiction, dysregulation of reward, and allostasis. *Neuropsychopharmacology* 24 (2):97-129
- Koob GF, Volkow ND (2010) Neurocircuitry of addiction. *Neuropsychopharmacology* 35 (1):217-238. doi:10.1038/npp.2009.110
- Koob GF, Volkow ND (2016) Neurobiology of addiction: a neurocircuitry analysis. *Lancet Psychiatry* 3 (8):760-773. doi:10.1016/S2215-0366(16)00104-8
- Koranda JL, Cone JJ, McGehee DS, Roitman MF, Beeler JA, Zhuang X (2014) Nicotinic receptors regulate the dynamic range of dopamine release in vivo. *J Neurophysiol* 111 (1):103-111. doi:10.1152/jn.00269.2013
- Kourrich S, Rothwell PE, Klug JR, Thomas MJ (2007) Cocaine experience controls bidirectional synaptic plasticity in the nucleus accumbens. *J Neurosci* 27 (30):7921-7928. doi:10.1523/JNEUROSCI.1859-07.2007
- Koyama S, Brodie MS, Appel SB (2007) Ethanol inhibition of m-current and ethanol-induced direct excitation of ventral tegmental area dopamine neurons. *J Neurophysiol* 97 (3):1977-1985. doi:10.1152/jn.00270.2006
- Kreek MJ, Nielsen DA, Butelman ER, LaForge KS (2005) Genetic influences on impulsivity, risk taking, stress responsivity and vulnerability to drug abuse and addiction. *Nat Neurosci* 8 (11):1450-1457
- Kuhar MJ, Pilotte NS (1996) Neurochemical changes in cocaine withdrawal. *Trends Pharmacol Sci* 17 (7):260-264
- Kunko PM, French D, Izenwasser S (1998) Alterations in locomotor activity during chronic cocaine administration: effect on dopamine receptors and interaction with opioids. *J Pharmacol Exp Ther* 285 (1):277-284

- Kunz G, Englisch HJ, Wenzel J (1976) [Spine-distribution of pyramidal neurons of the CA1-region of the rat hippocampus following long-term oral alcohol administration]. *J Hirnforsch* 17 (4):351-363
- Kupchik YM, Brown RM, Heinsbroek JA, Lobo MK, Schwartz DJ, Kalivas PW (2015) Coding the direct/indirect pathways by D1 and D2 receptors is not valid for accumbens projections. *Nat Neurosci* 18 (9):1230-1232. doi:10.1038/nn.4068
- Lafragette A, Bardo MT, Lardeux V, Solinas M, Thiriet N (2017) Reduction of Cocaine-Induced Locomotor Effects by Enriched Environment Is Associated with Cell-Specific Accumulation of DeltaFosB in Striatal and Cortical Subregions. *Int J Neuropsychopharmacol* 20 (3):237-246. doi:10.1093/ijnp/pyw097
- Lammel S, Ion DI, Roeper J, Malenka RC (2011) Projection-specific modulation of dopamine neuron synapses by aversive and rewarding stimuli. *Neuron* 70 (5):855-862
- Laurier LG, Corrigall WA, George SR (1994) Dopamine receptor density, sensitivity and mRNA levels are altered following self-administration of cocaine in the rat. *Brain Res* 634 (1):31-40
- Lawrence AD, Evans AH, Lees AJ (2003) Compulsive use of dopamine replacement therapy in Parkinson's disease: reward systems gone awry? *Lancet neurology* 2 (10):595-604
- Le Foll B, Collo G, Rabiner EA, Boileau I, Merlo Pich E, Sokoloff P (2014) Dopamine D3 receptor ligands for drug addiction treatment: update on recent findings. *Prog Brain Res* 211:255-275. doi:10.1016/B978-0-444-63425-2.00011-8
- Le Foll B, Gallo A, Le Strat Y, Lu L, Gorwood P (2009) Genetics of dopamine receptors and drug addiction: a comprehensive review. *Behav Pharmacol* 20 (1):1-17. doi:10.1097/FBP.0b013e3283242f05
- Lecca D, Cacciapaglia F, Valentini V, Di Chiara G (2006a) Monitoring extracellular dopamine in the rat nucleus accumbens shell and core during acquisition and maintenance of intravenous WIN 55,212-2 self-administration. *Psychopharmacology (Berl)* 188 (1):63-74. doi:10.1007/s00213-006-0475-3
- Lecca D, Cacciapaglia F, Valentini V, Gronli J, Spiga S, Di Chiara G (2006b) Preferential increase of extracellular dopamine in the rat nucleus accumbens shell as compared to that in the core during acquisition and maintenance of intravenous nicotine self-administration. *Psychopharmacology (Berl)* 184 (3-4):435-446. doi:10.1007/s00213-005-0280-4
- Lecca D, Valentini V, Cacciapaglia F, Acquas E, Di Chiara G (2007) Reciprocal effects of response contingent and noncontingent intravenous heroin on in vivo nucleus accumbens shell versus core dopamine in the rat: a repeated sampling microdialysis study. *Psychopharmacology (Berl)* 194 (1):103-116. doi:10.1007/s00213-007-0815-y
- Lecca S, Melis M, Luchicchi A, Muntoni AL, Pistis M (2012) Inhibitory inputs from rostromedial tegmental neurons regulate spontaneous activity of midbrain dopamine cells and their responses to drugs of abuse. *Neuropsychopharmacology* 37 (5):1164-1176. doi:10.1038/npp.2011.302
- Lecca S, Meye FJ, Mamei M (2014) The lateral habenula in addiction and depression: an anatomical, synaptic and behavioral overview. *Eur J Neurosci* 39 (7):1170-1178. doi:10.1111/ejn.12480
- Lee BR, Ma YY, Huang YH, Wang X, Otaka M, Ishikawa M, Neumann PA, Graziane NM, Brown TE, Suska A, Guo C, Lobo MK, Sesack SR, Wolf ME, Nestler EJ, Shaham Y, Schluter OM, Dong Y (2013) Maturation of silent synapses in amygdala-

- accumbens projection contributes to incubation of cocaine craving. *Nat Neurosci* 16 (11):1644-1651. doi:10.1038/nn.3533
- Lee KW, Kim Y, Kim AM, Helmin K, Nairn AC, Greengard P (2006) Cocaine-induced dendritic spine formation in D1 and D2 dopamine receptor-containing medium spiny neurons in nucleus accumbens. *Proc Natl Acad Sci U S A* 103 (9):3399-3404. doi:10.1073/pnas.0511244103
- Lee TH, Gao WY, Davidson C, Ellinwood EH (1999) Altered activity of midbrain dopamine neurons following 7-day withdrawal from chronic cocaine abuse is normalized by D2 receptor stimulation during the early withdrawal phase. *Neuropsychopharmacology* 21 (1):127-136. doi:10.1016/S0893-133X(99)00011-1
- Letchworth SR, Daunais JB, Hedgecock AA, Porrino LJ (1997) Effects of chronic cocaine administration on dopamine transporter mRNA and protein in the rat. *Brain Res* 750 (1-2):214-222
- Letchworth SR, Nader MA, Smith HR, Friedman DP, Porrino LJ (2001) Progression of changes in dopamine transporter binding site density as a result of cocaine self-administration in rhesus monkeys. *J Neurosci* 21 (8):2799-2807
- Leyton M, Boileau I, Benkelfat C, Diksic M, Baker G, Dagher A (2002) Amphetamine-induced increases in extracellular dopamine, drug wanting, and novelty seeking: a PET/[11C]raclopride study in healthy men. *Neuropsychopharmacology* 27 (6):1027-1035. doi:10.1016/S0893-133X(02)00366-4
- Liggins J, Pihl RO, Benkelfat C, Leyton M (2012) The dopamine augmenter L-DOPA does not affect positive mood in healthy human volunteers. *PloS one* 7 (1):e28370. doi:10.1371/journal.pone.0028370
- Liu J, Nickolenko J, Sharp FR (1994) Morphine induces c-fos and junB in striatum and nucleus accumbens via D1 and N-methyl-D-aspartate receptors. *Proc Natl Acad Sci U S A* 91 (18):8537-8541
- Liu X, Weiss F (2002) Reversal of ethanol-seeking behavior by D1 and D2 antagonists in an animal model of relapse: differences in antagonist potency in previously ethanol-dependent versus nondependent rats. *J Pharmacol Exp Ther* 300 (3):882-889
- Lobo MK, Covington HE, 3rd, Chaudhury D, Friedman AK, Sun H, Damez-Werno D, Dietz DM, Zaman S, Koo JW, Kennedy PJ, Mouzon E, Mogri M, Neve RL, Deisseroth K, Han MH, Nestler EJ (2010) Cell type-specific loss of BDNF signaling mimics optogenetic control of cocaine reward. *Science* 330 (6002):385-390. doi:10.1126/science.1188472
- Lobo MK, Nestler EJ (2011) The striatal balancing act in drug addiction: distinct roles of direct and indirect pathway medium spiny neurons. *Front Neuroanat* 5:41. doi:10.3389/fnana.2011.00041
- Lodge DJ, Grace AA (2005) Acute and chronic corticotropin-releasing factor 1 receptor blockade inhibits cocaine-induced dopamine release: correlation with dopamine neuron activity. *J Pharmacol Exp Ther* 314 (1):201-206. doi:10.1124/jpet.105.084913
- Lodge DJ, Grace AA (2006) The laterodorsal tegmentum is essential for burst firing of ventral tegmental area dopamine neurons. *Proc Natl Acad Sci U S A* 103 (13):5167-5172. doi:10.1073/pnas.0510715103
- Lodge DJ, Grace AA (2008) Amphetamine activation of hippocampal drive of mesolimbic dopamine neurons: a mechanism of behavioral sensitization. *J Neurosci* 28 (31):7876-7882. doi:10.1523/JNEUROSCI.1582-08.2008

- Lu W, Wolf ME (1999) Repeated amphetamine administration alters AMPA receptor subunit expression in rat nucleus accumbens and medial prefrontal cortex. *Synapse* 32 (2):119-131. doi:10.1002/(SICI)1098-2396(199905)32:2<119::AID-SYN5>3.0.CO;2-F
- Luscher C, Malenka RC (2011) Drug-evoked synaptic plasticity in addiction: from molecular changes to circuit remodeling. *Neuron* 69 (4):650-663. doi:10.1016/j.neuron.2011.01.017
- Luscher C, Ungless MA (2006) The mechanistic classification of addictive drugs. *PLoS Med* 3 (11):e437. doi:10.1371/journal.pmed.0030437
- Lynd-Balta E, Haber SN (1994) The organization of midbrain projections to the striatum in the primate: sensorimotor-related striatum versus ventral striatum. *Neuroscience* 59 (3):625-640
- Ma YY, Lee BR, Wang X, Guo C, Liu L, Cui R, Lan Y, Balcita-Pedicino JJ, Wolf ME, Sesack SR, Shaham Y, Schluter OM, Huang YH, Dong Y (2014) Bidirectional modulation of incubation of cocaine craving by silent synapse-based remodeling of prefrontal cortex to accumbens projections. *Neuron* 83 (6):1453-1467. doi:10.1016/j.neuron.2014.08.023
- MacAskill AF, Little JP, Cassel JM, Carter AG (2012) Subcellular connectivity underlies pathway-specific signaling in the nucleus accumbens. *Nat Neurosci* 15 (12):1624-1626. doi:10.1038/nn.3254
- Macey DJ, Rice WN, Freedland CS, Whitlow CT, Porrino LJ (2004) Patterns of functional activity associated with cocaine self-administration in the rat change over time. *Psychopharmacology (Berl)* 172 (4):384-392. doi:10.1007/s00213-003-1676-7
- Maggos CE, Spangler R, Zhou Y, Schlussman SD, Ho A, Kreek MJ (1997) Quantitation of dopamine transporter mRNA in the rat brain: mapping, effects of "binge" cocaine administration and withdrawal. *Synapse* 26 (1):55-61. doi:10.1002/(SICI)1098-2396(199705)26:1<55::AID-SYN6>3.0.CO;2-D
- Mahler SV, Vazey EM, Beckley JT, Keistler CR, McGlinchey EM, Kaufling J, Wilson SP, Deisseroth K, Woodward JJ, Aston-Jones G (2014) Designer receptors show role for ventral pallidum input to ventral tegmental area in cocaine seeking. *Nat Neurosci* 17 (4):577-585. doi:10.1038/nn.3664
- Mailleux P, Verslype M, Preud'homme X, Vanderhaeghen JJ (1994) Activation of multiple transcription factor genes by tetrahydrocannabinol in rat forebrain. *Neuroreport* 5 (10):1265-1268
- Maldonado R, Saiardi A, Valverde O, Samad TA, Roques BP, Borrelli E (1997) Absence of opiate rewarding effects in mice lacking dopamine D2 receptors. *Nature* 388 (6642):586-589. doi:10.1038/41567
- Mameli M, Halbout B, Creton C, Engblom D, Parkitna JR, Spanagel R, Luscher C (2009) Cocaine-evoked synaptic plasticity: persistence in the VTA triggers adaptations in the NAc. *Nat Neurosci* 12 (8):1036-1041
- Mantsch JR, Baker DA, Funk D, Le AD, Shaham Y (2016) Stress-Induced Reinstatement of Drug Seeking: 20 Years of Progress. *Neuropsychopharmacology* 41 (1):335-356. doi:10.1038/npp.2015.142
- Marinelli M, Piazza PV (2002) Interaction between glucocorticoid hormones, stress and psychostimulant drugs. *Eur J Neurosci* 16 (3):387-394
- Markou A, Koob GF (1991) Postcocaine anhedonia. An animal model of cocaine withdrawal. *Neuropsychopharmacology* 4 (1):17-26
- Martin BJ, Naughton BJ, Thirtamara-Rajamani K, Yoon DJ, Han DD, Devries AC, Gu HH (2011) Dopamine transporter inhibition is necessary for cocaine-induced

- increases in dendritic spine density in the nucleus accumbens. *Synapse* 65 (6):490-496. doi:10.1002/syn.20865
- Martin M, Chen BT, Hopf FW, Bowers MS, Bonci A (2006) Cocaine self-administration selectively abolishes LTD in the core of the nucleus accumbens. *Nat Neurosci* 9 (7):868-869
- Martinez D, Gil R, Slifstein M, Hwang DR, Huang Y, Perez A, Kegeles L, Talbot P, Evans S, Krystal J, Laruelle M, Abi-Dargham A (2005) Alcohol dependence is associated with blunted dopamine transmission in the ventral striatum. *Biol Psychiatry* 58 (10):779-786. doi:10.1016/j.biopsych.2005.04.044
- Mathews TA, John CE, Lapa GB, Budygin EA, Jones SR (2006) No role of the dopamine transporter in acute ethanol effects on striatal dopamine dynamics. *Synapse* 60 (4):288-294. doi:10.1002/syn.20301
- Matsumoto M, Hikosaka O (2007) Lateral habenula as a source of negative reward signals in dopamine neurons. *Nature* 447 (7148):1111-1115. doi:10.1038/nature05860
- May LJ, Kuhr WG, Wightman RM (1988) Differentiation of dopamine overflow and uptake processes in the extracellular fluid of the rat caudate nucleus with fast-scan in vivo voltammetry. *J Neurochem* 51 (4):1060-1069
- Maze I, Covington HE, 3rd, Dietz DM, LaPlant Q, Renthal W, Russo SJ, Mechanic M, Mouzon E, Neve RL, Haggarty SJ, Ren Y, Sampath SC, Hurd YL, Greengard P, Tarakhovskiy A, Schaefer A, Nestler EJ (2010) Essential role of the histone methyltransferase G9a in cocaine-induced plasticity. *Science* 327 (5962):213-216. doi:10.1126/science.1179438
- Mazei-Robison MS, Couch RS, Shelton RC, Stein MA, Blakely RD (2005) Sequence variation in the human dopamine transporter gene in children with attention deficit hyperactivity disorder. *Neuropharmacology* 49 (6):724-736. doi:10.1016/j.neuropharm.2005.08.003
- McBride WJ, Murphy JM, Ikemoto S (1999) Localization of brain reinforcement mechanisms: intracranial self-administration and intracranial place-conditioning studies. *Behav Brain Res* 101 (2):129-152
- McClung CA, Ulerly PG, Perrotti LI, Zachariou V, Berton O, Nestler EJ (2004) DeltaFosB: a molecular switch for long-term adaptation in the brain. *Brain Res Mol Brain Res* 132 (2):146-154
- McGeorge AJ, Faull RL (1989) The organization of the projection from the cerebral cortex to the striatum in the rat. *Neuroscience* 29 (3):503-537
- McGinty JF, Shi XD, Schwendt M, Saylor A, Toda S (2008) Regulation of psychostimulant-induced signaling and gene expression in the striatum. *J Neurochem* 104 (6):1440-1449. doi:10.1111/j.1471-4159.2008.05240.x
- McQuade JA, Xu M, Woods SC, Seeley RJ, Benoit SC (2003) Ethanol consumption in mice with a targeted disruption of the dopamine-3 receptor gene. *Addict Biol* 8 (3):295-303. doi:10.1080/13556210310001602202
- Mello NK, Negus SS (1996) Preclinical evaluation of pharmacotherapies for treatment of cocaine and opioid abuse using drug self-administration procedures. *Neuropsychopharmacology* 14 (6):375-424. doi:10.1016/0893-133X(95)00274-H
- Mereu G, Gessa GL (1984) Ethanol excites dopamine (DA) neurons and inhibits non-dopamine (non-DA) neurons in the Substantia nigra of rats. *Ann Ist Super Sanita* 20 (1):11-15

- Mergy MA, Gowrishankar R, Gresch PJ, Gantz SC, Williams J, Davis GL, Wheeler CA, Stanwood GD, Hahn MK, Blakely RD (2014) The rare DAT coding variant Val559 perturbs DA neuron function, changes behavior, and alters in vivo responses to psychostimulants. *Proc Natl Acad Sci U S A* 111 (44):E4779-4788. doi:10.1073/pnas.1417294111
- Micevych PE, Meisel RL (2017) Integrating Neural Circuits Controlling Female Sexual Behavior. *Front Syst Neurosci* 11:42. doi:10.3389/fnsys.2017.00042
- Milella MS, Fotros A, Gravel P, Casey KF, Larcher K, Verhaeghe JA, Cox SM, Reader AJ, Dagher A, Benkelfat C, Leyton M (2016) Cocaine cue-induced dopamine release in the human prefrontal cortex. *J Psychiatry Neurosci* 41 (5):322-330. doi:10.1503/jpn.150207
- Miner LL, Drago J, Chamberlain PM, Donovan D, Uhl GR (1995) Retained cocaine conditioned place preference in D1 receptor deficient mice. *Neuroreport* 6 (17):2314-2316
- Moratalla R, Robertson HA, Graybiel AM (1992) Dynamic regulation of NGFI-A (zif268, egr1) gene expression in the striatum. *J Neurosci* 12 (7):2609-2622
- Morgan D, Grant KA, Gage HD, Mach RH, Kaplan JR, Prioleau O, Nader SH, Buchheimer N, Ehrenkaufer RL, Nader MA (2002) Social dominance in monkeys: dopamine D2 receptors and cocaine self-administration. *Nat Neurosci* 5 (2):169-174. doi:10.1038/nn798
- Morice E, Denis C, Giros B, Nosten-Bertrand M (2010) Evidence of long-term expression of behavioral sensitization to both cocaine and ethanol in dopamine transporter knockout mice. *Psychopharmacology (Berl)* 208 (1):57-66. doi:10.1007/s00213-009-1707-0
- Moron JA, Brockington A, Wise RA, Rocha BA, Hope BT (2002) Dopamine uptake through the norepinephrine transporter in brain regions with low levels of the dopamine transporter: evidence from knock-out mouse lines. *J Neurosci* 22 (2):389-395
- Nader MA, Czoty PW, Gould RW, Riddick NV (2008) Review. Positron emission tomography imaging studies of dopamine receptors in primate models of addiction. *Philos Trans R Soc Lond B Biol Sci* 363 (1507):3223-3232. doi:10.1098/rstb.2008.0092
- Nader MA, Daunais JB, Moore T, Nader SH, Moore RJ, Smith HR, Friedman DP, Porrino LJ (2002) Effects of cocaine self-administration on striatal dopamine systems in rhesus monkeys: initial and chronic exposure. *Neuropsychopharmacology* 27 (1):35-46. doi:10.1016/S0893-133X(01)00427-4
- Nader MA, Morgan D, Gage HD, Nader SH, Calhoun TL, Buchheimer N, Ehrenkaufer R, Mach RH (2006) PET imaging of dopamine D2 receptors during chronic cocaine self-administration in monkeys. *Nat Neurosci* 9 (8):1050-1056. doi:10.1038/nn1737
- Nausieda PA (1985) Sinemet "abusers". *Clin Neuropharmacol* 8 (4):318-327
- Nestler EJ (1996) Under siege: The brain on opiates. *Neuron* 16 (5):897-900
- Nestler EJ (2001) Molecular basis of long-term plasticity underlying addiction. *Nat Rev Neurosci* 2 (2):119-128
- Nestler EJ (2004) Molecular mechanisms of drug addiction. *Neuropharmacology* 47 Suppl 1:24-32. doi:10.1016/j.neuropharm.2004.06.031
- Nestler EJ, Barrot M, Self DW (2001) DeltaFosB: a sustained molecular switch for addiction. *Proc Natl Acad Sci U S A* 98 (20):11042-11046

- Newman AH, Blaylock BL, Nader MA, Bergman J, Sibley DR, Skolnick P (2012) Medication discovery for addiction: translating the dopamine D3 receptor hypothesis. *Biochem Pharmacol* 84 (7):882-890. doi:10.1016/j.bcp.2012.06.023
- Nicolas C, Tauber C, Lepelletier FX, Chalon S, Belujon P, Galineau L, Solinas M (2017) Longitudinal Changes in Brain Metabolic Activity after Withdrawal from Escalation of Cocaine Self-Administration. *Neuropsychopharmacology* 42 (10):1981-1990. doi:10.1038/npp.2017.109
- Norman AB, Norman MK, Hall JF, Tsibulsky VL (1999) Priming threshold: a novel quantitative measure of the reinstatement of cocaine self-administration. *Brain Res* 831 (1-2):165-174
- Norrholm SD, Bibb JA, Nestler EJ, Ouimet CC, Taylor JR, Greengard P (2003) Cocaine-induced proliferation of dendritic spines in nucleus accumbens is dependent on the activity of cyclin-dependent kinase-5. *Neuroscience* 116 (1):19-22
- Nutt DJ, Lingford-Hughes A, Erritzoe D, Stokes PR (2015) The dopamine theory of addiction: 40 years of highs and lows. *Nat Rev Neurosci* 16 (5):305-312. doi:10.1038/nrn3939
- O'Donnell P (2003) Dopamine gating of forebrain neural ensembles. *Eur J Neurosci* 17 (3):429-435
- O'Donnell P, Grace AA (1995) Synaptic interactions among excitatory afferents to nucleus accumbens neurons: hippocampal gating of prefrontal cortical input. *J Neurosci* 15 (5 Pt 1):3622-3639
- O'Neill B, Tilley MR, Han DD, Thirtamara-Rajamani K, Hill ER, Bishop GA, Zhou FM, During MJ, Gu HH (2014) Behavior of knock-in mice with a cocaine-insensitive dopamine transporter after virogenetic restoration of cocaine sensitivity in the striatum. *Neuropharmacology* 79:626-633. doi:10.1016/j.neuropharm.2013.12.023
- Olds J, Milner P (1954) Positive reinforcement produced by electrical stimulation of septal area and other regions of rat brain. *J Comp Physiol Psychol* 47 (6):419-427
- Onn SP, Grace AA (2000) Amphetamine withdrawal alters bistable states and cellular coupling in rat prefrontal cortex and nucleus accumbens neurons recorded in vivo. *J Neurosci* 20 (6):2332-2345
- Ortinski PI, Vassoler FM, Carlson GC, Pierce RC (2012) Temporally dependent changes in cocaine-induced synaptic plasticity in the nucleus accumbens shell are reversed by D1-like dopamine receptor stimulation. *Neuropsychopharmacology* 37 (7):1671-1682. doi:10.1038/npp.2012.12
- Pang TY, Hannan AJ, Lawrence AJ (2018) Novel approaches to alcohol rehabilitation: Modification of stress-responsive brain regions through environmental enrichment. *Neuropharmacology*. doi:10.1016/j.neuropharm.2018.02.021
- Parsons LH, Smith AD, Justice JB, Jr. (1991) Basal extracellular dopamine is decreased in the rat nucleus accumbens during abstinence from chronic cocaine. *Synapse* 9 (1):60-65. doi:10.1002/syn.890090109
- Paterson NE, Myers C, Markou A (2000) Effects of repeated withdrawal from continuous amphetamine administration on brain reward function in rats. *Psychopharmacology (Berl)* 152 (4):440-446
- Pattison LP, McIntosh S, Sexton T, Childers SR, Hemby SE (2014) Changes in dopamine transporter binding in nucleus accumbens following chronic self-administration cocaine: heroin combinations. *Synapse* 68 (10):437-444. doi:10.1002/syn.21755

- Patton MH, Bizup BT, Grace AA (2013) The infralimbic cortex bidirectionally modulates mesolimbic dopamine neuron activity via distinct neural pathways. *J Neurosci* 33 (43):16865-16873. doi:10.1523/JNEUROSCI.2449-13.2013
- Pecina S, Cagniard B, Berridge KC, Aldridge JW, Zhuang X (2003) Hyperdopaminergic mutant mice have higher "wanting" but not "liking" for sweet rewards. *J Neurosci* 23 (28):9395-9402
- Perrotti LI, Weaver RR, Robison B, Renthal W, Maze I, Yazdani S, Elmore RG, Knapp DJ, Selley DE, Martin BR, Sim-Selley L, Bachtell RK, Self DW, Nestler EJ (2008) Distinct patterns of DeltaFosB induction in brain by drugs of abuse. *Synapse* 62 (5):358-369. doi:10.1002/syn.20500
- Pezzella FR, Colosimo C, Vanacore N, Di Rezze S, Chianese M, Fabbrini G, Meco G (2005) Prevalence and clinical features of hedonistic homeostatic dysregulation in Parkinson's disease. *Mov Disord* 20 (1):77-81. doi:10.1002/mds.20288
- Phillips PE, Robinson DL, Stuber GD, Carelli RM, Wightman RM (2003a) Real-time measurements of phasic changes in extracellular dopamine concentration in freely moving rats by fast-scan cyclic voltammetry. *Methods Mol Med* 79:443-464
- Phillips PE, Stuber GD, Heien ML, Wightman RM, Carelli RM (2003b) Subsecond dopamine release promotes cocaine seeking. *Nature* 422 (6932):614-618. doi:10.1038/nature01476
- Piazza PV, Le Moal ML (1996) Pathophysiological basis of vulnerability to drug abuse: role of an interaction between stress, glucocorticoids, and dopaminergic neurons. *Annu Rev Pharmacol Toxicol* 36:359-378. doi:10.1146/annurev.pa.36.040196.002043
- Pierce RC, Kalivas PW (1997) A circuitry model of the expression of behavioral sensitization to amphetamine-like psychostimulants. *Brain Res Brain Res Rev* 25 (2):192-216
- Pierce RC, Kumaresan V (2006) The mesolimbic dopamine system: the final common pathway for the reinforcing effect of drugs of abuse? *Neurosci Biobehav Rev* 30 (2):215-238. doi:10.1016/j.neubiorev.2005.04.016
- Pierce RC, O'Brien CP, Kenny PJ, Vanderschuren LJ (2012) Rational development of addiction pharmacotherapies: successes, failures, and prospects. *Cold Spring Harb Perspect Med* 2 (6):a012880. doi:10.1101/cshperspect.a012880
- Pierce RC, Wolf ME (2013) Psychostimulant-induced neuroadaptations in nucleus accumbens AMPA receptor transmission. *Cold Spring Harb Perspect Med* 3 (2):a012021. doi:10.1101/cshperspect.a012021
- Pontieri FE, Tanda G, Di Chiara G (1995) Intravenous cocaine, morphine, and amphetamine preferentially increase extracellular dopamine in the "shell" as compared with the "core" of the rat nucleus accumbens. *Proc Natl Acad Sci U S A* 92 (26):12304-12308
- Pontieri FE, Tanda G, Orzi F, Di Chiara G (1996) Effects of nicotine on the nucleus accumbens and similarity to those of addictive drugs. *Nature* 382 (6588):255-257. doi:10.1038/382255a0
- Porrino LJ, Daunais JB, Smith HR, Nader MA (2004) The expanding effects of cocaine: studies in a nonhuman primate model of cocaine self-administration. *Neurosci Biobehav Rev* 27 (8):813-820. doi:10.1016/j.neubiorev.2003.11.013
- Pothos E, Rada P, Mark GP, Hoebel BG (1991) Dopamine microdialysis in the nucleus accumbens during acute and chronic morphine, naloxone-precipitated withdrawal and clonidine treatment. *Brain Res* 566 (1-2):348-350

- Purgianto A, Scheyer AF, Loweth JA, Ford KA, Tseng KY, Wolf ME (2013) Different adaptations in AMPA receptor transmission in the nucleus accumbens after short vs long access cocaine self-administration regimens. *Neuropsychopharmacology* 38 (9):1789-1797. doi:10.1038/npp.2013.78
- Rabinak CA, Nirenberg MJ (2010) Dopamine agonist withdrawal syndrome in Parkinson disease. *Arch Neurol* 67 (1):58-63. doi:
- Ranaldi R, Pocock D, Zereik R, Wise RA (1999) Dopamine fluctuations in the nucleus accumbens during maintenance, extinction, and reinstatement of intravenous D-amphetamine self-administration. *J Neurosci* 19 (10):4102-4109
- Rasmussen K, Czachura JF (1995) Nicotine withdrawal leads to increased firing rates of midbrain dopamine neurons. *Neuroreport* 7 (1):329-332
- Ren T, Sagar SM (1992) Induction of c-fos immunostaining in the rat brain after the systemic administration of nicotine. *Brain Res Bull* 29 (5):589-597
- Ren Z, Sun WL, Jiao H, Zhang D, Kong H, Wang X, Xu M (2010) Dopamine D1 and N-methyl-D-aspartate receptors and extracellular signal-regulated kinase mediate neuronal morphological changes induced by repeated cocaine administration. *Neuroscience* 168 (1):48-60. doi:10.1016/j.neuroscience.2010.03.034
- Renthal W, Nestler EJ (2008) Epigenetic mechanisms in drug addiction. *Trends Mol Med* 14 (8):341-350
- Rickhag M, Hansen FH, Sorensen G, Strandfelt KN, Andresen B, Gotfryd K, Madsen KL, Vestergaard-Klewe I, Ammendrup-Johnsen I, Eriksen J, Newman AH, Fuchtbauer EM, Gomeza J, Woldbye DP, Wortwein G, Gether U (2013) A C-terminal PDZ domain-binding sequence is required for striatal distribution of the dopamine transporter. *Nat Commun* 4:1580. doi:10.1038/ncomms2568
- Riddle JL, Rokosik SL, Napier TC (2012) Pramipexole- and methamphetamine-induced reward-mediated behavior in a rodent model of Parkinson's disease and controls. *Behav Brain Res* 233 (1):15-23. doi:10.1016/j.bbr.2012.04.027
- Riley JN, Walker DW (1978) Morphological alterations in hippocampus after long-term alcohol consumption in mice. *Science* 201 (4356):646-648
- Risinger FO, Freeman PA, Rubinstein M, Low MJ, Grandy DK (2000) Lack of operant ethanol self-administration in dopamine D2 receptor knockout mice. *Psychopharmacology (Berl)* 152 (3):343-350
- Robinson DL, Venton BJ, Heien ML, Wightman RM (2003) Detecting subsecond dopamine release with fast-scan cyclic voltammetry in vivo. *Clin Chem* 49 (10):1763-1773
- Robinson TE, Berridge KC (1993) The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Res Brain Res Rev* 18 (3):247-291
- Robinson TE, Berridge KC (2008) Review. The incentive sensitization theory of addiction: some current issues. *Philos Trans R Soc Lond B Biol Sci* 363 (1507):3137-3146
- Robinson TE, Gorny G, Mitton E, Kolb B (2001) Cocaine self-administration alters the morphology of dendrites and dendritic spines in the nucleus accumbens and neocortex. *Synapse* 39 (3):257-266. doi:10.1002/1098-2396(20010301)39:3<257::AID-SYN1007>3.0.CO;2-1
- Robinson TE, Kolb B (1997) Persistent structural modifications in nucleus accumbens and prefrontal cortex neurons produced by previous experience with amphetamine. *J Neurosci* 17 (21):8491-8497

- Robinson TE, Kolb B (1999) Morphine alters the structure of neurons in the nucleus accumbens and neocortex of rats. *Synapse* 33 (2):160-162. doi:10.1002/(SICI)1098-2396(199908)33:2<160::AID-SYN6>3.0.CO;2-S
- Robinson TE, Kolb B (2004) Structural plasticity associated with exposure to drugs of abuse. *Neuropharmacology* 47 Suppl 1:33-46. doi:10.1016/j.neuropharm.2004.06.025
- Rocha BA, Fumagalli F, Gainetdinov RR, Jones SR, Ator R, Giros B, Miller GW, Caron MG (1998) Cocaine self-administration in dopamine-transporter knockout mice. *Nat Neurosci* 1 (2):132-137. doi:10.1038/381
- Rossetti ZL, Hmaidan Y, Gessa GL (1992) Marked inhibition of mesolimbic dopamine release: a common feature of ethanol, morphine, cocaine and amphetamine abstinence in rats. *Eur J Pharmacol* 221 (2-3):227-234
- Rossi MA, Yin HH (2015) Elevated dopamine alters consummatory pattern generation and increases behavioral variability during learning. *Front Integr Neurosci* 9:37. doi:10.3389/fnint.2015.00037
- Rothman RB, Baumann MH, Prisinzano TE, Newman AH (2008) Dopamine transport inhibitors based on GBR12909 and benztropine as potential medications to treat cocaine addiction. *Biochem Pharmacol* 75 (1):2-16. doi:10.1016/j.bcp.2007.08.007
- Rouge-Pont F, Usiello A, Benoit-Marand M, Gonon F, Piazza PV, Borrelli E (2002) Changes in extracellular dopamine induced by morphine and cocaine: crucial control by D2 receptors. *J Neurosci* 22 (8):3293-3301. doi:20026322
- Rubinstein M, Phillips TJ, Bunzow JR, Falzone TL, Dziewczapolski G, Zhang G, Fang Y, Larson JL, McDougall JA, Chester JA, Saez C, Pugsley TA, Gershanik O, Low MJ, Grandy DK (1997) Mice lacking dopamine D4 receptors are supersensitive to ethanol, cocaine, and methamphetamine. *Cell* 90 (6):991-1001
- Ruesink GB, Georgiadis JR (2017) Brain Imaging of Human Sexual Response: Recent Developments and Future Directions. *Curr Sex Health Rep* 9 (4):183-191. doi:10.1007/s11930-017-0123-4
- Russo SJ, Dietz DM, Dumitriu D, Morrison JH, Malenka RC, Nestler EJ (2010) The addicted synapse: mechanisms of synaptic and structural plasticity in nucleus accumbens. *Trends Neurosci* 33 (6):267-276. doi:10.1016/j.tins.2010.02.002
- Salahpour A, Ramsey AJ, Medvedev IO, Kile B, Sotnikova TD, Holmstrand E, Ghisi V, Nicholls PJ, Wong L, Murphy K, Sesack SR, Wightman RM, Gainetdinov RR, Caron MG (2008) Increased amphetamine-induced hyperactivity and reward in mice overexpressing the dopamine transporter. *Proc Natl Acad Sci U S A* 105 (11):4405-4410. doi:10.1073/pnas.0707646105
- Salamone JD, Correa M, Farrar A, Mingote SM (2007) Effort-related functions of nucleus accumbens dopamine and associated forebrain circuits. *Psychopharmacology (Berl)* 191 (3):461-482. doi:10.1007/s00213-006-0668-9
- Schmidt HD, Anderson SM, Pierce RC (2006) Stimulation of D1-like or D2 dopamine receptors in the shell, but not the core, of the nucleus accumbens reinstates cocaine-seeking behaviour in the rat. *Eur J Neurosci* 23 (1):219-228. doi:10.1111/j.1460-9568.2005.04524.x
- Schulteis G, Markou A, Cole M, Koob GF (1995) Decreased brain reward produced by ethanol withdrawal. *Proc Natl Acad Sci U S A* 92 (13):5880-5884
- Schulteis G, Markou A, Gold LH, Stinus L, Koob GF (1994) Relative sensitivity to naloxone of multiple indices of opiate withdrawal: a quantitative dose-response analysis. *J Pharmacol Exp Ther* 271 (3):1391-1398

- Schultz W (1997) Dopamine neurons and their role in reward mechanisms. *Curr Opin Neurobiol* 7 (2):191-197
- Schultz W (1998) The phasic reward signal of primate dopamine neurons. *Adv Pharmacol* 42:686-690
- Schultz W (2011) Potential vulnerabilities of neuronal reward, risk, and decision mechanisms to addictive drugs. *Neuron* 69 (4):603-617. doi:10.1016/j.neuron.2011.02.014
- Schultz W (2016) Dopamine reward prediction error coding. *Dialogues Clin Neurosci* 18 (1):23-32
- Schultz W, Dayan P, Montague PR (1997) A neural substrate of prediction and reward. *Science* 275 (5306):1593-1599
- See RE, Elliott JC, Feltenstein MW (2007) The role of dorsal vs ventral striatal pathways in cocaine-seeking behavior after prolonged abstinence in rats. *Psychopharmacology (Berl)* 194 (3):321-331. doi:10.1007/s00213-007-0850-8
- Sesack SR, Grace AA (2010) Cortico-Basal Ganglia reward network: microcircuitry. *Neuropsychopharmacology* 35 (1):27-47
- Shaham Y, Shalev U, Lu L, De Wit H, Stewart J (2003) The reinstatement model of drug relapse: history, methodology and major findings. *Psychopharmacology (Berl)* 168 (1-2):3-20. doi:10.1007/s00213-002-1224-x
- Shalev U, Grimm JW, Shaham Y (2002) Neurobiology of relapse to heroin and cocaine seeking: a review. *Pharmacol Rev* 54 (1):1-42
- Sharp T, Zetterstrom T, Ungerstedt U (1986) An in vivo study of dopamine release and metabolism in rat brain regions using intracerebral dialysis. *J Neurochem* 47 (1):113-122
- Shepard JD, Chuang DT, Shaham Y, Morales M (2006) Effect of methamphetamine self-administration on tyrosine hydroxylase and dopamine transporter levels in mesolimbic and nigrostriatal dopamine pathways of the rat. *Psychopharmacology (Berl)* 185 (4):505-513. doi:10.1007/s00213-006-0316-4
- Shepard PD, Bunney BS (1988) Effects of apamin on the discharge properties of putative dopamine-containing neurons in vitro. *Brain Res* 463 (2):380-384
- Silvia CP, Jaber M, King GR, Ellinwood EH, Caron MG (1997) Cocaine and amphetamine elicit differential effects in rats with a unilateral injection of dopamine transporter antisense oligodeoxynucleotides. *Neuroscience* 76 (3):737-747
- Sinha R (2001) How does stress increase risk of drug abuse and relapse? *Psychopharmacology (Berl)* 158 (4):343-359
- Sinha R (2007) The role of stress in addiction relapse. *Curr Psychiatry Rep* 9 (5):388-395
- Sinha R (2008) Chronic stress, drug use, and vulnerability to addiction. *Ann N Y Acad Sci* 1141:105-130. doi:10.1196/annals.1441.030
- Smith ID, Grace AA (1992) Role of the subthalamic nucleus in the regulation of nigral dopamine neuron activity. *Synapse* 12 (4):287-303. doi:10.1002/syn.890120406
- Solinas M, Goldberg SR, Piomelli D (2008) The endocannabinoid system in brain reward processes. *Br J Pharmacol* 154 (2):369-383. doi:10.1038/bjp.2008.130
- Solinas M, Scherma M, Tanda G, Wertheim CE, Fratta W, Goldberg SR (2007) Nicotinic facilitation of delta9-tetrahydrocannabinol discrimination involves endogenous anandamide. *J Pharmacol Exp Ther* 321 (3):1127-1134. doi:10.1124/jpet.106.116830
- Solinas M, Tanda G, Wertheim CE, Goldberg SR (2010a) Dopaminergic augmentation of delta-9-tetrahydrocannabinol (THC) discrimination: possible involvement of

- D(2)-induced formation of anandamide. *Psychopharmacology (Berl)* 209 (2):191-202. doi:10.1007/s00213-010-1789-8
- Solinas M, Thiriet N, Chauvet C, Jaber M (2010b) Prevention and treatment of drug addiction by environmental enrichment. *Prog Neurobiol* 92 (4):572-592. doi:S0301-0082(10)00145-0 [pii]
10.1016/j.pneurobio.2010.08.002
- Solinas M, Thiriet N, El Rawas R, Lardeux V, Jaber M (2009) Environmental enrichment during early stages of life reduces the behavioral, neurochemical, and molecular effects of cocaine. *Neuropsychopharmacology* 34 (5):1102-1111. doi:10.1038/npp.2008.51
- Solomon RL, Corbit JD (1974) An opponent-process theory of motivation. I. Temporal dynamics of affect. *Psychol Rev* 81 (2):119-145
- Sora I, Wichems C, Takahashi N, Li XF, Zeng Z, Revay R, Lesch KP, Murphy DL, Uhl GR (1998) Cocaine reward models: conditioned place preference can be established in dopamine- and in serotonin-transporter knockout mice. *Proc Natl Acad Sci U S A* 95 (13):7699-7704
- Sousa FC, Gomes PB, Macedo DS, Marinho MM, Viana GS (1999) Early withdrawal from repeated cocaine administration upregulates muscarinic and dopaminergic D2-like receptors in rat neostriatum. *Pharmacol Biochem Behav* 62 (1):15-20
- Spielewoy C, Gonon F, Roubert C, Fauchey V, Jaber M, Caron MG, Roques BP, Hamon M, Betancur C, Maldonado R, Giros B (2000) Increased rewarding properties of morphine in dopamine-transporter knockout mice. *Eur J Neurosci* 12 (5):1827-1837
- Spiga S, Mulas G, Piras F, Diana M (2014a) The "addicted" spine. *Front Neuroanat* 8:110. doi:10.3389/fnana.2014.00110
- Spiga S, Talani G, Mulas G, Licheri V, Fois GR, Muggironi G, Masala N, Cannizzaro C, Biggio G, Sanna E, Diana M (2014b) Hampered long-term depression and thin spine loss in the nucleus accumbens of ethanol-dependent rats. *Proc Natl Acad Sci U S A* 111 (35):E3745-3754. doi:10.1073/pnas.1406768111
- Stairs DJ, Bardo MT (2009) Neurobehavioral effects of environmental enrichment and drug abuse vulnerability. *Pharmacol Biochem Behav* 92 (3):377-382. doi:10.1016/j.pbb.2009.01.016
- Stefanski R, Ladenheim B, Lee SH, Cadet JL, Goldberg SR (1999) Neuroadaptations in the dopaminergic system after active self-administration but not after passive administration of methamphetamine. *Eur J Pharmacol* 371 (2-3):123-135
- Stefanski R, Lee SH, Yasar S, Cadet JL, Goldberg SR (2002) Lack of persistent changes in the dopaminergic system of rats withdrawn from methamphetamine self-administration. *Eur J Pharmacol* 439 (1-3):59-68
- Stuber GD, Hopf FW, Hahn J, Cho SL, Guillory A, Bonci A (2008) Voluntary ethanol intake enhances excitatory synaptic strength in the ventral tegmental area. *Alcohol Clin Exp Res* 32 (10):1714-1720. doi:10.1111/j.1530-0277.2008.00749.x
- Suckling J, Nestor LJ (2017) The neurobiology of addiction: the perspective from magnetic resonance imaging present and future. *Addiction* 112 (2):360-369. doi:10.1111/add.13474
- Sung KW, Choi S, Lovinger DM (2001) Activation of group I mGluRs is necessary for induction of long-term depression at striatal synapses. *J Neurophysiol* 86 (5):2405-2412. doi:10.1152/jn.2001.86.5.2405
- Swainson R, Rogers RD, Sahakian BJ, Summers BA, Polkey CE, Robbins TW (2000) Probabilistic learning and reversal deficits in patients with Parkinson's disease or

- frontal or temporal lobe lesions: possible adverse effects of dopaminergic medication. *Neuropsychologia* 38 (5):596-612
- Szabo J (1979) Strionigral and nigrostriatal connections. *Anatomical studies. Appl Neurophysiol* 42 (1-2):9-12
- Tajuddin NF, Druse MJ (1996) Effects of chronic alcohol consumption and aging on dopamine D2 receptors in Fischer 344 rats. *Alcohol Clin Exp Res* 20 (1):144-151
- Tanda G, Loddo P, Di Chiara G (1999) Dependence of mesolimbic dopamine transmission on delta9-tetrahydrocannabinol. *Eur J Pharmacol* 376 (1-2):23-26
- Tanda G, Pontieri FE, Di Chiara G (1997) Cannabinoid and heroin activation of mesolimbic dopamine transmission by a common mu1 opioid receptor mechanism. *Science* 276 (5321):2048-2050
- Tepper JM, Lee CR (2007) GABAergic control of substantia nigra dopaminergic neurons. *Prog Brain Res* 160:189-208. doi:10.1016/S0079-6123(06)60011-3
- Tepper JM, Martin LP, Anderson DR (1995) GABAA receptor-mediated inhibition of rat substantia nigra dopaminergic neurons by pars reticulata projection neurons. *J Neurosci* 15 (4):3092-3103
- Terwilliger RZ, Beitner-Johnson D, Sevarino KA, Crain SM, Nestler EJ (1991) A general role for adaptations in G-proteins and the cyclic AMP system in mediating the chronic actions of morphine and cocaine on neuronal function. *Brain Res* 548 (1-2):100-110
- Thiriet N, Amar L, Toussay X, Lardeux V, Ladenheim B, Becker KG, Cadet JL, Solinas M, Jaber M (2008) Environmental enrichment during adolescence regulates gene expression in the striatum of mice. *Brain Res* 1222:31-41. doi:10.1016/j.brainres.2008.05.030
- Thiriet N, Aunis D, Zwiller J (2000) C-fos and egr-1 immediate-early gene induction by cocaine and cocaethylene in rat brain: a comparative study. *Ann N Y Acad Sci* 914:46-57
- Thiriet N, Jouvert P, Gobaille S, Solov'eva O, Gough B, Aunis D, Ali S, Zwiller J (2001a) C-type natriuretic peptide (CNP) regulates cocaine-induced dopamine increase and immediate early gene expression in rat brain. *Eur J Neurosci* 14 (10):1702-1708
- Thiriet N, Zwiller J, Ali SF (2001b) Induction of the immediate early genes egr-1 and c-fos by methamphetamine in mouse brain. *Brain Res* 919 (1):31-40
- Thomas MJ, Beurrier C, Bonci A, Malenka RC (2001) Long-term depression in the nucleus accumbens: a neural correlate of behavioral sensitization to cocaine. *Nat Neurosci* 4 (12):1217-1223. doi:10.1038/nn757
- Thomsen M, Han DD, Gu HH, Caine SB (2009) Lack of cocaine self-administration in mice expressing a cocaine-insensitive dopamine transporter. *J Pharmacol Exp Ther* 331 (1):204-211. doi:10.1124/jpet.109.156265
- Torres GE, Yao WD, Mohn AR, Quan H, Kim KM, Levey AI, Staudinger J, Caron MG (2001) Functional interaction between monoamine plasma membrane transporters and the synaptic PDZ domain-containing protein PICK1. *Neuron* 30 (1):121-134
- Tripathi A, Prensa L, Cebrian C, Mengual E (2010) Axonal branching patterns of nucleus accumbens neurons in the rat. *J Comp Neurol* 518 (22):4649-4673. doi:10.1002/cne.22484
- Tzavara ET, Li DL, Moutsimilli L, Bisogno T, Di Marzo V, Phebus LA, Nomikos GG, Giros B (2006) Endocannabinoids activate transient receptor potential vanilloid 1 receptors to reduce hyperdopaminergia-related hyperactivity: therapeutic implications. *Biol Psychiatry* 59 (6):508-515. doi:10.1016/j.biopsych.2005.08.019

- Ungerstedt U (1971) Adipsia and aphagia after 6-hydroxydopamine induced degeneration of the nigro-striatal dopamine system. *Acta Physiol Scand Suppl* 367:95-122
- Ungless MA, Argilli E, Bonci A (2010) Effects of stress and aversion on dopamine neurons: implications for addiction. *Neurosci Biobehav Rev* 35 (2):151-156. doi:10.1016/j.neubiorev.2010.04.006
- Ungless MA, Whistler JL, Malenka RC, Bonci A (2001) Single cocaine exposure in vivo induces long-term potentiation in dopamine neurons. *Nature* 411 (6837):583-587. doi:10.1038/35079077
- Unterwald EM, Fillmore J, Kreek MJ (1996) Chronic repeated cocaine administration increases dopamine D1 receptor-mediated signal transduction. *Eur J Pharmacol* 318 (1):31-35
- Unterwald EM, Ho A, Rubinfeld JM, Kreek MJ (1994) Time course of the development of behavioral sensitization and dopamine receptor up-regulation during binge cocaine administration. *J Pharmacol Exp Ther* 270 (3):1387-1396
- Urban NB, Kegeles LS, Slifstein M, Xu X, Martinez D, Sakr E, Castillo F, Moadel T, O'Malley SS, Krystal JH, Abi-Dargham A (2010) Sex differences in striatal dopamine release in young adults after oral alcohol challenge: a positron emission tomography imaging study with [(1)(1)C]raclopride. *Biol Psychiatry* 68 (8):689-696. doi:10.1016/j.biopsych.2010.06.005
- Valjent E, Bertran-Gonzalez J, Aubier B, Greengard P, Herve D, Girault JA (2010) Mechanisms of locomotor sensitization to drugs of abuse in a two-injection protocol. *Neuropsychopharmacology* 35 (2):401-415. doi:10.1038/npp.2009.143
- Valjent E, Corbille AG, Bertran-Gonzalez J, Herve D, Girault JA (2006) Inhibition of ERK pathway or protein synthesis during reexposure to drugs of abuse erases previously learned place preference. *Proc Natl Acad Sci U S A* 103 (8):2932-2937. doi:10.1073/pnas.0511030103
- Valjent E, Corvol JC, Pages C, Besson MJ, Maldonado R, Caboche J (2000) Involvement of the extracellular signal-regulated kinase cascade for cocaine-rewarding properties. *J Neurosci* 20 (23):8701-8709
- Valjent E, Mitchell JM, Besson MJ, Caboche J, Maldonado R (2002) Behavioural and biochemical evidence for interactions between Delta 9-tetrahydrocannabinol and nicotine. *Br J Pharmacol* 135 (2):564-578. doi:10.1038/sj.bjp.0704479
- van der Kooy D, Swerdlow NR, Koob GF (1983) Paradoxical reinforcing properties of apomorphine: effects of nucleus accumbens and area postrema lesions. *Brain Res* 259 (1):111-118
- Vanderschuren LJ, Di Ciano P, Everitt BJ (2005) Involvement of the dorsal striatum in cue-controlled cocaine seeking. *J Neurosci* 25 (38):8665-8670. doi:10.1523/JNEUROSCI.0925-05.2005
- Vanderschuren LJ, Everitt BJ (2004) Drug seeking becomes compulsive after prolonged cocaine self-administration. *Science* 305 (5686):1017-1019. doi:10.1126/science.1098975
- Verrico CD, Haile CN, Newton TF, Kosten TR, De La Garza R, 2nd (2013) Pharmacotherapeutics for substance-use disorders: a focus on dopaminergic medications. *Expert Opin Investig Drugs* 22 (12):1549-1568. doi:10.1517/13543784.2013.836488
- Volkow ND, Fowler JS, Wang GJ, Baler R, Telang F (2009) Imaging dopamine's role in drug abuse and addiction. *Neuropharmacology* 56 Suppl 1:3-8. doi:10.1016/j.neuropharm.2008.05.022

- Volkow ND, Fowler JS, Wang GJ, Hitzemann R, Logan J, Schlyer DJ, Dewey SL, Wolf AP (1993) Decreased dopamine D2 receptor availability is associated with reduced frontal metabolism in cocaine abusers. *Synapse* 14 (2):169-177. doi:10.1002/syn.890140210
- Volkow ND, Fowler JS, Wang GJ, Swanson JM (2004) Dopamine in drug abuse and addiction: results from imaging studies and treatment implications. *Mol Psychiatry* 9 (6):557-569
- Volkow ND, Koob GF, McLellan AT (2016) Neurobiologic Advances from the Brain Disease Model of Addiction. *N Engl J Med* 374 (4):363-371. doi:10.1056/NEJMra1511480
- Volkow ND, Wang GJ, Fowler JS, Logan J, Gatley SJ, Gifford A, Hitzemann R, Ding YS, Pappas N (1999) Prediction of reinforcing responses to psychostimulants in humans by brain dopamine D2 receptor levels. *Am J Psychiatry* 156 (9):1440-1443. doi:10.1176/ajp.156.9.1440
- Volkow ND, Wang GJ, Fowler JS, Thanos PP, Logan J, Gatley SJ, Gifford A, Ding YS, Wong C, Pappas N (2002) Brain DA D2 receptors predict reinforcing effects of stimulants in humans: replication study. *Synapse* 46 (2):79-82. doi:10.1002/syn.10137
- Volkow ND, Wang GJ, Fowler JS, Tomasi D, Telang F (2011) Addiction: beyond dopamine reward circuitry. *Proc Natl Acad Sci U S A* 108 (37):15037-15042. doi:10.1073/pnas.1010654108
- Volkow ND, Wang GJ, Telang F, Fowler JS, Logan J, Childress AR, Jayne M, Ma Y, Wong C (2006) Cocaine cues and dopamine in dorsal striatum: mechanism of craving in cocaine addiction. *J Neurosci* 26 (24):6583-6588. doi:10.1523/JNEUROSCI.1544-06.2006
- Volkow ND, Wang GJ, Tomasi D, Baler RD (2013) Unbalanced neuronal circuits in addiction. *Curr Opin Neurobiol* 23 (4):639-648. doi:10.1016/j.conb.2013.01.002
- Volkow ND, Wise RA, Baler R (2017) The dopamine motive system: implications for drug and food addiction. *Nat Rev Neurosci* 18 (12):741-752. doi:10.1038/nrn.2017.130
- Voon V, Fernagut PO, Wickens J, Baunez C, Rodriguez M, Pavon N, Juncos JL, Obeso JA, Bezard E (2009) Chronic dopaminergic stimulation in Parkinson's disease: from dyskinesias to impulse control disorders. *Lancet neurology* 8 (12):1140-1149. doi:10.1016/S1474-4422(09)70287-X
- Voon V, Napier TC, Frank MJ, Sgambato-Faure V, Grace AA, Rodriguez-Oroz M, Obeso J, Bezard E, Fernagut PO (2017) Impulse control disorders and levodopa-induced dyskinesias in Parkinson's disease: an update. *Lancet neurology* 16 (3):238-250. doi:10.1016/S1474-4422(17)30004-2
- Walaas I, Fonnum F (1980) Biochemical evidence for gamma-aminobutyrate containing fibres from the nucleus accumbens to the substantia nigra and ventral tegmental area in the rat. *Neuroscience* 5 (1):63-72
- Wall NR, De La Parra M, Callaway EM, Kreitzer AC (2013) Differential innervation of direct- and indirect-pathway striatal projection neurons. *Neuron* 79 (2):347-360. doi:10.1016/j.neuron.2013.05.014
- Wanat MJ, Bonci A (2008) Dose-dependent changes in the synaptic strength on dopamine neurons and locomotor activity after cocaine exposure. *Synapse* 62 (10):790-795. doi:10.1002/syn.20546
- Warren N, O'Gorman C, Lehn A, Siskind D (2017) Dopamine dysregulation syndrome in Parkinson's disease: a systematic review of published cases. *Journal of neurology*,

- neurosurgery, and psychiatry 88 (12):1060-1064. doi:10.1136/jnnp-2017-315985
- Weiss F, Markou A, Lorang MT, Koob GF (1992) Basal extracellular dopamine levels in the nucleus accumbens are decreased during cocaine withdrawal after unlimited-access self-administration. *Brain Res* 593 (2):314-318
- Weiss F, Parsons LH, Schulteis G, Hyytia P, Lorang MT, Bloom FE, Koob GF (1996) Ethanol self-administration restores withdrawal-associated deficiencies in accumbal dopamine and 5-hydroxytryptamine release in dependent rats. *J Neurosci* 16 (10):3474-3485
- White FJ, Kalivas PW (1998) Neuroadaptations involved in amphetamine and cocaine addiction. *Drug Alcohol Depend* 51 (1-2):141-153
- White FJ, Wolf ME (1991) Psychomotor stimulants. In: Pratt, JA

- Ed, *The Biological Basis of Drug Tolerance and Dependence* Academic Press, London:153-197
- Willuhn I, Burgeno LM, Everitt BJ, Phillips PE (2012) Hierarchical recruitment of phasic dopamine signaling in the striatum during the progression of cocaine use. *Proc Natl Acad Sci U S A* 109 (50):20703-20708. doi:10.1073/pnas.1213460109
- Willuhn I, Burgeno LM, Groblewski PA, Phillips PE (2014) Excessive cocaine use results from decreased phasic dopamine signaling in the striatum. *Nat Neurosci* 17 (5):704-709. doi:10.1038/nn.3694
- Wise RA (1978) Catecholamine theories of reward: a critical review. *Brain Res* 152 (2):215-247
- Wise RA (1984) Neural mechanisms of the reinforcing action of cocaine. *NIDA Res Monogr* 50:15-33
- Wise RA (2009) Roles for nigrostriatal--not just mesocorticolimbic--dopamine in reward and addiction. *Trends Neurosci* 32 (10):517-524. doi:10.1016/j.tins.2009.06.004
- Wise RA, Bozarth MA (1987) A psychomotor stimulant theory of addiction. *Psychol Rev* 94 (4):469-492
- Wise RA, Leone P, Rivest R, Leeb K (1995a) Elevations of nucleus accumbens dopamine and DOPAC levels during intravenous heroin self-administration. *Synapse* 21 (2):140-148. doi:10.1002/syn.890210207
- Wise RA, Munn E (1995) Withdrawal from chronic amphetamine elevates baseline intracranial self-stimulation thresholds. *Psychopharmacology (Berl)* 117 (2):130-136
- Wise RA, Newton P, Leeb K, Burnette B, Pocock D, Justice JB, Jr. (1995b) Fluctuations in nucleus accumbens dopamine concentration during intravenous cocaine self-administration in rats. *Psychopharmacology (Berl)* 120 (1):10-20
- Wise RA, Spindler J, deWit H, Gerberg GJ (1978) Neuroleptic-induced "anhedonia" in rats: pimozide blocks reward quality of food. *Science* 201 (4352):262-264
- Wolf ME (1998) The role of excitatory amino acids in behavioral sensitization to psychomotor stimulants. *Prog Neurobiol* 54 (6):679-720
- Wolf ME (2010) Regulation of AMPA receptor trafficking in the nucleus accumbens by dopamine and cocaine. *Neurotox Res* 18 (3-4):393-409. doi:10.1007/s12640-010-9176-0
- Wolf ME (2016) Synaptic mechanisms underlying persistent cocaine craving. *Nat Rev Neurosci* 17 (6):351-365. doi:10.1038/nrn.2016.39
- Wolf ME, Ferrario CR (2010) AMPA receptor plasticity in the nucleus accumbens after repeated exposure to cocaine. *Neurosci Biobehav Rev* 35 (2):185-211. doi:10.1016/j.neubiorev.2010.01.013
- Wolf ME, Tseng KY (2012) Calcium-permeable AMPA receptors in the VTA and nucleus accumbens after cocaine exposure: when, how, and why? *Front Mol Neurosci* 5:72. doi:10.3389/fnmol.2012.00072
- Wolf ME, White FJ, Nassar R, Brooderson RJ, Khansa MR (1993) Differential development of autoreceptor subsensitivity and enhanced dopamine release during amphetamine sensitization. *J Pharmacol Exp Ther* 264 (1):249-255
- Wong DF, Kuwabara H, Schretlen DJ, Bonson KR, Zhou Y, Nandi A, Brasic JR, Kimes AS, Maris MA, Kumar A, Contoreggi C, Links J, Ernst M, Rousset O, Zuckin S, Grace AA, Lee JS, Rohde C, Jasinski DR, Gjedde A, London ED (2006) Increased occupancy of dopamine receptors in human striatum during cue-elicited cocaine craving. *Neuropsychopharmacology* 31 (12):2716-2727. doi:10.1038/sj.npp.1301194

- Wu X, French ED (2000) Effects of chronic delta9-tetrahydrocannabinol on rat midbrain dopamine neurons: an electrophysiological assessment. *Neuropharmacology* 39 (3):391-398
- Xia J, Meyers AM, Beeler JA (2017) Chronic Nicotine Alters Corticostriatal Plasticity in the Striatopallidal Pathway Mediated By NR2B-Containing Silent Synapses. *Neuropsychopharmacology* 42 (12):2314-2324. doi:10.1038/npp.2017.87
- Xia Y, Goebel DJ, Kapatos G, Bannon MJ (1992) Quantitation of rat dopamine transporter mRNA: effects of cocaine treatment and withdrawal. *J Neurochem* 59 (3):1179-1182
- Xu M, Moratalla R, Gold LH, Hiroi N, Koob GF, Graybiel AM, Tonegawa S (1994) Dopamine D1 receptor mutant mice are deficient in striatal expression of dynorphin and in dopamine-mediated behavioral responses. *Cell* 79 (4):729-742
- Yager LM, Garcia AF, Wunsch AM, Ferguson SM (2015) The ins and outs of the striatum: role in drug addiction. *Neuroscience* 301:529-541. doi:10.1016/j.neuroscience.2015.06.033
- Yao WD, Gainetdinov RR, Arbuckle MI, Sotnikova TD, Cyr M, Beaulieu JM, Torres GE, Grant SG, Caron MG (2004) Identification of PSD-95 as a regulator of dopamine-mediated synaptic and behavioral plasticity. *Neuron* 41 (4):625-638
- Yokel RA, Wise RA (1975) Increased lever pressing for amphetamine after pimozide in rats: implications for a dopamine theory of reward. *Science* 187 (4176):547-549
- Zahniser NR, Peris J, Dwoskin LP, Curella P, Yasuda RP, O'Keefe L, Boyson SJ (1998) Sensitization to cocaine in the nigrostriatal dopamine system. *NIDA Res Monogr* 88:55-77
- Zald DH, Boileau I, El-Dearedy W, Gunn R, McGlone F, Dichter GS, Dagher A (2004) Dopamine transmission in the human striatum during monetary reward tasks. *J Neurosci* 24 (17):4105-4112. doi:10.1523/JNEUROSCI.4643-03.2004
- Zapata A, Minney VL, Shippenberg TS (2010) Shift from goal-directed to habitual cocaine seeking after prolonged experience in rats. *J Neurosci* 30 (46):15457-15463. doi:10.1523/JNEUROSCI.4072-10.2010
- Zhang L, Doyon WM, Clark JJ, Phillips PE, Dani JA (2009) Controls of tonic and phasic dopamine transmission in the dorsal and ventral striatum. *Mol Pharmacol* 76 (2):396-404. doi:10.1124/mol.109.056317
- Zhang L, Li J, Liu N, Wang B, Gu J, Zhang M, Zhou Z, Jiang Y, Zhang L, Zhang L (2012) Signaling via dopamine D1 and D3 receptors oppositely regulates cocaine-induced structural remodeling of dendrites and spines. *Neurosignals* 20 (1):15-34. doi:10.1159/000330743
- Zhang XF, Hu XT, White FJ, Wolf ME (1997) Increased responsiveness of ventral tegmental area dopamine neurons to glutamate after repeated administration of cocaine or amphetamine is transient and selectively involves AMPA receptors. *J Pharmacol Exp Ther* 281 (2):699-706
- Zhou FC, Anthony B, Dunn KW, Lindquist WB, Xu ZC, Deng P (2007) Chronic alcohol drinking alters neuronal dendritic spines in the brain reward center nucleus accumbens. *Brain Res* 1134 (1):148-161. doi:10.1016/j.brainres.2006.11.046
- Zhuang X, Oosting RS, Jones SR, Gainetdinov RR, Miller GW, Caron MG, Hen R (2001) Hyperactivity and impaired response habituation in hyperdopaminergic mice. *Proc Natl Acad Sci U S A* 98 (4):1982-1987. doi:10.1073/pnas.98.4.1982

- Zilverstand A, Huang AS, Alia-Klein N, Goldstein RZ (2018) Neuroimaging Impaired Response Inhibition and Salience Attribution in Human Drug Addiction: A Systematic Review. *Neuron* 98 (5):886-903. doi:10.1016/j.neuron.2018.03.048
- Zweifel LS, Fadok JP, Argilli E, Garelick MG, Jones GL, Dickerson TM, Allen JM, Mizumori SJ, Bonci A, Palmiter RD (2011) Activation of dopamine neurons is critical for aversive conditioning and prevention of generalized anxiety. *Nat Neurosci* 14 (5):620-626. doi:10.1038/nn.2808

Figure 1- Simplified schematic representation of the afferent regulation of DA neurons in the VTA and the SNc. Inset – Activity population of DA neurons. About half of VTA neurons are firing spontaneously, and only neurons that fire spontaneously can fire in bursts of action potentials, due to activation of PPTg afferent. Abbreviation: VTA, ventral tegmental area; SNc, substantia nigra pars compacta; SNr, substantia nigra pars reticulata; BLA, basolateral amygdala, vSub, ventral subiculum of the hippocampus; mPFC, medial prefrontal cortex, NAc, nucleus accumbens; VP, ventral pallidum; STN, subthalamic nucleus; PPTg, pedunculo-pontine tegmental nucleus; LDTg, laterodorsal tegmental nucleus; RMTg, rostromedial tegmental nucleus; LHb, lateral habenula.

