



Biodistribution, Imaging and Metabolism Studies of 64-Copper Radiolabelled Monoclonal Antibody : Comparison of DOTA and TE1PA Chelating Agents in a Murine Model of Multiple Myeloma

A. Navarro, T. Le Bihan, Patricia Remaud-Le Saëc, Sébastien Gouard, N. Le
Bris, Clément Bailly, Catherine Sai-Maurel, Benjamin Chalopin,
Jean-François Gustin, Michel Chérel, et al.

► To cite this version:

A. Navarro, T. Le Bihan, Patricia Remaud-Le Saëc, Sébastien Gouard, N. Le Bris, et al.. Biodistribution, Imaging and Metabolism Studies of 64-Copper Radiolabelled Monoclonal Antibody : Comparison of DOTA and TE1PA Chelating Agents in a Murine Model of Multiple Myeloma. *European Journal of Nuclear Medicine and Molecular Imaging*, 2018, 45 (1), pp.S119-S120. inserm-02344286

HAL Id: inserm-02344286

<https://inserm.hal.science/inserm-02344286>

Submitted on 4 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biodistribution, Imaging and Metabolism Studies of 64-Copper Radiolabelled Monoclonal Antibody: Comparison of DOTA and TE1PA Chelating Agents in a Murine Model of Multiple Myeloma

A. Navarro^{1,2}, T. Le Bihan³, P. Le Saëc¹, S. Gouard¹, N. Le Bris³, C. Bailly^{1,2}, C. Sai-Maurel¹, B. Chalopin¹, J. Gestin¹, M. Chérel^{1,4}, R. Tripier³, A. Faivre-Chauvet^{1,2,4}; ¹Inserm 1232 CRCINA, University of Nantes, Nantes, FRANCE, ²University Hospital, Nantes, FRANCE, ³CNRS 6521 / IBSAM, University of Brest, Brest, FRANCE, ⁴Cyclotron Arronax, Saint-Herblain, FRANCE.

Purpose: TE1PA is a C-functionalized monopicolinate cyclam designed for 64-copper chelation. It was proven to have excellent Cu(II) and Cu(I) chelation properties, with fast kinetics, high thermodynamic stability and resistance to transchelation *in vitro*, and a usefulness for phenotypic imagery when coupled to antibodies. This work presents *in vivo* studies (biodistribution, metabolism and imaging) of TE1PA in a syngenic multiple myeloma model on mice, compared to DOTA, its main competitor.

Materials and methods: *p*-SCN-Bn-TE1PA and DOTA-NHS ester were grafted on 9E7.4 rat IgG_{2a} antibody, targeting murine CD-138. Immunoconjugates were then radiolabelled with ⁶⁴Cu. Both ⁶⁴Cu-9E7.4-*p*-SCN-Bn-TE1PA and ⁶⁴Cu-9E7.4-NHS-DOTA were injected in 12 mice previously grafted with 5T33 cells, allowing the development of subcutaneous tumors expressing CD-138. Each mouse was injected with 100 µg of radioimmunoconjugates corresponding to 10 MBq of ⁶⁴Cu. For each time studied (2h, 24h and 48h post-injection), a biodistribution and a liver metabolism studies were conducted on mice for both radiolabelled antibodies. Additionally, a micro-PET scan was performed on 4 mice injected with ⁶⁴Cu-9E7.4-*p*-SCN-Bn-TE1PA at those 3 times. **Results:** Biodistribution study shows an excellent hepatic clearance of the ⁶⁴Cu-9E7.4-*p*-SCN-Bn-TE1PA over time. Significantly higher radioactivity was found in blood, lungs and heart at 48h PI for ⁶⁴Cu-9E7.4-NHS-DOTA, which suggests a release of ⁶⁴Cu from antibody, in parallel with a higher intestinal elimination. This was correlated by the liver metabolism study, which shows a better *in vivo* resistance to transchelation for ⁶⁴Cu-9E7.4-*p*-SCN-Bn-TE1PA. The imaging study of ⁶⁴Cu-9E7.4-*p*-SCN-Bn-TE1PA shows an increasing targeting of the tumors over time and confirms the hepatic clearance at 24h and 48h PI.

Conclusion: ⁶⁴Cu-9E7.4-*p*-NCS-Bn-TE1PA has shown a very high tumor targeting, associated to an increasing of tumor-to-tissues ratio over time that suggests an overall clearance from healthy tissues. Compared to DOTA, TE1PA displays an *in vivo* stability and resistance to transchelation significantly superior. This confirms the previous results obtained *in vitro* and ⁶⁴Cu-labelled TE-1PA proved once again its usefulness for immuno-PET imaging in a preclinical model.