



HAL
open science

Myeloma Cell Self-Renewal Depends on JAG2 Expression and Is Mediated by IGF1 or SCF Loop

David Chiron, Sophie Maiga, Géraldine Descamps, Severine Marionneau-Lambot, Catherine Pellat-Deceunynck

► **To cite this version:**

David Chiron, Sophie Maiga, Géraldine Descamps, Severine Marionneau-Lambot, Catherine Pellat-Deceunynck. Myeloma Cell Self-Renewal Depends on JAG2 Expression and Is Mediated by IGF1 or SCF Loop. ASH meeting 2010, Dec 2010, Orlando, United States. pp.127, 10.1182/blood.V116.21.127.127 . inserm-02459063

HAL Id: inserm-02459063

<https://inserm.hal.science/inserm-02459063v1>

Submitted on 29 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



MYELOMA - BIOLOGY AND PATHOPHYSIOLOGY, EXCLUDING THERAPY I | NOVEMBER 19, 2010

Myeloma Cell Self-Renewal Depends on JAG2 Expression and Is Mediated by IGF1 or SCF Loop

David Chiron, PhD, in, training, Sophie Maiga, Géraldine Descamps, Severine Marionneau, Thilbaud Ouiller, Jerome Moreaux, Bernard Klein, PhD, MD, Philippe Moreau, Jean-Luc Harousseau, Regis Bataille, MD, PhD, Martine Amiot, Catherine Pellat-Deceunynck, PhD



Blood (2010) 116 (21): 127.

<https://doi.org/10.1182/blood.V116.21.127.127>

Split-Screen Share  Tools 

Abstract

Abstract 127

The purpose of this study was to identify the pathways associated with the ability of human myeloma cells (HMCLs) to spontaneous self-renew in a serum-free semi-solid human collagen-based assay. Among 32 HMCLs analyzed, 8 were able to grow spontaneously (from 5% to 35% of seeded cells) without any addition of cytokines or growth factors and this capacity to grow correlated with the presence of *RAS* mutations ($p=0.04$). Gene expression profile analysis of HMCLs identified one gene, *JAG2*, overexpressed in HMCLs that are able to self-renew. Interestingly, flow cytometry analysis of *JAG2* expression showed that the level of membrane *JAG2* expression positively correlated ($r=0.87$) to the percentage of colony formation ($p=0.004$). Blocking Jag-Notch interactions with Notch-Fc chimeric molecules impaired self-colony formation underlying a role for Jag-Notch pathway in colony formation. Furthermore, direct *JAG2* silencing in two independent HMCLs (KMM1 and JJN3) prevented colony formation. Moreover, xenografts in SCID mice showed that *JAG2* silencing fully impaired tumor growth of both KMM1 and JJN3. RT-PCR evaluation of *JAG2* expression showed that 20 of 30 CD138+ purified primary myeloma cells expressed *JAG2* and a Jag2+ subpopulation was identified by flow cytometry within primary CD138+ MM cells of patients at diagnosis or relapse.



We further identified the growth factors involved in the self-renewal. By using blocking anti-IGF1R Ab or C-KIT inhibitor (imatinib mesylate), we showed that self-renewal of HMCLs was dependent on IGF1/IGF1R (5 of 8) or on C-KIT/SCF (1) or on both loops (2 of 8). Of note, C-KIT⁺ HMCLs expressed high JAG2 level at the cell membrane that was decreased by imatinib mesylate. Interestingly, none HMCL self-renewal was dependent on IL6/IL6R loop despite the high efficiency of paracrine IL6 to induce colony formation in most HMCLs. To address expression of *C-KIT/SCF* and *IGF1R/IGF1* in primary myeloma cells, we used public data from patients at diagnosis published by Arkansas University. Expression of *C-KIT* and *IGF1R* was found in 56% and 50% of patients at diagnosis, respectively: 53% of patients express one or the other receptor, 27% express both and 20% express none. Expression of receptors is not similar with regard to the molecular classification of patients as previously shown by cytometry: indeed, MS patients underexpress *C-KIT* ($p < 0.001$) but overexpress *IGF1R* ($p < 0.001$), in full contrast to HY patients who overexpress *C-KIT* ($p < 0.001$) but underexpress *IGF1R* ($p < 0.001$). Moreover, *IGF1R* expression is lower on *C-KIT*⁺ patients as compared with *C-KIT*⁻ ones ($p = 0.049$). Of note, CD-1 and CD-2 patients underexpress both *C-KIT* ($p = 0.039$) and *IGF1R* ($p < 0.001$). IGF1 and SCF are produced by the microenvironment although *IGF1* (but not *SCF*) mRNA was found in myeloma cells too. Altogether, these data suggest that IGF1 and SCF could be the main growth factors for 80% of the patients. Blocking these two tyrosine kinase receptors (in good agreement with their expression in patients) as well as Jag2/Notch interactions could decrease myeloma progression and/or relapse and thus increase survival.

Disclosures:

No relevant conflicts of interest to declare.

Topics: insulin-like growth factor i, myeloma cells, proto-oncogene protein c-kit, insulin-like-growth factor i receptor, growth factor, flow cytometry, imatinib mesylate, interleukin-6, cytokine, gene expression profiling

Author notes

*Asterisk with author names denotes non-ASH members.

© 2010 by The American Society of Hematology

Volume 116, Issue 21

November 19 2010

[Skip to Main Content](#)



Abstract



Abstract Potential Articles of Interest

Hypoxia-Inducible Factor (HIF)-1 α Is a Therapeutic Target in Myeloma-Induced Angiogenesis

Paola Storti et al., Blood, 2011

Galectin-1 Is Highly Expressed By Myeloma Cells and the Bone Marrow Microenvironment and Its Suppression Delineates a New Therapeutic in Vitro and in Vivo Strategy in Multiple Myeloma

Paola Storti et al., Blood, 2014

Beta-Catenin Depended and Independent Effects Induced by Myeloma Cells in Human and Murine Osteoblasts and Osteoblast Progenitors.

Francesca Morandi et al., Blood, 2006

Immune control of hair growth

Lisa D. Chong, Science, 2017

OSU Researchers Report Antitumor Role, Link to p53 for miR-486

GenomeWeb, 2013

Role of IGF1R in PET hybridization imaging of ER positive human breast cancer xenografts using Cu-64 PNA

Kaijun Zhang et al., J Nucl Med, 2007

Powered by



I consent to the use of Google Analytics and related cookies across the TrendMD network (widget, website, blog). [Learn more](#)

Yes

No

[View Metrics](#)

Cited By

Google Scholar

[Skip to Main Content](#)



Abstract



Abstract

Article Activity Alert

Latest Issue Alert



[Current Issue](#)

[First edition](#)

[All Issues](#)

[Collections](#)

[Abstracts](#)

[Video Library](#)

[About Blood](#)

[Subscriptions](#)

[Permissions](#)

[Submit to Blood](#)

[Alerts](#)

[Contact Us](#)

[Authors](#)

[Institutions/Librarians](#)

[Blood Classifieds](#)

[Advertising in Blood](#)

[Terms and Conditions](#)

[Twitter](#)

[Skip to Main Content](#) **American Society of Hematology** / 2021 L Street NW, Suite 900 / Washington, DC 20036 /
TEL +1 202-776-0544 / FAX +1 202-776-0545



Abstract



Abstract

ASH Publications

Blood

Blood Advances

Hematology, ASH Education Program

ASH Clinical News

ASH-SAP

The Hematologist

American Society of Hematology

ASH Home

Research

Education

Advocacy

Meetings

Publications

ASH Store

Copyright ©2020 by American Society of Hematology

[Privacy Policy](#)

[Cookie Policy](#)

[Terms of Use](#)

[Contact Us](#)

