



**HAL**  
open science

## Possible influence of Plasmodium/Trypanosoma co-infections on the vectorial capacity of Anopheles mosquitoes

Maty Fofana, Christian Mitri, Diawo Diallo, Brice Rotureau, Cheikh Tidiane Diagne, Alioune Gaye, Yamar Ba, Constantin Dieme, Mawlouth Diallo, Ibrahima Dia

### ► To cite this version:

Maty Fofana, Christian Mitri, Diawo Diallo, Brice Rotureau, Cheikh Tidiane Diagne, et al.. Possible influence of Plasmodium/Trypanosoma co-infections on the vectorial capacity of Anopheles mosquitoes. BMC Research Notes, 2020, 13 (1), pp.127. 10.1186/s13104-020-04977-8. inserm-02513036

**HAL Id: inserm-02513036**

**<https://inserm.hal.science/inserm-02513036v1>**

Submitted on 20 Mar 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH NOTE

Open Access



# Possible influence of *Plasmodium/Trypanosoma* co-infections on the vectorial capacity of *Anopheles* mosquitoes

Maty Fofana<sup>1</sup>, Christian Mitri<sup>2</sup>, Diawo Diallo<sup>1</sup>, Brice Rotureau<sup>3</sup>, Cheikh Tidiane Diagne<sup>1</sup>, Alioune Gaye<sup>1</sup>, Yamar Ba<sup>1</sup>, Costentin Dieme<sup>2,4</sup>, Mawlouth Diallo<sup>1</sup> and Ibrahima Dia<sup>1\*</sup>

## Abstract

**Objective:** In tropical Africa, trypanosomiasis is present in endemic areas with many other diseases including malaria. Because malaria vectors become more anthro-zoophilic under the current insecticide pressure, they may be exposed to trypanosome parasites. By collecting mosquitoes in six study sites with distinct malaria infection prevalence and blood sample from cattle, we tried to assess the influence of malaria-trypanosomiasis co-endemicity on the vectorial capacity of *Anopheles*.

**Results:** Overall, all animal infections were due to *Trypanosoma vivax* (infection rates from 2.6 to 10.5%) in villages where the lowest *Plasmodium* prevalence were observed at the beginning of the study. *An. gambiae* s.l. displayed trophic preferences for human-animal hosts. Over 84 mosquitoes, only one was infected by *Plasmodium falciparum* (infection rate: 4.5%) in a site that displayed the highest prevalence at the beginning of the study. Thus, *Anopheles* could be exposed to *Trypanosoma* when they feed on infected animals. No *Plasmodium* infection was observed in the *Trypanosoma*-infected animals sites. This can be due to an interaction between both parasites as observed in mice and highlights the need of further studies considering *Trypanosoma/Plasmodium* mixed infections to better characterize the role of these infections in the dynamic of malaria transmission and the mechanisms involved.

**Keywords:** *Trypanosoma*, *Plasmodium*, Mixed infection, *Anopheles*, Senegal

## Introduction

Malaria and African Trypanosomiases also known as Human African Trypanosomiasis (HAT) or sleeping sickness in humans and Animal African Trypanosomiasis (AAT) or Nagana in animals, are endemic diseases in tropical Africa [1–3]. They are respectively caused by protists of the genera *Plasmodium* and *Trypanosoma* that are transmitted to humans and animals by the infective

bites of the blood-feeding insect vectors of the genera *Anopheles* and *Glossina*, respectively. In many parts of tropical Africa, sleeping sickness occurs in areas where malaria is endemic [1]. Although published reports of co-infections are scarce [4–6], *Anopheles* and *Glossina* vectors can theoretically feed on hosts carrying *Plasmodium* and/or *Trypanosoma*, and thus become exposed simultaneously or consecutively to these parasites. We reasoned that such a co-exposure may impact on the development of *Plasmodium* in *Anopheles* vectors and consequently on its transmission.

In Senegal, although no HAT case has been reported for at least one decade [7, 8], AAT remains endemic,

\*Correspondence: Ibrahima.Dia@pasteur.sn

<sup>1</sup> Pôle de Zoologie Médicale, Institut Pasteur de Dakar, 36 Avenue Pasteur, BP 220, Dakar, Sénégal

Full list of author information is available at the end of the article



mainly in the Kédougou region. The last survey from 1971 to assess the distribution of the disease, reported the presence of *T. congolense* and *T. brucei* in cattle, horses, donkeys and carnivores, and the presence of *T. vivax* infections was reported in cattle and horses [9]. As in many African countries, malaria is still a public health problem in Senegal. Although valuable efforts were undertaken to reduce its burden, the Kédougou area remains one with the highest incidence of malaria [10]. Indeed, the ecology of the region is favourable to the maintenance of high seasonal malaria transmission with entomological inoculation rates raising up to 200 infective bites and *Plasmodium* incidence rates greater than 25 per 1000 inhabitants [11].

In a recent study, Sanches-Vaz and others [12] showed through a co-infection model, that mice primarily infected with *T. brucei*, followed by the administration of *P. berghei* sporozoites, are protected from experimental cerebral malaria and presented increased host survival. In addition, subsequent infection of *An. coluzzii* in mice infected with *Trypanosoma brucei brucei* and *Plasmodium yoelii* showed a significant impact of infection as compared to mosquitoes fed on *Plasmodium yoelii* mono-infected mice [13]. These observations raise the question of the possible impact of malaria-trypanosomiasis co-endemicity on the *Plasmodium* transmission by *Anopheles* in the field.

The present study was conducted in the Kédougou region where both malaria and trypanosomiasis are endemic in six study sites of this region selected for their distinct malaria infection prevalence. The objective was to update the last epidemiological data on AAT published more than 40 years ago and further investigate the possible link of *Anopheles* exposure to *Trypanosoma* on their vectorial capacity for the malaria parasite *Plasmodium*.

## Main text

### Methods

#### Selection of the study sites

The study was conducted in the Kédougou region where malaria and AAT co-circulate. This area is situated in a transition zone between the dry tropical forest and the savannah belt. There is one rainy season that lasts from June to November. Mean temperatures vary from 33 to 39.5 °C during the year. The population is predominantly rural and is estimated at 141,226 inhabitants with 55% of people under 20 years [14, 15]. The average population density is estimated at 8 inhabitants per km<sup>2</sup>, mostly living in small-scattered villages. The incidence of malaria in this region is among the highest in Senegal (15 per 1000 inhabitants) with the highest prevalence of *P. falciparum* (14%) among children under 5 years [10]. For trypanosomiasis, no human case has been recorded for at least one

decade. However, AAT is known to be endemic and only a specific cattle race (trypanotolerant Ndama) is adapted in the area. *T. vivax*, *T. congolense* and, in a lesser extent, *T. brucei* are also recorded in neighbouring countries. Based on the data available from 48 different health centers within the Kédougou region [15], six villages were purposively selected based on *P. falciparum* prevalence infection outcomes from outpatients (Fig. 1, Additional file 1). Mako, Bantaco and Tomboronkoto exhibited the lowest *Plasmodium* prevalence (respectively 46.2%, 36.5% and 36.1%) while Silling, Ndébou and Boundoucondi displayed the highest *Plasmodium* prevalence (respectively 73.2%, 73.5% and 75.4%).

#### Blood sample collection in animals

At the beginning of the study in October 2015, before mosquito collection, blood was collected from cattle (bovines, sheeps and goats) in each of the six sites, into a vacuum tube containing EDTA. After collection, the tubes were kept at room temperature for at least 2 h and the serum was recovered after centrifugation and stored at - 80 °C in the laboratory.

#### Mosquito collections

Adult mosquitoes were collected using two sampling methods: (1) CDC miniature light traps hung next to a sleeper under untreated bed nets indoor and outdoor from 7:00 pm to 7:00 am during three consecutive nights; (2) Indoor PSC in October 2015, October 2016 and December 2016 in ten bedrooms in each of the six sites.


Upon collection, mosquitoes were sorted and identified morphologically following the key of Gillies and de Meillon [16]. Bloodmeals from fed specimens collected by PSC were blotted onto a filter paper for host source identification in the laboratory. All mosquitoes were stored individually in numbered tubes with desiccant for laboratory processing.

#### Laboratory processing

The origin of blood meals from fully blood fed mosquito females was identified as human, bovine, ovine, sheep and horse using the ELISA from the procedure of Beier et al. [17].

Females from the *Anopheles gambiae* complex were identified at the species level using the method of Fanello et al. [18].

*Plasmodium* and *Trypanosoma* infections were detected using respectively the nested-PCR of Snounou et al. [19] and the ITS1 "Touchdown" PCR from the procedure of Tran et al. [20].



**Data analysis**

For *Plasmodium* and *Trypanosoma* infection, the infection rates were calculated as the proportion of mosquito specimens or animals found positive by PCR. The anthropophilic/zoophilic rates were calculated as the proportion of human/animal blood to the total blood meals identified by ELISA.

For statistical analysis, Chi square and Fisher exact tests were used with P values < 0.05 considered as significant.

**Results**

**Mosquito collections**

Overall, 252 anopheline specimens were collected during the three sampling sessions. *An. gambiae* s.l. was the predominant species in each of the six villages with a mean frequency of 75% (range 53.8–100%). It was followed by *An. funestus* (13%), *An. rufipes* (6%), *An. flavicosta* (2%), *An. squamosus* (2%), *An. nili* (1.2%), *An. domicola* and *An. pharoensis* (0.4% each).

Within the *An. gambiae* complex, *An. arabiensis* was the predominant species representing 60.4% followed by *An. gambiae* (37.9%) and *An. coluzzii* (1.7%).

**Trophic preferences**

A total of 191 blood meals from *An. gambiae* females were tested by ELISA. Overall, 44% of the identified

blood meals originated from single blood meals (either from human or animal sources). The others were represented by mixed blood meals either from human and animal hosts (34.8%) or from two animal hosts (21.2%).

At the site level, *An. gambiae* were seen to be anthropo-zoophagic in all villages, excepted in Ndébou where they were highly anthropophilic (Table 1).

**Plasmodium and Trypanosoma infections**

In *An. gambiae* s.l. populations, no infection neither by *Trypanosoma* nor by *Plasmodium* parasite was detected by PCR in mosquitoes collected in 2015. However, from a total of 84 specimens collected in 2016, one *An. arabiensis* from the village of Boundoucondi was found harbouring *Plasmodium falciparum* (Table 2). The estimated *P. falciparum* infection rate was 4.5% (95% CI 0.1–22.8%).

Regarding the infectious status of the animals, a total of 284 sera were tested by PCR for *Trypanosoma* detection. Among positive samples, only *Trypanosoma vivax* was detected in 10 animals from 3 different villages (Table 2). The trypanosome infection rates were estimated respectively to 9.1% (95% CI 3–19.9%) in Bantaco, 10.5% (95% CI 2.9–24.8%) in Tomboronkoto and 2.6% (95% CI 0.1–13.8%) in Mako. These villages presented the lowest *Plasmodium* infection rates at the beginning of the study. In Bantaco and Mako, all the infected animals were goats,

**Table 1 Trophic preferences of *An. gambiae* females from the 6 sites visited**

Sites	Tested	Single blood meals		Mixed blood meals		Undet.
		Human	Animal	HA	AA	
Bantaco	8	3 (37.5%)	3 (37.5%)	2 (25%)	0 (0%)	1 (11.1%)
Tomboronkoto	17	10 (58.8%)	1 (5.9%)	2 (11.8%)	4 (23.5%)	0 (0%)
Mako	60	11 (18.3%)	14 (23.3%)	22 (36.7%)	13 (21.7%)	1 (1.6%)
Boundoucondi	46	13 (28.3%)	6 (13%)	17 (37%)	10 (20%)	4 (8%)
Ndébou	16	15 (93.8%)	0 (0%)	0 (0%)	1 (6.3%)	0 (0%)
Silling	37	1 (2.7%)	4 (10.8%)	21 (56.8%)	11 (29.7%)	0 (0%)

HA human-animal, AA animal-animal, Undet. undetermined

**Table 2 *P. falciparum* and *T. vivax* infection rates respectively in *An. gambiae* females and animals in the different villages visited**

Sites	<i>An. gambiae</i> s.l.			Animals		
	<i>P. falciparum</i>			<i>T. vivax</i>		
	Tested	Positive	%	Tested	Positive	%
Bantaco	7	0	0	55	5	9.1 (3–19.9)
Tomboronkoto	11	0	0	38	4	10.5 (2.9–24.8)
Mako	24	0	0	38	1	2.6 (0.1–13.8)
Ndébou	15	0	0	47	0	0
Silling	5	0	0	57	0	0
Boundoucondi	22	1	4.5 (0.1–22.8)	49	0	0

Values in brackets indicate the 95% confidence interval

whereas in Tomboronkoto, three sheep and one goat were found infected. The animal infection rates were significantly different between the 6 villages (Fisher exact test,  $p = 0.005$ ).

## Discussion

The objective of this study was to update the epidemiological profile of AAT in South-East Senegal and to investigate a possible link between *Trypanosoma* infection and *Anopheles* vectorial capacities for malaria.

The presence of *Trypanosoma* was investigated in domestic animals from each of the 6 selected villages. Our results showed the presence of *T. vivax* in ten animals out of 284 in Bantaco, Mako and Tomboronkoto; the three sites that showed the lowest *Plasmodium* infections in human at the beginning of the study. In these sites, no *Plasmodium* infection was observed in mosquitoes. Several studies conducted in the area have shown the presence of trypanotolerant animals [9, 21–23]. The detection of *T. vivax* in animals from the Kédougou region confirms the data already obtained by Touré [9] who showed that this species was mainly found in cattle and horses, not only in this area but also across all the country and also the presence of *T. congolense* and *T.*

*brucei* in cattle, horses, donkeys and carnivores in tsetse-infested areas. A more recent cross-sectional study conducted between June 2011 and September 2012 in Dakar, Sine Saloum, Kédougou and Casamance has revealed the presence of *T. congolense* and *T. brucei* in dogs, donkeys, horses, cattle, sheep, goats with an average infection rate of 3.4% [24]. Concerning *T. vivax*, only one goat from the village of Dielmo was found infected.

The entomological findings showed the predominance of *An. arabiensis* exhibiting variables trophic preferences with mixed blood meals from both human and domestic animals as well as from several animals. *Trypanosoma* parasites were not detected in *Anopheles* mosquitoes. These observations were made from mosquitoes freshly fed on animals or both human and animals. The lack of detection of *Trypanosoma* from mosquito could possibly be due to the short lag of time that the parasite can survive in its organism. Indeed, during their studies Dieme et al, (personal communication) have found that *Trypanosoma* parasites can only survive in the mosquito mid-gut for about 24 h after blood feeding.

*P. falciparum* was detected in the village of Boundoucondi (from 1 *An. arabiensis*) which presented, with Ndébou and Silling, the highest *P. falciparum* infection

rates at the beginning of our study. While infected mosquitoes were observed only in the village of Boundoucondi, the estimated infection rate in this village was higher than that observed by Dia et al, [25] in the area. The absence of *P. falciparum*-infected mosquitoes in the majority of village could be due to the low observed anthropophilic rates combined to the relative important numbers of mixed blood meals on human-animal and animal–animal vertebrate hosts. These results indicate, however, that *Anopheles* mosquitoes could be exposed to *Trypanosoma* parasites when they feed on infected animals. Thus, as observed by Sanches-Vaz and others [12] in mice, this can conduct to a reduction of infection in *Anopheles* mosquitoes. Indeed, the absence of plasmodial infection in the sites where animals were found infected with *T. vivax* (sites with low malaria prevalence) and the presence of plasmodial infections in sites without trypanosome-infected animals, would be in favour of the involvement of trypanosomal infection in the plasmodial infection.

This study confirms the presence of AAT in the South-East region of Senegal and provides indications that *Anopheles* could be exposed to *Trypanosoma* parasites when they feed on infected animals. It highlights the need of further studies taking into account *Trypanosoma* and *Plasmodium* mixed infections to better characterize the role of these infections in the dynamic of malaria transmission and the mechanisms involved.

### Limitations

This work is not an comprehensive and thorough mechanistic study of the interaction of the two parasites and cannot therefore provide a deeper insight into the mechanisms involved. It shows however that *Anopheles* mosquitoes could be exposed to *Trypanosoma* parasites when they feed on infected animals that in turn could reduce *Plasmodium* infection. We would like to investigate this assumption through a study of the impact of *Trypanosoma* infection in the dynamic of malaria transmission and the mechanisms involved experimentally using artificial co-infection with trypanosomes and gametocytes.

### Supplementary information

**Supplementary information** accompanies this paper at <https://doi.org/10.1186/s13104-020-04977-8>.

**Additional file 1.** Localisation of the selected villages in the Kédougou region. Red and yellow colors denote respectively the villages with highest and lowest *P. falciparum* prevalence. The size of the bubble is proportional to the observed prevalences. This map was built using a shapefile from the free domain of the Geographic Information System (<http://www.diva-gis.org>) with the R software version 3.3.1 and the package rgdal.

### Abbreviations

CDC: Centers for Disease Control; PCR: Polymerase chain reaction; HAT: Human African Trypanosomiasis; AAT: Animal African Trypanosomiasis; EDTA: Ethylene diamine tetraacetic acid; PSC: Pyrethrum spray collection; ELISA: Enzyme-Linked Immunosorbent Assay; ITS: Internal transcribed spacer; CI: Confidence interval.

### Acknowledgements

The authors thank Mr Kaly Haya Boubane, Head of the Veterinary Services of Kédougou Region for his support and cooperation in conducting this study and the habitants of the visited villages for their cooperation.

### Authors' contributions

ID, MD, CM and BR conceived and designed the study. DD, CTD, AG and MF performed field studies. MF and ID performed laboratory tests. MF, ID, MD, CM, BR, YB and CD analysed, interpreted the data and wrote the manuscript. All authors read and approved the final manuscript.

### Funding

This work was supported by the Institut Pasteur (PTR 542-2015) through the PTR programme.

### Availability of data and materials

All data generated or analysed during this study are included in this manuscript.

### Ethics approval and consent to participate

The collection of animal blood samples was done by the head of the veterinary services of Kédougou region as part of his routine activities. The owners of the domestic animals were informed about the objectives of this study. Oral informed consents were obtained from them.

### Consent to publish

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup> Pôle de Zoologie Médicale, Institut Pasteur de Dakar, 36 Avenue Pasteur, BP 220, Dakar, Sénégal. <sup>2</sup> Unité Génétique et Génomique des Insectes Vecteurs, Institut Pasteur, Paris, France. <sup>3</sup> Trypanosome Transmission Group, Trypanosome Cell Biology Unit, INSERM U1201 & Department of Parasites and Insect Vectors, Institut Pasteur, Paris, France. <sup>4</sup> Wadsworth Center, New York State Department of Health, Slingerlands, NY, USA.

Received: 29 January 2020 Accepted: 25 February 2020

Published online: 04 March 2020

### References

1. Simarro PP, Cecchi G, Franco JR, Paone M, Diarra A, Ruiz-Postigo A, Fèvre EM, Mattioli RC, Jannin G. Estimating and mapping the population at risk of sleeping sickness. *PLoS Negl Trop Dis*. 2012;6(10):e1859.
2. Rotureau B, Abbeele JVD. Through the dark continent: African trypanosomes development in the tsetse fly. *Front Cell Infect Microbiol*. 2013;3:53.
3. Büscher P, Cecchi G, Jamonneau V, Priotto G. Human African trypanosomiasis. *Lancet*. 2017;390(10110):2397–409.
4. Blum J, Schmid C, Burri C. Clinical aspects of 2541 patients with second stage human African trypanosomiasis. *Acta Trop*. 2006;97(1):55–64.
5. Kuepfer I, Hhary EP, Allan M, Edielu A, Burri C, Blum JA. Clinical presentation of *T.b. rhodesiense* sleeping sickness in second stage patients from Tanzania and Uganda. *PLoS Negl Trop Dis*. 2011;5(3):e968.
6. Kagira JM, Maina N, Njenga J, Karanja SM, Karori SM, Ngotho JM. Prevalence and types of coinfections in sleeping sickness patients in Kenya (2000/2009). *J Trop Med*. 2011. <https://doi.org/10.1155/2011/248914>.
7. WHO. Mapping the distribution of human African trypanosomiasis; 2018. [https://www.who.int/trypanosomiasis\\_african/country/foci\\_AFRO/en/](https://www.who.int/trypanosomiasis_african/country/foci_AFRO/en/). Accessed 16 Apr 2019.

8. Franco JR, Cecchi G, Priotto G, Paone M, Diarra A, Grout L, Mattioli RC, Argaw D. Monitoring the elimination of human african trypanosomiasis: update to 2014. *PLoS Negl Trop Dis*. 2017;11(5):e0005585. <https://doi.org/10.1371/journal.pntd.0005585>.
9. Touré SM. Les trypanosomiasis animales au Sénégal. *Epizootiologie et moyens de lutte*. Bulletin de l'Office International des Epizooties. 1971;76:235–41.
10. Ndiaye Y, Ndiaye JL, Cissé B, Blanas D, Bassene J, Manga IA, Ndiath M, Faye SL, Bocoum M, Ndiaye M, Thior PM, Sene D, Milligan P, Gaye O, Schellenberg D. Community case management in malaria: review and perspectives after four years of operational experience in Saraya district, south-east Senegal. *Malar J*. 2013;12:240.
11. National Malaria Control Programme. National Strategic Framework 2016–2010; 2016. [http://www.pnlp.sn/telechargements/Documents-Strategiques/PNLP\\_CADRE\\_STRATEGIQUE.pdf](http://www.pnlp.sn/telechargements/Documents-Strategiques/PNLP_CADRE_STRATEGIQUE.pdf). Accessed 21 Dec 2018.
12. Sanches-Vaz M, Temporao A, Luis R, Nunes-Cabac H, Mendes AM, Goellner S, Carvalho T, Figueiredo LM, Prudencio M. *Trypanosoma brucei* infection protects mice against malaria. *PLoS Pathog*. 2019;15:e1008145.
13. Dieme C, Zmarlak NM, Brito-Fravallo E, Travaillé C, Pain A, Cherrier F, Genève C, Calvo-Alvarez E, Riehle MM, Vernick KD, Rotureau B, Christian Mitri C. Exposure of anopheles mosquitoes to trypanosomes reduces reproductive fitness and enhances susceptibility to Plasmodium. *PLoS Negl Trop Dis*. 2020;14(2):e000805.
14. Agence Nationale de la Statistique et de la Démographie. Situation économique et sociale régionale 2013. Service Régional de la Statistique et de la Démographie de Sédhiou. <http://www.ansd.sn/ressources/ses/SES-Sedhiou-2013.pdf>. Accessed 11 Apr 2018.
15. Sow A, Loucoubar C, Diallo D, Faye O, Ndiaye Y, Senghor CS, Tal-Dia A, Faye O, Weaver SC, Diallo M, Malvy D, Sall AA. Concurrent malaria and arbovirus infections in Kédougou, southeastern Senegal. *Malar J*. 2016;15:47.
16. Gillies MT, De Meillon B. The Anophelinae of Africa South of the Sahara. *Publ South Afr Ins Med Res*. 1968;54:343.
17. Beier JC, Perkins PV, Wirtz RA, Koros J, Diggs D, Gargan TP II, Koech DK. Bloodmeal identification by direct enzyme-linked immunosorbent assay (ELISA) tested on *Anopheles* (Diptera: Culicidae) in Kenya. *J Med Entomol*. 1988;25:9–16.
18. Fanello C, Santolamazza F, della Torre A. Simultaneous identification of species and molecular forms of *An. gambiae* complex by PCR-RFLP. *Med Vet Entomol*. 2002;16(4):461–4.
19. Snounou G, Viriyakosol S, Jarra W, Thaitong S, Brown KN. Identification of the four human malaria parasite species in field samples by the polymerase chain reaction and detection of a high prevalence of mixed infections. *Mol Biochem Parasitol*. 1993;58:283–92.
20. Tran T, Napier G, Rowan T, Cordel C, Labushagne M, Delespaulx V, Reet NV, Erasmus H, Joubert A, Bücher P. Development and evaluation of an ITS1 "Touchdown" PCR for assessment of drug efficacy against animal African trypanosomiasis. *Vet Parasitol*. 2014;202:164–70.
21. Touré SM. Le parc national du Niokolo-koba (Sénégal). *Mem Ins Fond Afri Noire*. 1969;84:397–400.
22. Hoste CH, Chalon E, d'Ieteren G, Trail JCM. Le bétail trypanotolérant en Afrique occidentale et centrale, Etude FAO production et santé animales. 1988;20/3:217.
23. Diouf MN, Seck MT, Thevenon S, Diop M, Sow RS, Seck MM, Sissoko M, Diop M, Skilton R, Njahira M, Kyalo M, Wanjala B, Mbanjo G, Kaduma E, Nzuki I, Wamonje F, Ndila M, Kemp S. Taurin Ndama au Sénégal: diversité des trois sous populations. *Série fiches techniques ISRA*. 2017;15(3):5.
24. Ravel S, Mediannikov O, Bossard G, Desquesnes M, Cuny G, Davoust B. A study on African animal trypanosomiasis in four areas of Senegal. *Folia Parasitol*. 2015;62:44.
25. Dia I, Diop T, Rakotoarivony I, Kengne P, Fontenille D. Bionomics of *Anopheles gambiae* Giles, *An. arabiensis* Patton, *An. funestus* Giles and *An. nili* (Theobald) (Diptera: Culicidae) and transmission of *Plasmodium falciparum* in a Sudano-Guinean Zone (Ngari, Senegal). *J Med Entomol*. 2003;40(3):279–83.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

