

Global Telemedicine Implementation and Integration Within Health Systems to Fight the COVID-19 Pandemic: A Call to Action

Robin Ohannessian, Tu Anh Duong, Anna Odone

▶ To cite this version:

Robin Ohannessian, Tu Anh Duong, Anna Odone. Global Telemedicine Implementation and Integration Within Health Systems to Fight the COVID-19 Pandemic: A Call to Action. JMIR Public Health and Surveillance, 2020, 6 (2), pp.e18810. 10.2196/18810. inserm-02549310

HAL Id: inserm-02549310 https://inserm.hal.science/inserm-02549310

Submitted on 31 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Viewpoint

Global Telemedicine Implementation and Integration Within Health Systems to Fight the COVID-19 Pandemic: A Call to Action

Robin Ohannessian¹, MD, MPH; Tu Anh Duong^{2,3}, MD, PhD; Anna Odone^{4,5,6}, MD, MPH, PhD

Corresponding Author:

Robin Ohannessian, MD, MPH Télémédecine 360 TLM360 55 Avenue Marceau Paris, 75116 France

Phone: 33 647814410

Email: robin.ohannessian@telemedecine-360.com

Abstract

On March 11, 2020, the World Health Organization declared the coronavirus disease 2019 (COVID-19) outbreak as a pandemic, with over 720,000 cases reported in more than 203 countries as of 31 March. The response strategy included early diagnosis, patient isolation, symptomatic monitoring of contacts as well as suspected and confirmed cases, and public health quarantine. In this context, telemedicine, particularly video consultations, has been promoted and scaled up to reduce the risk of transmission, especially in the United Kingdom and the United States of America. Based on a literature review, the first conceptual framework for telemedicine implementation during outbreaks was published in 2015. An updated framework for telemedicine in the COVID-19 pandemic has been defined. This framework could be applied at a large scale to improve the national public health response. Most countries, however, lack a regulatory framework to authorize, integrate, and reimburse telemedicine services, including in emergency and outbreak situations. In this context, Italy does not include telemedicine in the essential levels of care granted to all citizens within the National Health Service, while France authorized, reimbursed, and actively promoted the use of telemedicine. Several challenges remain for the global use and integration of telemedicine into the public health response to COVID-19 and future outbreaks. All stakeholders are encouraged to address the challenges and collaborate to promote the safe and evidence-based use of telemedicine during the current pandemic and future outbreaks. For countries without integrated telemedicine in their national health care system, the COVID-19 pandemic is a call to adopt the necessary regulatory frameworks for supporting wide adoption of telemedicine.

(JMIR Public Health Surveill 2020;6(2):e18810) doi: 10.2196/18810

KEYWORDS

telemedicine; telehealth; digital health; digital medicine; COVID-19; coronavirus; SARS-CoV-2; public health; surveillance; outbreak; pandemic

On March 11, 2020, the World Health Organization declared the coronavirus disease 2019 (COVID-19) outbreak as a pandemic, with over 720,000 cases reported in more than 203 countries as of 31 March. This announcement followed the declaration of a Public Health Emergency of International Concern (PHEIC) on January 30. The response strategy included early diagnosis, patient isolation, symptomatic monitoring of

contacts, as well as suspected and confirmed cases, and a public health quarantine. The confinement of population and the outbreak impact on health care systems is disrupting routine care for non COVID-19 patients. In this context, telemedicine, particularly video consultations, has been promoted and scaled up to reduce the risk of transmission, especially in the United Kingdom [1] and the United States of America [2,3]. Telemental



¹Télémédecine 360, TLM360, Paris, France

²Dermatology Department, AP-HP, Henri Mondor Hospital, Créteil, France

³Chaire Avenir Sante numérique, équipe 8 IMRB, Inserm, Créteil, France

⁴School of Public Health, Faculty of Medicine, University Vita-Salute San Raffaele, Milan, Italy

⁵Clinical Epidemiology and HTA, IRCCS San Raffaele Scientific Institute, Milan, Italy

⁶Digital Health Section, European Public Health Association, Milan, Italy


health services have been reported in China [4] and Australia as well [5].

Telemedicine was shown to be helpful in previous outbreaks, including former coronavirus outbreaks such as SARS-CoV (severe acute respiratory syndrome—associated coronavirus) and MERS-CoV (Middle East respiratory syndrome coronavirus), or PHEICs related to Ebola and Zika viruses [6,7]. Based on a literature review, the first conceptual framework for telemedicine implementation during outbreaks was published

in 2015 [7]. The framework included tele-expertise, remote patient monitoring of contact cases, and teleconsultation for triage and isolated cases.

An updated framework for telemedicine during the COVID-19 pandemic has been defined in Figure 1. This framework could be applied at a large scale to improve national public health response, and should be shaped on the basis of scientific evidence arising from implemented telemedicine activities.

Figure 1. Conceptual framework of telemedicine for the coronavirus disease 2019 (COVID-19) pandemic. GP: general practitioner; ICU: intensive care unit.



Technological improvements and cost reduction of telemedicine solutions combined with both the high-speed internet and mass spread of smartphones makes it possible to apply this framework and quickly deploy video teleconsultations from a patient's home.

Most countries, however, lack a regulatory framework to authorize, integrate, and reimburse telemedicine in their care delivery for all patients, particularly in emergency and outbreak situations [8]. Two possibilities are currently available for patients: (1) direct-to-consumer telemedicine with private providers mostly relying on out-of-pocket or private insurance payment and (2) free solutions, mainly from US-based companies (for example, WhatsApp, Skype, or Facetime), that may not respect national health data privacy and security requirements. Although these solutions may be useful to support and alleviate the pressure on health care systems during the outbreak, to date, they are mostly unintegrated within national health care systems and not sharing data with public health authorities for epidemiological surveillance.

With the second largest burden of COVID-19 in the world, Italy does not include telemedicine in the essential levels of care granted to all citizens within the National Health Service. No formal input was given on telemedicine by health authorities, despite high pressure on health services during the first phase of the epidemic [9,10]; not until an open call for telemedicine and monitoring system technologies proposals on March 24th was jointly issued by the Ministry for Technological Innovation and Digitalization, the Ministry of Health, the National Institute of Health and the WHO [11].

In France, the Ministry of Health signed a decree on March 9, 2020, allowing the reimbursement of video teleconsultations and tele-expertise by the National Health Insurance (NHI), for patients with COVID-19 symptoms and those confirmed with COVID-19 throughout the country, without the need to know the patient beforehand [12]. The decree was aimed to decrease unnecessary travel for medical consultations, limit the number of individuals grouping in waiting rooms, screen and detect suspected patients, and allow follow-up of mild confirmed cases from home. As the outbreak worsened, temporary funding for follow-up by nurses via video or phone as well as video teleconsultations by midwives (March 19, 2020) and speech therapists (March 25, 2020) was legally allowed.

The pre-existing telemedicine regulations also enabled primary care and hospital doctors to switch scheduled face-to-face patients consultations with known to reimbursed teleconsultations, when suitable. This model was activated in the largest national public academic hospital (AP-HP) in Paris, to encourage mass use of outpatient teleconsultations to reduce patient visits to the hospital (March 13, 2020). This has been reinforced by the High Council of Public Health, which recommended prioritization of teleconsultations for people with risk factors for severe disease in primary care (March 14, 2020) [13], followed by clinical and practical guidelines for patient examination by video consultation published by the Ministry of Health (March 16, 2020) [14]. Between 23 and 29 March, on the second week of national confinement, 486,369 teleconsultations were invoiced to the NHI, representing around 11% of all consults of the week [15]. Among general practitioners, 44% conducted at least one teleconsultation. Until



early March, less than 10,000 teleconsultations a week were invoiced to the NHI.

In this context, several challenges remain for telemedicine to be globally used and integrated into the public health response to COVID-19 and future outbreaks:

- The integration of telemedicine into international and national guidelines for public health preparedness (in keeping with International Health Regulations, 2005) and response [16]
- The definition of national regulations and funding frameworks for telemedicine in the context of public health emergencies
- A strategy to quickly define telemedicine frameworks; use case scenarios; develop clinical guidelines; and standardize triage auto questionnaire and remote patient-monitoring algorithms for any outbreaks at local, national, or global scales

- 4. A strategy and operational plan guiding health care providers to switch to outpatient teleconsultations and increase tele-expertise and remote patient monitoring
- 5. A communication toolkit to inform and educate the population on the recommended use of telemedicine
- 6. A data-sharing mechanism to integrate telemedicine providers' data with epidemiological surveillance
- A scientific evaluation framework and dedicated research funds to describe and assess the impact of telemedicine during outbreaks

All stakeholders are encouraged to address the challenges and collaborate to promote the safe and evidence-based use of telemedicine during the current pandemic and future outbreaks. For countries without integrated telemedicine within their national health care system, the COVID-19 pandemic is a call to adopt the necessary regulatory changes supporting wide adoption of telemedicine.

Acknowledgments

The authors would like to thank Sarina Yaghobian for her contribution and all public and private stakeholders actively involved in the implementation of telemedicine in France during the COVID-19 outbreak.

Conflicts of Interest

None declared.

References

- 1. Greenhalgh T, Wherton J, Shaw S, Morrison C. Video consultations for covid-19. BMJ 2020 Mar 12;368:m998. [doi: 10.1136/bmj.m998] [Medline: 32165352]
- 2. Hollander JE, Carr BG. Virtually Perfect? Telemedicine for Covid-19. N Engl J Med 2020 Mar 11. [doi: 10.1056/nejmp2003539]
- 3. Dorsey ER, Topol EJ. Telemedicine 2020 and the next decade. The Lancet 2020 Mar;395(10227):859. [doi: 10.1016/s0140-6736(20)30424-4]
- 4. Liu S, Yang L, Zhang C, Xiang Y, Liu Z, Hu S, et al. Online mental health services in China during the COVID-19 outbreak. The Lancet Psychiatry 2020 Apr;7(4):e17-e18. [doi: 10.1016/s2215-0366(20)30077-8]
- 5. Zhou X, Snoswell CL, Harding LE, Bambling M, Edirippulige S, Bai X, et al. The Role of Telehealth in Reducing the Mental Health Burden from COVID-19. Telemed J E Health 2020 Mar 23. [doi: 10.1089/tmj.2020.0068] [Medline: 32202977]
- 6. Ohannessian R, Mattos S, Paranhos-Baccalà G, Vanhems P. The role of telemedicine in response to the Zika virus outbreak in Brazil. 2016 Presented at: Zika Summit; April 25, 2016; Paris, France.
- 7. Ohannessian R. Telemedicine: Potential applications in epidemic situations. European Research in Telemedicine/La Recherche Européenne en Télémédecine 2015 Sep;4(3):95-98. [doi: 10.1016/j.eurtel.2015.08.002]
- 8. Smith A, Thomas E, Snoswell C, Haydon H, Mehrotra A, Clemensen J, et al. Telehealth for global emergencies: Implications for coronavirus disease 2019 (COVID-19). J Telemed Telecare 2020 Mar 20:1357633X2091656. [doi: 10.1177/1357633x20916567]
- 9. Paterlini M. On the front lines of coronavirus: the Italian response to covid-19. BMJ 2020 Mar 16;368:m1065. [doi: 10.1136/bmj.m1065] [Medline: 32179517]
- 10. Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? The Lancet 2020 Mar 13. [doi: 10.1016/s0140-6736(20)30627-9]
- 11. Innova per l'Italia. La tecnologia e l'innovazione per la lotta al Coronavirus: Le fast call URL: https://innovaperlitalia.agid.gov.it/call2action/index.html [accessed 2020-03-26]
- 12. Legifrance.gouv.fr. Décret n° 2020-227 du 9 mars 2020 adaptant les conditions du bénéfice des prestations en espèces d'assurance maladie et de prise en charge des actes de télémédecine pour les personnes exposées au covid-19 URL: https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000041704122&categorieLien=id [accessed 2020-03-13]
- 13. Haut Conseil de la santé publique. URL: https://solidarites-sante.gouv.fr/IMG/pdf/new-hcsp-sars-cov-2 patients fragiles v3. pdf [accessed 2020-03-14]
- Lignes directrices pour la prise en charge en ville des patients symptomatiques en phase épidémique de covid-19. 16th March 2020. 2020 Mar 17. URL: https://solidarites-sante.gouv.fr/IMG/pdf/covid-19 doctrine ville v16032020finalise.pdf [accessed 2020-03-17]



- 15. Assurance maladie. 2020 Mar 31. Croissance record du recours à la téléconsultation en mars URL: https://www.ameli.fr/fileadmin/user-upload/documents/20200331 -CP Teleconsultations Covid 19.pdf
- 16. World Health Organization Regional Office for Europe. Strengthening the Health Systems Response to COVID-19. Technical guidance #1: Maintaining continuity of essential health care services while mobilizing the health workforce for COVID-19 response URL: https://euro.sharefile.com/share/view/sbc0659718fd4c8aa [accessed 2020-03-26]

Abbreviations

COVID-19: coronavirus disease 2019

MERS-CoV: Middle East respiratory syndrome coronavirus **PHEIC:** Public Health Emergency of International Concern

SARS-CoV: severe acute respiratory syndrome—associated coronavirus

Edited by G Eysenbach, T Sanchez; submitted 19.03.20; peer-reviewed by C Wu, Y Liu; comments to author 26.03.20; revised version received 27.03.20; accepted 27.03.20; published 02.04.20

Please cite as:

Ohannessian R, Duong TA, Odone A

Global Telemedicine Implementation and Integration Within Health Systems to Fight the COVID-19 Pandemic: A Call to Action

JMIR Public Health Surveill 2020;6(2):e18810 URL: https://publichealth.jmir.org/2020/2/e18810

doi: <u>10.2196/18810</u> PMID: <u>32238336</u>

©Robin Ohannessian, Tu Anh Duong, Anna Odone. Originally published in JMIR Public Health and Surveillance (http://publichealth.jmir.org), 02.04.2020. This is an open-access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Public Health and Surveillance, is properly cited. The complete bibliographic information, a link to the original publication on http://publichealth.jmir.org, as well as this copyright and license information must be included.

