HAL
open science

Inhibitory Plasticity Permits the Recruitment of CA2 Pyramidal Neurons by CA3

Kaoutsar Nasrallah, Rebecca Piskorowski, Vivien Chevaleyre

To cite this version:

Kaoutsar Nasrallah, Rebecca Piskorowski, Vivien Chevaleyre. Inhibitory Plasticity Permits the Recruitment of CA2 Pyramidal Neurons by CA3. eNeuro, 2015, 2 (4), pp.ENEURO.0049-15.2015. 10.1523/ENEURO.0049-15.2015 . inserm-02769407

HAL Id: inserm-02769407

https://inserm.hal.science/inserm-02769407v1

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

This Accepted Manuscript has not been copyedited and formatted. The final version may differ from this version.

eNeuro

Research Article: New Research | Neuronal Excitability

Inhibitory plasticity permits the recruitment of CA2 pyramidal neurons by CA3

Dis-inhibitory plasticity in area CA2

Kaoutsar Nasrallah, Rebecca A. Piskorowski and Vivien Chevaleyre

CNRS UMR8118, Team Synaptic Plasticity and Neural Networks, FR3636, Université Paris Descartes, Sorbonne Paris Cité, 45 rue des Saints-Pères, 75006 Paris, France

DOI: 10.1523/ENEURO.0049-15.2015

Received: 11 May 2015

Revised: 1 July 2015

Accepted: 6 July 2015

Published: 15 July 2015

Funding: Agence Nationale de la Recherche: ANR-12-BSV4-0021-01. Agence Nationale de la Recherche: ANP-13-JSV4-0002. CNRS ATIP programme; Ville de Paris Programme Emergence;

Conflict of Interest: The authors report no conflict of interests

Contributions: KN, RAP, VC designed research, performed experiments, analyzed data; RAP and VC wrote the paper.

Funding sources: We acknowledge the ATIP/Avenir and the Agence Nationale de la Recherche (ANR-12-BSV4-0021-01, ANR-13-JSV4-0002) and Ville de Paris program Emergence for financial support.

Correspondence should be addressed to: Vivien Chevaleyre, vivien.chevaleyre@parisdescartes.fr

Cite as: eNeuro 2015; 10.1523/ENEURO.0049-15.2015

Alerts: Sign up at eneuro.org/alerts to receive customized email alerts when the fully formatted version of this article is published.

Accepted manuscripts are peer-reviewed but have not been through the copyediting, formatting, or proofreading process.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Copyright © 2015 Society for Neuroscience

eNeuro Accepted Manuscript

eNeuro Accepted Manuscript

eNeuro

http://eneuro.msubmit.net

eN-NWR-0049-15R1

Inhibitory plasticity permits the recruitment of CA2 pyramidal neurons by CA3

This is a confidential document and must not be discussed with others, forwarded in any form, or posted on websites without the express written consent of eNeuro.

eNeuro Accepted Manuscript

1. Title : Inhibitory plasticity permits the recruitment of CA2 pyramidal neurons by CA3

2. Abbreviated title : Dis-inhibitory plasticity in area CA2

3. Authors names: Kaoutsar Nasrallah, Rebecca A. Piskorowski and Vivien Chevaleyre

Affiliation : CNRS UMR8118, Team Synaptic Plasticity and Neural Networks, FR3636, Université Paris Descartes, Sorbonne Paris Cité, 45 rue des Saints-Pères, 75006 Paris, France

4. Contributions : KN, RAP, VC : designed research, performed experiments, analyzed data; RAP and VC wrote the paper

5. Correspondence should be addressed to : Vivien Chevaleyre (vivien.chevaleyre@parisdescartes.fr)

6. Number of Figures : 6

7. Number of tables : 0

8. Number of Multimedia : 0

9. Number or words for Abstract : 149

10. Number or words for significance statement: 120

11. Number or words for Introduction : 521

12. Number or words for Discussion : 1678

13. Acknowledgements :

14. Conflict of Interest : A : The authors report no conflict of interests

15. Funding sources: We acknowledge the ATIP/Avenir and the Agence Nationale de la Recherche (ANR-12-BSV4-0021-01, ANR-13-JSV4-0002) and Ville de Paris program Emergence for financial support.

eNeuro Accepted Manuscript

ABSTRACT

Area CA2 is emerging as an important region for hippocampal memory formation. However, how CA2 pyramidal neurons (PNs) are engaged by intra-hippocampal inputs remains unclear. Excitatory transmission between CA3 and CA2 is strongly inhibited and is not plastic. We show in mice that different patterns of activity can in fact increase the excitatory drive between CA3 and CA2. We provide evidence that this effect is mediated by a long-term depression at inhibitory synapses (iLTD), as it is evoked by the same protocols and shares the same pharmacology. In addition, we show that the net excitatory drive of distal inputs is also increased following iLTD induction. The dis-inhibitory increase in excitatory drive is sufficient to allow CA3 inputs to evoke action potential firing in CA2 PNs. Thus, these data reveal that the output of CA2 PNs can be gated by the unique activity-dependent plasticity of inhibitory neurons in area CA2.

SIGNIFICANCE STATEMENT

Long overlooked, recent work has demonstrated that area CA2 of the hippocampus is a critical region for social memory and aggression. How area CA2 integrates in the hippocampal circuit is not understood. While CA2 pyramidal neurons (PNs) receive excitatory input from CA3 PNs, these inputs cannot drive action potentials in CA2 PNs because of a large feed-forward inhibition. Furthermore, CA2 PNs do not express long-term potentiation, so it is unclear if CA2 PNs can be engaged by CA3. We demonstrate that a unique activity-dependent plasticity of CA2 interneurons can increase the excitatory/inhibitory balance between CA3 and CA2 PNs and allow the recruitment of CA2 PNs. Therefore, our results reveal a mechanism by witch CA2 PNs can be engaged by CA3.

eNeuro Accepted Manuscript

50 **INTRODUCTION**

51 Recent findings have revealed that hippocampal area CA2 plays an important role in learning
52 and memory. Area CA2 has been shown to be necessary for social memory formation (Hitti
53 and Siegelbaum, 2014; Stevenson and Caldwell, 2014) and may act to detect discrepancies
54 between stored memories and current sensory information (Wintzer et al., 2014). It has
55 recently been found that the receptive fields of CA2 pyramidal neuron (PN) place cells
56 change rapidly with time, indicating that this region is playing a role in hippocampal learning
57 that may be separate from spatial orientation (Mankin et al., 2015). Neurons in area CA2
58 seem poised to modulate hippocampal activity, given the striking number of receptors that are
59 enriched or uniquely expressed in this area (Jones and McHugh, 2011). A clearer
60 understanding of the connectivity in area CA2 has recently evolved (Chevaleyre and
61 Siegelbaum, 2010; Cui et al., 2012; Rowland et al., 2013; Kohara et al., 2014); however, a
62 better understanding of the plasticity of these connections is critical for understanding how
63 this region contributes to hippocampal learning.

64 Axons from CA3 PNs, i.e. the Schaffer collaterals (SC), connect both CA2 and CA1
65 PNs. However, unlike the well-described SC-CA1 synapses, the SC-CA2 connection does not
66 display activity-dependent long term potentiation (LTP) (Zhao et al., 2007; Chevaleyre and
67 Siegelbaum, 2010) due to the unique expression of calcium buffering proteins (Simons et al.,
68 2009) and of the regulator of G-protein signaling RGS14 (Lee et al., 2010). Furthermore,
69 there is a higher density of interneurons in area CA2 as compared to CA1 (Benes et al., 1998;
70 Andrioli et al., 2007; Piskorowski and Chevaleyre, 2013; Botcher et al., 2014). While
71 inhibition is known to control the strength and plasticity of excitatory transmission in the
72 hippocampus (Chevaleyre and Piskorowski, 2014), the control exerted by inhibition in area
73 CA2 is extensive. When inhibitory transmission is intact, electrical (Chevaleyre and

eNeuro Accepted Manuscript

Siegelbaum, 2010) or selective optogenetic (Kohara et al., 2014) stimulation of the SC inputs results in a very small depolarizing post-synaptic potential (PSP) comprised of both excitatory and inhibitory potentials (EPSPs, IPSPs). The depolarizing EPSP is abruptly truncated by a large hyperpolarizing component due to inhibitory currents from feed-forward inhibition. This very large feed-forward inhibition completely prevents CA3 excitatory inputs from evoking action potentials (APs) in CA2 PNs (Chevaleyre and Siegelbaum, 2010). Following pharmacological block of inhibition, the amplitude of the PSP becomes ~5 fold larger and is sufficient to drive action potentials in CA2 PNs (Chevaleyre and Siegelbaum, 2010; Piskorowski and Chevaleyre, 2013).

Inhibitory transmission from Parvalbumin expressing (PV+) interneurons in area CA2 undergoes a long-term depression (iLTD) mediated by Delta opioid receptor (DOR) activation (Piskorowski and Chevaleyre, 2013). However, it is unknown whether the synapses that undergo iLTD are playing any role in controlling the net strength of SC-CA2 transmission. We show that the balance between excitation and inhibition between CA3 and CA2 is persistently altered following the induction of DOR-mediated iLTD. Furthermore, induction of iLTD by SC stimulation also permits a net increase in the excitation/inhibition ratio at distal CA2 inputs. Finally, this dis-inhibitory mechanism is sufficient to allow SC inputs to drive AP firing in CA2 PNs, thereby engaging CA2 in the tri-synaptic circuit.

MATERIALS AND METHODS

Slice preparation. All animal procedures were performed in accordance with the Author University animal care committee's regulations. 400 μM transverse hippocampal slices were prepared from 6- to 9-week-old C57BL6 male mice. Animals were euthanized under

eNeuro Accepted Manuscript

anesthesia with isofluorane. Hippocampi were removed and placed upright into an agar mold and cut with a vibratome (Leica VT1200S, Germany) in ice-cold extracellular solution containing (in mM): 10 NaCl, 195 sucrose, 2.5 KCl, 15 glucose, 26 $NaHCO_3$, 1.25 NaH_2PO_4, 1 $CaCl_2$ and 2 $MgCl_2$). A cut was made between CA3 and CA2 with a scalpel in some of the slices before being transferred to 30°C ACSF (in mM: 125 NaCl, 2.5 KCl, 10 glucose, 26 $NaHCO_3$, 1.25 NaH_2PO_4, 2 Na Pyruvate, 2 $CaCl_2$ and 1 $MgCl_2$) for 30 min and kept at room temperature for at least 1.5 hr before recording. All experiments were performed at 33°C. Cutting and recording solutions were both saturated with 95% O2 and 5% CO2 (pH 7.4).

Electrophysiological recordings and analysis. Field recordings of PSPs were performed in current clamp mode with a recording patch pipette (3–5 MΩ) containing 1M of NaCl and positioned in the middle of SR or *stratum pyramidale* (SP) in CA1 or CA2. Whole-cell recordings were obtained from CA2 PNs in current clamp mode held at -73 mV with a patch pipette (3–5 MΩ) containing (in mM): 135 KMethylSulfate, 5 KCl, 0.1 EGTA-Na, 10 HEPES, 2 NaCl, 5 ATP, 0.4 GTP, 10 phosphocreatine and 5 µM biocytin (pH 7.2; 280–290 mOsm). Inhibitory currents were recorded with pipette solution containing 135 CsMethylSulfate instead of KMethylSulfate. The liquid junction potential was ~1.2 mV and membrane potentials were corrected for this junction potential. Some recordings were performed with the perforated patch technique. For these experiments, 75 µg/ml gramicidin was added to the intracellular solution along with an additional 4 mM calcium to ensure recordings were acquired in perforated configuration only. Series resistance (typically 12–18 MΩ for whole cell recordings and 65.5 ± 2 MΩ for perforated patch recordings) was monitored throughout each experiment; cells with more than 15% change in series resistance were excluded from analysis. Synaptic potentials were evoked by monopolar stimulation with a patch pipette filled with ACSF and positioned in the middle of CA1 SR or SLM. When

eNeuro Accepted Manuscript

121 evoking PSPs in both SR and SLM in the same experiment, we tested the independence of the
122 inputs prior to the experiment. Using the same stimulation intensity used for the experiment,
123 we compared the PSP amplitude evoked by SR stimulation with the PSP amplitude evoked by
124 SR stimulation 100 ms after SLM stimulation. We also performed the reverse measurement,
125 comparing the PSP amplitude evoked by SLM stimulation to a PSP evoked with SLM
126 stimulation following SR stimulation by 100 ms. The stimulation of the SR and SLM inputs
127 were considered independent if a preceding evoked PSP in the separate pathway had no effect
128 on the amplitude of the second evoked PSP. When axons of CA2 pyramidal neurons were
129 directly recruited by the stimulation pipette, as observed with a back-propagating AP in the
130 recorded CA2 neuron, the stimulating pipette was moved until the direct activation of the
131 axon disappeared.

132
133 A HFS (100 pulses at 100Hz repeated twice), a 10Hz stimulation (100 pulses at 10Hz
134 repeated twice) or DPDPE ([D-Pen^{2,5}]Enkephalin) was applied following 15 – 20 min of
135 stable baseline. Before beginning whole cell experiments, we identified the CA2 PNs by
136 somatic location and size. Furthermore, the cell type was confirmed by several
137 electrophysiological properties (input resistance, membrane capacitance, resting membrane
138 potential, sag amplitude, action potential amplitude and duration). For several experiments,
139 particularly when Cs^+ was in the pipette solution, the slices were fixed following the
140 recording with 4% paraformaldehyde and the neurons were identified by biocytin-streptavidin
141 labeling and immunohistochemistry to label the CA2-specific protein RGS-14.

142 The amplitudes of the PSPs were normalized to the baseline amplitude. The magnitude of
143 plasticity was estimated by comparing averaged responses at 30-40 min for whole cell and at
144 50-60 min for extracellular recordings after the induction protocol with baseline-averaged
145 responses 0–10 min before the induction protocol. All drugs were bath-applied following

eNeuro Accepted Manuscript

dilution into the external solution from concentrated stock solutions. We used Axograph X software for data acquisition and Origin Pro for data analysis. Student's t-test was performed for statistical comparisons and results are reported as mean ± SEM.

RESULTS

Inhibition in area CA2 has been shown to play a very prominent role in controlling the size of the depolarizing component of the compound EPSP-IPSP following stimulation of the SC (Piskorowski and Chevaleyre, 2013). Furthermore, inhibition has been shown to be highly plastic, undergoing an iLTD mediated by DORs (Piskorowski and Chevaleyre, 2013). We asked whether this plasticity of inhibitory transmission might be sufficient to modulate the level of excitatory drive at SC-CA2 synapses. To address this question, we first recorded extracellular post-synaptic potentials (fPSPs) in CA2 *stratum radiatum* (SR) in response to electrical stimulation of SC fibers. These fPSPs are a compound readout of both the local excitatory and inhibitory post-synaptic potentials. After a stable baseline, we applied either a high frequency stimulation (HFS) protocol (100 pulses at 100 Hz repeated twice) or a 10 Hz protocol (100 pulses at 10 Hz repeated twice). These two protocols efficiently induce iLTD of inhibitory inputs in area CA2 (Piskorowski and Chevaleyre, 2013). We found that both the HFS and 10 Hz protocol evoked a lasting increase in the amplitude of the compound fPSP (Figure 1A; with 100Hz: 160.5 ± 4.2% of fPSP amplitude, $p < 0.00001$ $n = 10$; with 10 Hz: Figure 1B; 144.8 ± 9.9% of fPSP amplitude, $p = 0.0034$ $n = 8$). We also observed a significant but smaller increase in the compound fPSP when measuring the slope (136.2 ± 6.4% of fPSP slope, $p = 0.0009$ $n = 8$). We expected a smaller change in the fPSP when measuring the slope, as most of the inhibition evoked by the stimulation is recruited by the SC. The extra synaptic delay of this feed-forward inhibition onto CA2 PNs (compared to the direct SC

eNeuro Accepted Manuscript

170 transmission) will result in a larger control of the peak rather than of the slope of the fPSP.
171 Therefore, the amplitude of the PSP was used for the analysis of the subsequent experiments.

172 To address whether a change in excitability of CA3 PNs was responsible for the increase in
173 fEPSP in CA2, we measured the amplitude of the fiber volley (a measure of the number of
174 axons firing an action potential). As shown in figure 1A and B, we did not detect any
175 significant increase in the fiber volley amplitude after HFS (112.0 ± 5.3%, p = 0.06) or after
176 10 Hz stimulation (105.8 ± 4.1, p = 0.19). In addition, we also applied HFS in slices that had a
177 detached CA3. In this condition, we found a similar increase of the compound fPSP amplitude
178 (Figure 1A, 158.9 ± 6.2% of fPSP amplitude, n = 4, p = 0.0027 p = 0.84 with uncut slices).
179 These data indicate that the long-term increase in the amplitude of the compound fPSP in area
180 CA2 is not due to recurrent activation of CA3 neurons.

181 We also tested the effect of HFS and 10 Hz stimulation in whole cell current clamp recordings
182 of CA2 PNs to ascertain that the increased amplitude of the fPSP was a result of an increased
183 excitatory/inhibitory ratio onto CA2 PNs. We found that both the HFS and 10 Hz protocols
184 triggered a large increase in the PSP recorded in this condition (Figure 1 C,D, 223.4 ± 14.1%
185 after HFS, n = 9; 195.5 ± 26.3% after 10 Hz, n = 6). The larger increase in PSP amplitude
186 observed in whole cell as compared to extracellular recording is not surprising as DORs have
187 been reported to be expressed in interneurons that target PN soma and proximal dendrites
188 (Erbs et al., 2012). A potential iLTD at somatic inputs will likely have a minor contribution in
189 regulating the size of fPSP amplitude recorded in the dendritic area, but will have a large
190 impact on PSP amplitude measured by whole-cell recordings. Consistent with this idea, we
191 found that fPSPs monitored in the somatic region show a larger increase after HFS compared
192 to fPSP monitored simultaneously in the dendritic area (fPSP in soma: 222.3 ± 28.4 % vs

eNeuro Accepted Manuscript

193 fPSP in dendrite: 164.3 ± 7.1 %, n = 5, p =0.047). Thus, our data show different patterns of
194 activity can increase the excitatory drive between CA3 and CA2.

195 **Inhibitory transmission is mandatory for the HFS-induced long-term increase of the SC-**
196 **CA2 PSP.**

197 We hypothesize that the increase in the amplitude of the PSP we measured is the consequence
198 of the previously described DOR-mediated iLTD at inhibitory synapses (Piskorowski and
199 Chevaleyre, 2013). That is, this increase in amplitude is due to a dis-inhibitory mechanism,
200 and not a result of a potentiation of excitatory transmission. If this prediction is true, then the
201 lasting increase in the compound PSP amplitude can only occur if inhibitory transmission is
202 left intact.

203 To test this hypothesis, we recorded extracellular compound fPSPs in areas CA1 and CA2 in
204 response to HFS of SC inputs either in presence or in the absence of $GABA_A$ and $GABA_B$
205 receptor antagonists (1µM SR 95531 and 2µM CGP 55845). In area CA1, the HFS induced a
206 small increase in the fPSP amplitude when inhibition was intact (Figure 2A; 128.2 ± 4.1% of
207 fPSP amplitude, p = 0.0020, n = 5). As expected, this increase was significantly enhanced
208 when the recordings were performed in the continuous presence of GABA receptor
209 antagonists (Figure 2A; 167.9 ± 13.4%, p = 0.0017 and p = 0.0447 with intact inhibition, n =
210 8). In area CA2, a robust and lasting increase of the fPSP was induced with inhibition intact
211 (Figure 2B; 160.5 ± 4.2% of PSP basal amplitude, p < 0.00001 n = 10). Furthermore, unlike
212 what was observed in area CA1, the HFS did not evoke any lasting change in the fPSP
213 amplitude when inhibitory transmission was blocked (Figure 2B; 104.9± 2.3% of EPSP
214 amplitude, p = 0.09 n = 10 and p < 0.0001 with intact inhibition).

eNeuro Accepted Manuscript

We also performed the same experiment using whole-cell current clamp recordings of CA2 PNs. Consistent with the extracellular recordings, the HFS induced an increase in the amplitude of the depolarizing response when inhibition was intact (Figure 2C; 207.4 ± 12.6% of PSP amplitude, $p < 0.00001$, $n = 10$), but not in presence of GABA receptor antagonists (Figure 2C; 103.1 ± 5.7% of EPSP amplitude, $p = 0.6997$ and $p = 0.000006$ with intact inhibition, $n = 7$). Altogether, these results show that in area CA2, intact inhibitory transmission is a requirement for the long-term increase of the compound PSP amplitude.

If the increase in the PSP after HFS indeed results from a decrease in inhibition, then it is possible to make the following prediction: pharmacological block of inhibition should increase the amplitude of the PSP, and part of this increase should be occluded by a previous HFS. In order to test this, we performed current clamp recordings of CA2 PNs and applied GABA receptor blockers following a HFS of SC inputs, or in control experiments with no tetanus. We found that when the GABA receptor blockers were applied without the HFS, the amplitude of the PSP amplitude was strongly increased (Figure 3A; 472.8 ± 85.5% of the PSP amplitude, $p = 0.0026$, $n = 6$), confirming that GABAergic transmission exerts a strong negative control on the PSP amplitude in CA2 PNs (Piskorowski and Chevaleyre, 2013). When the GABA receptor blockers were applied after the HFS, the increase in the PSP amplitude was smaller than the effect of the blockers applied without the HFS. (Figure 3A; 185.8 ± 7.8% of increase of the PSP amplitude, $p< 0.00001$ and $p = 0.0074$ compared to blockers without HFS, $n = 6$). However, the final PSP amplitude after the HFS and GABA blocker application was identical to the amplitude of the PSP after GABA blocker application alone (Figure 3A, $p = 0.7$, $n = 6$). Altogether, these results indicate that the HFS-induced increase of the PSP amplitude is mediated by a decrease in GABAergic transmission.

eNeuro Accepted Manuscript

It was previously reported that the SC-CA2 synapse does not express LTP when inhibition is pharmacologically blocked, but also when inhibition is kept intact (Zhao et al., 2007). One potential explanation for the apparent discrepancy with our results could be the difference of chloride used in our internal solution and that of the previous study. The intracellular [Cl^-] of our pipette solution is 7 mM, making an E_{Cl} of -77 mV. The previous study used 16 mM (Zhao et al., 2007), with an E_{Cl} of -55 mV, which is more depolarized than the resting membrane potential of the cell. With the higher [Cl^-] in the intracellular solution, a decrease in GABAergic transmission would likely not increase the compound PSP amplitude. To test whether the difference in E_{Cl} could explain the discrepancy between our results and those of Zhao *et al*., we performed whole-cell current clamp recordings of CA2 PNs with an internal solution containing 16 mM Cl^-. We found that a HFS did not evoke an increase in the amplitude of the PSP in CA2 PNs when using high concentration of chloride in the pipette even with intact GABAergic transmission (Figure 3B; 111.0 ± 16.9% of PSP basal amplitude, $p = 0.4103$ and $p = 0.006$ with control slices $n = 5$). Finally, in order to directly monitor the compound EPSP/IPSP in pyramidal neurons without affecting intracellular [Cl^-], we performed perforated patch recordings with gramicidin in the pipette. In these conditions, we found that HFS evoked a large increase in the amplitude of the PSP that was not different than the one observed with whole cell recordings (Figure 3C; 200.8 ± 19.9% of baseline, $n = 5$, $p = 0.008$ with baseline, $p = 0.7$ compared to whole cell experiments). Altogether, these data strongly indicate that the dis-inhibitory increase in PSP amplitude can occur during physiological conditions.

The long-term increase in the CA2 compound PSP amplitude requires the activation of delta-opioid receptors

eNeuro Accepted Manuscript

261 Our results show that the potentiation of the PSP amplitude in area CA2 induced by a HFS
262 results from a decrease in inhibition, rather than a direct potentiation of excitatory inputs. If
263 iLTD in area CA2 indeed mediates the increase in the compound PSP, both iLTD and the
264 increase in PSP amplitude should share similar pharmacology. Because the induction of iLTD
265 in area CA2 requires the activation of DORs (Piskorowski and Chevaleyre, 2013), we tested
266 whether HFS-mediated increase in PSP amplitude is also dependent on activation of these
267 receptors.

268 First we performed extracellular and whole cell current clamp recordings of CA2 PNs in
269 response to SC stimulation and applied a HFS in the presence of a DOR competitive
270 antagonist, ICI 74864. We found that the HFS did not significantly change the CA2 PN PSP
271 amplitude if DORs were not activated during the tetanus (Figure 4A; ICI 74864: 119.6 ±
272 12.9% of the PSP amplitude, n = 7, p = 0.135 vs. interleaved control: 215.5 ± 20.6% of the
273 PSP amplitude, p = 0.005, n = 5). Similarly, using extracellular recordings, we found no
274 lasting change in the amplitude of the fPSPs following HFS in the presence of DOR
275 antagonists ICI 74864 or Naltrindol (Figure 4B; 108.7±4.6% of the fPSP basal amplitude, p =
276 0.0822 and p = 0.0006 with absence of DOR antagonist, n = 8). For these experiments, we
277 used two structurally distinct DOR antagonists (ICI 74864 and Naltrindol). Because the
278 increase in PSP amplitude was blocked with both antagonists, data were pooled together.

279 Together, these data show that DOR activation is needed for induction of the lasting increase
280 in CA2 PN PSP amplitude. We then asked whether activation of DORs would be sufficient to
281 mediate this plasticity. To address this question, we performed extracellular recordings in the
282 SR of area CA2 and applied the specific DOR agonist DPDPE for 15 minutes. We found that
283 the application of DPDPE induced a long-term increase of the fPSP amplitude (Figure 4C,
284 162.2 ± 17.7% of baseline, p = 0.02, n = 5). To make sure that this long-term potentiation is

eNeuro Accepted Manuscript

due to a DOR-dependent decrease in inhibition and not to a potential direct increase in SC-CA2 excitatory transmission, we performed the same experiment in the presence of GABA receptor blockers. We found that when inhibitory transmission was blocked, DOR activation did not induce any lasting increase in the PSP amplitude at SC-CA2 synapses (Figure 4C; 99.8 ± 4.2% of baseline, $p = 0.71$ $n = 6$). Altogether, these data show that DOR activation is necessary and sufficient to trigger a lasting increase in the amplitude of the PSP between SC and CA2 PNs.

iLTD evoked by SC input stimulation also alters the excitatory/inhibitory balance at distal inputs of CA2 PNs

CA2 PNs are strongly activated by distal inputs in *stratum lacunosum molecualare* (SLM). Thus, we wondered whether stimulation of proximal SC-inputs, via a dis-inhibitory mechanism, could also influence distal inputs in SLM. We used two stimulating electrodes to evoke PSPs in SC and SLM inputs, as diagramed in figure 5A, and checked for the independence of the two pathways before starting experiments (see methods). We found that tetanic stimulation in SR not only led to a large increase in the PSP amplitude of SC inputs, but also increased the amplitude of PSPs from SLM inputs (Figure 5B; 125.9 ± 5.9% of baseline, $p = 0.0017$ with baseline, $n = 10$). In contrast to SC inputs, SLM inputs have been described to express a LTP that is independent of inhibitory transmission (Chevaleyre and Siegelbaum, 2010). Thus, we wondered whether our observed increase in SLM PSP is a result of a direct LTP at these inputs or from a dis-inhibitory mechanism similar to what we have found at SC-CA2 inputs. We repeated the experiment in the presence of GABA receptor blockers and found that the tetanic stimulation in SR did not result in any increase of SLM-evoked PSPs (Figure 5C; 94.6 ± 7.9% of baseline, $p = 0.52$ with baseline, $n = 8$), consistent with a dis-inhibitory mechanism. We then tested if this dis-inhibitory mechanism at the SLM

eNeuro Accepted Manuscript

309 pathway also depends on DOR activation by applying the SR tetanus in the presence of
310 Naltrindol. We found that the increase in the PSP amplitude of SLM input was completely
311 blocked under these conditions (Figure 5C; 103.9 ± 3.4% of baseline, p = 0.31 with baseline,
312 n = 6).

313 We postulate that this increase in the strength of SLM inputs is a result of the same DOR-
314 dependent dis-inhibition that increases the strength of SC inputs. To test this hypothesis, we
315 directly monitored IPSCs evoked by stimulation in SR and SLM. We found that a tetanus in
316 SR evokes a lasting and significant iLTD in both SC and SLM inputs onto CA2 PNs (Figure
317 5D; 77.7 ± 4.8% of baseline, p = 0.005 for SR IPSCs and 87.0 ± 2.5% of baseline, p = 0.003,
318 for SLM IPSCs, n = 6).

319 **The dis-inhibition at SC-CA2 allows SC inputs to drive firing in CA2 PNs**

320 The large feed-forward inhibition recruited by SC stimulation completely prevents action
321 potential firing in CA2 PNs. However, when inhibition is pharmacologically blocked, SC
322 inputs are able to drive action potential firing in CA2 PNs (Chevaleyre and Siegelbaum,
323 2010). Therefore, we asked whether the decrease in inhibition following iLTD, and the
324 consequent increase in compound PSP amplitude would sufficiently increase the net
325 excitatory drive between CA3 and CA2 PNs to allow SC to drive firing in CA2 PNs.

326 To test this hypothesis, we monitored the magnitude of the population spike (PS) in the
327 somatic layer of area CA2 before and after HFS. We found that before HFS, there was no PS
328 or a PS of very small amplitude measurable only at the highest intensity stimulation (30 V).
329 This observation confirms that the stimulation of CA3 SC axons was not capable of driving
330 the firing of PNs in the CA2 region. Interestingly, a large PS was revealed in CA2 after HFS
331 (Figure 6A, B, C; with 20V stimulation: PS amplitude increases from 0.035 ± 0.024 mV to

eNeuro Accepted Manuscript

332 0.37 ± 0.11mV after HFS, p = 0.01, n = 5; with 30V stimulation: PS amplitude increases from
333 0.09 ± 0.04 mV to 0.59 ± 0.14 mV after HFS, p = 0.006, n = 5). This result shows that the
334 plasticity induced by the HFS is sufficient to allow CA3 inputs to evoke firing of cells in
335 CA2.

336 To determine if the pyramidal cells in area CA2 experience an increased firing probability
337 following HFS of SC, we performed whole-cell current clamp recordings of CA2 PNs and
338 measured the probability of AP firing in response to a series of 5 pulses at 100 Hz at different
339 stimulus intensities before and after HFS. Before HFS, no APs were evoked in CA2 PNs.
340 However, after HFS, CA2 PNs were able to generate APs in response to SC stimulation. Both
341 the number of APs per train and the proportion of cell firing at least one AP during the train
342 were significantly increased after HFS (Figure 6D, E, F; with 30 V stimulation: from 0 to 1.15
343 ± 0.3 AP per train after HFS, p = 0.012, n = 6 and from 0 to 80% of cell firing APs after
344 HFS). To ensure that this increase in firing results from the DOR-mediated plasticity at
345 inhibitory synapses evoked by HFS, we performed the same experiment in presence of the
346 DOR antagonist ICI 174864 (2 µM). When HFS was applied in presence of the drug, no
347 significant change in action potential firing was evoked by HFS (Figure 6G, H, I; with 30V
348 stimulation: from 0 to 0.15 ± 0.15 AP per train after HFS, p = 0.37, n = 5 and from 0 to 20%
349 of cell firing APs after HFS). These results show that the DOR-dependent dis-inhibitory
350 increase in PSP is sufficient to allow CA3 inputs to drive AP firing in CA2 PNs.

351 **DISCUSSION**

352 In this study, we have shown that the net excitatory drive between CA3 and CA2 can be
353 persistently increased in an activity dependent manner, even though the CA3-CA2 excitatory
354 transmission does not express a direct LTP. Our results show that the increase in the
355 excitatory drive between CA3 and CA2 is dependent on inhibition and results from a DOR-

eNeuro Accepted Manuscript

356 mediated iLTD at inhibitory synapses. Furthermore, we show that this dis-inhibition also
357 increases the excitatory drive between SLM inputs and CA2. Finally, this plasticity
358 sufficiently increases the net excitatory drive between CA3 and CA2 to allow SC inputs to
359 drive action potential firing in CA2 PNs.

360 **Activity induces a long-term increase of CA3-CA2 transmission via a dis-inhibitory**
361 **mechanism.**

362 There are several examples of dis-inhibitory plasticity in the hippocampus. For instance, a
363 pairing protocol induces a shift in GABA reversal potential, thereby increasing synaptic
364 strength at CA3-CA1 synapses (Ormond and Woodin, 2009). Similarly, a pairing protocol
365 between distal and proximal inputs strongly facilitates CA3-CA1 synapses, an effect
366 dependent on a decrease in feed-forward inhibition recruited by SC synapses (Basu et al.,
367 2013). In both cases, the decrease in inhibition facilitates a plasticity that can occur
368 independently of a change in inhibition. In our study, the increase in excitatory/inhibitory
369 balance between CA3 and CA2 is entirely dependent on a dis-inhibition mechanism. Indeed,
370 SC-CA2 excitatory synapses do not express LTP after HFS or other induction protocols (Zhao
371 et al., 2007; Simons et al., 2009; Chevaleyre and Siegelbaum, 2010; Lee et al., 2010).
372 However, we found that HFS or 10Hz stimulation could trigger a lasting increase in the net
373 excitatory drive between CA3 and CA2. Several experiments indicate that this increase is
374 dependent on GABAergic transmission and results from a dis-inhibitory mechanism. First, the
375 lasting increase in PSP amplitude was absent when inhibitory transmission was
376 pharmacologically blocked. Second, increasing the reversal potential for Cl^- in the recorded
377 cell also prevented induction of this plasticity. Third, the increase in the amplitude of the PSP
378 after application of GABA receptor blockers was partially occluded by previous HFS. Fourth,
379 the activity-dependent increase in PSP shares the same pharmacology as the iLTD, i.e., a

eNeuro Accepted Manuscript

strict dependence on DOR activation. And finally, application of a DOR agonist, which induces a lasting decrease in inhibition but does not alter isolated excitatory transmission, was sufficient to evoke the lasting increase in PSP amplitude.

Our results are in agreement with several studies showing a lack of plasticity at SC-CA2 synapses when inhibition was blocked. Nonetheless, they seem to differ from a previous study where no plasticity was evoked at SC synapses with inhibitory transmission left intact (Zhao et al., 2007). We believe that the recording conditions likely account for this apparent discrepancy. While the previous study used a high concentration of Cl^- in the recording solution (16 mM), we used 7 mM of Cl^-. In accordance with this idea, we found no increase in PSP amplitude with 16 mM Cl^- in the recording pipette. The precise physiological concentration of intracellular Cl^- in CA2 PNs is unknown. However, the fact that we observe an increase in PSP amplitude when intracellular Cl^- is not perturbed (using gramicidin perforated patch recordings) and when both intracellular ionic composition and resting membrane potential are not affected (using extracellular recordings), strongly indicates that the dis-inhibitory-mediated increase in PSP amplitude can occur in an intact system. Furthermore, we are confident that the fPSP we record in CA2 is not contaminated by PSP in CA1 because we do not observe LTP in CA2 in presence of GABA blockers, while a large potentiation is observed with simultaneous fPSP recordings in CA1.

The dis-inhibitory increase in CA3-CA2 transmission is dependent on DOR

Our results strongly indicate that the DOR-mediated iLTD recently described (Piskorowski and Chevaleyre, 2013) is involved in the dis-inhibitory increase in PSP between CA3 and CA2. First, the dis-inhibitory potentiation was abolished in presence of two different selective antagonists of DORs (ICI 74864 and Naltrindol). Second, direct activation of DORs with the selective agonist DPDPE was sufficient to trigger a lasting increase in the PSP amplitude.

eNeuro Accepted Manuscript

404 Therefore, these data show that DORs are necessary and sufficient to increase the PSP
405 amplitude between CA3 and CA2.

406 Opioids have long been known to increase excitability in the hippocampus and to facilitate the
407 induction of plasticity at excitatory synapses. For instance, activation of mu opioid receptors
408 (MORs) and DORs is required for LTP induction at the perforant path – granual cell synapse
409 in the dentate gyrus (Bramham and Sarvey, 1996) and activation of MORs facilitates LTD
410 induction at the SC-CA1 PN synapse (Wagner et al., 2001). In both cases, the observed
411 increase in excitatory drive is believed to result from a transient decrease in inhibition during
412 the induction protocol, hence facilitating the induction of the plasticity at excitatory inputs.

413 Our results differ in several ways from the previously reported action of opioids. First, in
414 contrast to the transient effect of DORs on GABAergic transmission in CA1, activation of
415 DORs was shown to induce a lasting depression of inhibition in area CA2 (Piskorowski and
416 Chevaleyre, 2013). In addition, in contrast to the well-described facilitatory action of opioids
417 on plasticity at excitatory synapses, our results show that DOR activation is mandatory for the
418 change in excitatory/inhibitory balance between CA3 and CA2.

419 Our results show that stimulation in SR increases synaptic strength of both proximal and
420 distal inputs. The increase in the strength of excitatory transmission was dependent on
421 GABAergic transmission. There are several reports of hetero-synaptic plasticity between
422 proximal and distal inputs onto hippocampal PNs. For example, the potentiation of SC inputs
423 onto CA1 PNs following theta-burst stimulation of distal dendritic inputs (Han and
424 Heinemann, 2013), and potentiation of SC inputs and depression of distal dendritic inputs
425 following low frequency stimulation of distal inputs (Wöhrl et al., 2007). To our knowledge,
426 there is no report of a lasting increase in synaptic strength at distal dendritic input that is
427 induced by stimulation in the proximal dendritic region. The simplest explanation for our

eNeuro Accepted Manuscript

428 results is that stimulation in SR evokes iLTD, and both SC and SLM stimulation recruit the
429 depressed inhibitory inputs. In agreement with this idea are the findings that PV+ basket cells
430 in area CA2 have dendrites that traverse SR and SLM (Mercer et al., 2012; Tukker et al.,
431 2013). Furthermore, it was recently shown that synaptic transmission from PV+ interneurons
432 in area CA2 is depressed during DOR-mediated iLTD (Piskorowski and Chevaleyre, 2013).
433 Together, our results highlight a new mechanism by which activity in SR can enhance the net
434 excitatory drive of synapses in SLM. This interplay is likely to have important consequences
435 on information transfer by extrahippocampal (SLM) and intrahippocampal (SC) inputs onto
436 CA2 PNs.

437

438 **Physiological and pathological consequences of the dis-inhibitory action of DOR in CA2**

439 When inhibition is intact, SC EPSPs are very small and fail to induce AP in CA2 PNs. Our
440 results show that following the induction of DOR-mediated iLTD, stimulation of SC inputs
441 can evoke APs in CA2 PNs. This was observed both in whole-cell and extracellular
442 recordings, indicating that it can occur when the resting potential of CA2 PNs is not affected.
443 This result is consistent with a previous study using voltage sensitive dyes to study
444 information propagation through the hippocampus that reported that dentate gyrus stimulation
445 resulted in either a fast propagation of activity between CA3 and CA1, or a slower
446 propagation successively recruiting CA3, CA2 and CA1 PNs (Sekino et al., 1997).
447 Interestingly, blockade of GABA receptors allowed recruitment of CA2 from slices in which
448 activation of this region was initially not detected. Our results provide a mechanism for the
449 result observed in this previous study and show that the dis-inhibitory-dependent recruitment
450 of CA2 can be evoked in an activity-dependent manner. While HFS (100 Hz) might not be
451 very physiological, the activity-dependent increase in PSP amplitude can also be evoked with

eNeuro Accepted Manuscript

452 a 10 Hz stimulation, a frequency that falls within the range of theta oscillations. Furthermore,
453 it was previously shown that the depression of inhibitory transmission underlying the increase
454 in PSP can be evoked with 100 and 10 Hz, but also with a theta burst stimulation protocol.
455 Therefore, we think it is likely that physiological patterns of activity will be efficient to
456 trigger the increase in excitatory drive between CA3 and CA2, thus allowing CA2 to be
457 recruited by CA3. We propose that inhibition in area CA2 is acting as a gate to control
458 information flow between CA3 and CA2, and DOR activation can be considered as one of the
459 keys to open this gate.

460 While our results suggest that an activity-dependent decrease in inhibition could be
461 relevant for information transfer by allowing SC inputs to activate CA2 PNs, it is likely that a
462 more global decrease in inhibition in CA2 is detrimental for the hippocampus. For instance, a
463 decrease in the density of PV+ interneurons has been reported to occur uniquely in area CA2
464 during schizophrenia (Knable et al., 2004). Similarly, the density of PV+ cells is considerably
465 reduced in CA2 during epilepsy (Andrioli et al., 2007) and physiological recordings in the
466 hippocampi of human epileptic patients revealed an important decrease (Wittner et al., 2009)
467 or a complete loss of inhibition onto CA2 PNs (Williamson and Spencer, 1994). Our results
468 provide a potential explanation for why a persistent decrease in the inhibitory gate in CA2
469 could alter the tri-synaptic circuit during schizophrenia (Benes, 1999), and why CA2 might be
470 the locus of epileptiform activity generation both in rodents (Knowles et al., 1987) and in
471 humans (Wittner et al., 2009).

472 In summary, our results show that while there is no LTP at the SC-CA2 PN synapse, the
473 excitatory/inhibitory balance can be persistently shifted toward excitation in an activity-
474 dependent manner through a dis-inhibition mechanism. Interestingly, this dis-inhibition
475 results in an increase in net excitatory drive at both SC inputs as well as distal dendritic

eNeuro Accepted Manuscript

476 inputs. Furthermore, our results reveal that the decrease in inhibitory transmission sufficiently
477 increases the excitatory drive from SC inputs to allow recruitment of CA2 by CA3 PNs. Thus,
478 our data add complexity to the hippocampal circuitry and reveal how CA2 PNs can be
479 engaged by intra-hippocampal inputs following activity.

480

481

482 **REFENCES**

483 Andrioli A, Alonso-Nanclares L, Arellano J, DeFelipe J (2007) Quantitative analysis of
484 parvalbumin-immunoreactive cells in the human epileptic hippocampus. Neuroscience
485 149:131–143.

486 Basu J, Srinivas KV, Cheung SK, Taniguchi H, Huang ZJ, Siegelbaum SA (2013) A cortico-
487 hippocampal learning rule shapes inhibitory microcircuit activity to enhance hippocampal
488 information flow. Neuron 79:1208–1221.

489 Benes FM (1999) Evidence for altered trisynaptic circuitry in schizophrenic hippocampus.
490 Biol Psychiatry 46:589–599.

491 Benes FM, Kwok EW, Vincent SL, Todtenkopf MS (1998) A reduction of nonpyramidal cells
492 in sector CA2 of schizophrenics and manic depressives. Biol Psychiatry 44:88–97.

493 Botcher NA, Falck JE, Thomson AM, Mercer A (2014) Distribution of interneurons in the
494 CA2 region of the rat hippocampus. Front Neuroanat 8:104.

495 Bramham CR, Sarvey JM (1996) Endogenous activation of mu and delta-1 opioid receptors is
496 required for long-term potentiation induction in the lateral perforant path: dependence on

eNeuro Accepted Manuscript

497 GABAergic inhibition. J Neurosci 16:8123–8131.

498 Chevaleyre V, Piskorowski RA (2014) Modulating excitation through plasticity at inhibitory
499 synapses. Front Cell Neurosci 8.

500 Chevaleyre V, Siegelbaum SA (2010) Strong CA2 pyramidal neuron synapses define a
501 powerful disynaptic cortico-hippocampal loop. Neuron 66:560–572.

502 Cui Z, Gerfen CR, Young WS (2012) Hypothalamic and other connections with the dorsal
503 CA2 area of the mouse hippocampus. J Comp Neurol.

504 Erbs E, Faget L, Scherrer G, Kessler P, Hentsch D, Vonesch J-L, Matifas A, Kieffer BL,
505 Massotte D (2012) Distribution of delta opioid receptor-expressing neurons in the mouse
506 hippocampus. Neuroscience 221:203–213.

507 Han EB, Heinemann SF (2013) Distal dendritic inputs control neuronal activity by
508 heterosynaptic potentiation of proximal inputs. Journal of Neuroscience 33:1314–1325.

509 Hitti FL, Siegelbaum SA (2014) The hippocampal CA2 region is essential for social memory.
510 Nature 508:88–92.

511 Jones MW, McHugh TJ (2011) Updating hippocampal representations: CA2 joins the circuit.
512 Trends Neurosci 34:526–535.

513 Knable MB, Barci BM, Webster MJ, Meador-Woodruff J, Torrey EF, Stanley
514 Neuropathology Consortium (2004) Molecular abnormalities of the hippocampus in
515 severe psychiatric illness: postmortem findings from the Stanley Neuropathology
516 Consortium. Mol Psychiatry 9:609–20–544.

517 Knowles WD, Traub RD, Strowbridge BW (1987) The initiation and spread of epileptiform

eNeuro Accepted Manuscript

518 bursts in the in vitro hippocampal slice. Neuroscience 21:441–455.

519 Kohara K, Pignatelli M, Rivest AJ, Jung H-Y, Kitamura T, Suh J, Frank D, Kajikawa K, Mise
520 N, Obata Y, Wickersham IR, Tonegawa S (2014) Cell type-specific genetic and
521 optogenetic tools reveal hippocampal CA2 circuits. Nature Neuroscience 17:269–279.

522 Lee SE, Simons SB, Heldt SA, Zhao M, Schroeder JP, Vellano CP, Cowan DP, Ramineni S,
523 Yates CK, Feng Y, Smith Y, Sweatt JD, Weinshenker D, Ressler KJ, Dudek SM, Hepler
524 JR (2010) RGS14 is a natural suppressor of both synaptic plasticity in CA2 neurons and
525 hippocampal-based learning and memory. Proc Natl Acad Sci U S A 107:16994–16998.

526 Mankin EA, Diehl GW, Sparks FT, Leutgeb S, Leutgeb JK (2015) Hippocampal CA2
527 Activity Patterns Change over Time to a Larger Extent than between Spatial Contexts.
528 Neuron 85:190–201.

529 Mercer A, Eastlake K, Trigg HL, Thomson AM (2012) Local circuitry involving
530 parvalbumin-positive basket cells in the CA2 region of the hippocampus. Hippocampus
531 22:43–56.

532 Ormond J, Woodin MA (2009) Disinhibition mediates a form of hippocampal long-term
533 potentiation in area CA1. Tell F, ed. PLoS One 4:e7224.

534 Piskorowski RA, Chevaleyre V (2013) Delta-Opioid Receptors Mediate Unique Plasticity
535 onto Parvalbumin-Expressing Interneurons in Area CA2 of the Hippocampus. Journal of
536 Neuroscience 33:14567–14578.

537 Rowland DC, Weible AP, Wickersham IR, Wu H, Mayford M, Witter MP, Kentros CG
538 (2013) Transgenically Targeted Rabies Virus Demonstrates a Major Monosynaptic
539 Projection from Hippocampal Area CA2 to Medial Entorhinal Layer II Neurons. Journal

eNeuro Accepted Manuscript

540 of Neuroscience 33:14889–14898.

541 Sekino Y, Obata K, Tanifuji M, Mizuno M, Murayama J (1997) Delayed signal propagation
542 via CA2 in rat hippocampal slices revealed by optical recording. Journal of
543 Neurophysiology 78:1662–1668.

544 Simons SB, Escobedo Y, Yasuda R, Dudek SM (2009) Regional differences in hippocampal
545 calcium handling provide a cellular mechanism for limiting plasticity. Proc Natl Acad Sci
546 USA 106:14080–14084.

547 Stevenson EL, Caldwell HK (2014) Lesions to the CA2 region of the hippocampus impair
548 social memory in mice. Eur J Neurosci.

549 Tukker JJ, Lasztóczi B, Katona L, Roberts JDB, Pissadaki EK, Dalezios Y, Márton L, Zhang
550 L, Klausberger T, Somogyi P (2013) Distinct dendritic arborization and in vivo firing
551 patterns of parvalbumin-expressing basket cells in the hippocampal area CA3. Journal of
552 Neuroscience 33:6809–6825.

553 Wagner JJ, Etemad LR, Thompson AM (2001) Opioid-mediated facilitation of long-term
554 depression in rat hippocampus. J Pharmacol Exp Ther 296:776–781.

555 Williamson A, Spencer DD (1994) Electrophysiological characterization of CA2 pyramidal
556 cells from epileptic humans. Hippocampus 4:226–237.

557 Wintzer ME, Boehringer R, Polygalov D, McHugh TJ (2014) The Hippocampal CA2
558 Ensemble Is Sensitive to Contextual Change. Journal of Neuroscience 34:3056–3066.

559 Wittner L, Huberfeld G, Clémenceau S, Eross L, Dezamis E, Entz L, Ulbert I, Baulac M,
560 Freund TF, Maglóczky Z, Miles R (2009) The epileptic human hippocampal cornu

eNeuro Accepted Manuscript

ammonis 2 region generates spontaneous interictal-like activity in vitro. Brain 132:3032–3046.

Wöhrl R, Haebler von D, Heinemann U (2007) Low-frequency stimulation of the direct cortical input to area CA1 induces homosynaptic LTD and heterosynaptic LTP in the rat hippocampal-entorhinal cortex slice preparation. Eur J Neurosci 25:251–258.

Zhao M, Choi Y-S, Obrietan K, Dudek SM (2007) Synaptic plasticity (and the lack thereof) in hippocampal CA2 neurons. J Neurosci 27:12025–12032.

FIGURE LEGENDS

Figure 1: HFS and 10 Hz stimulations induce a long-term increase of SC-CA2 PSP amplitude.

Time course of the average normalized PSP amplitude obtained by extracellular recording **(A, B)** in CA2 SR or whole-cell current clamp recording **(C, D)** of CA2 PNs, in response to SC stimulation. Both a HFS protocol (two sets of 100 pulses at 100 Hz; A: $p < 0.00001$, $n = 10$; C: $p = 0.00018$, $n = 9$) and a 10 Hz protocol (two sets of 100 pulses at 10 Hz; B: $p = 0.0009$, $n = 8$; D: $p = 0.01596$, $n = 6$) induce a long-term increase in the SC PSP amplitude in CA2. The fiber volley (FV, a measure of the number of axons firing an action potential) was not significantly increased after HFS (A; $p = 0.06$) or 10Hz stimulation (B; $p = 0.19$). Panel A also shows that making a cut between CA3 and CA2 does not affect the magnitude of the potentiation evoked by HFS ($p = 0.59$ with uncut slices, $n = 5$). Upper right-hand corner in all panels: averaged PSP traces of a representative experiment corresponding to time points before (a) and 60 minutes (b) (A, B) or 40 minutes (C, D) after the stimulation protocol. Error

eNeuro Accepted Manuscript

584 bars show SEM in all panels.

585

586 **Figure 2: The HFS-induced long-term potentiation of the PSP in CA2 is dependent on**
587 **GABAergic transmission.**

588 **(A)** Time course of average normalized fPSP amplitude recorded in CA1 SR in response to
589 SC stimulation, showing that the HFS-induced increase in PSP amplitude in control
590 conditions (open circles, $p = 0.0020$, $n = 5$) was facilitated in continuous presence of the
591 GABA A and B receptor antagonists, 1μM SR 95531 and 2μM CGP 55845 (filled circles, $p =$
592 0.0017, $n = 8$). **(B, C)** In CA2, HFS does not trigger long-lasting increase in the SC PSP
593 amplitude in continuous presence of GABA receptor antagonists (filled circles, B:
594 extracellular recordings, $p = 0.09$, $n = 10$; C: whole cell recording, $p = 0.6997$, $n = 10$), but
595 evokes a large and lasting increase in the PSP amplitude in control experiments (open circles,
596 B: extracellular recordings, $p < 0.00001$, $n = 10$; C: whole cell recording, $p < 0.00001$, $n =$
597 10). In all panels, averaged PSP traces corresponding to the time points before (a) and after
598 (b) HFS are shown on the right. Error bars show SEM in all panels.

599

600 **Figure 3: HFS induces a long-term increase of the PSP amplitude in CA2 via a dis-**
601 **inhibition mechanism.**

602 **(A)** Two representative examples of normalized PSP time course from CA2 PN whole-cell
603 recordings illustrating how a HFS partially occludes the effect of GABAA and GABAB
604 receptor blocker application on PSP amplitude. Top right: PSP traces corresponding to the
605 time points before (a), after HFS (b) and after application of GABA receptor blockers (c),
606 with or without HFS. Bottom right: summary histograms showing the percentage of increase
607 in the PSP amplitude induced by HFS applied alone (1), GABA receptor blocker application

eNeuro Accepted Manuscript

608 after HFS (2), HFS plus GABA blockers (1+2) and GABA blockers applied without previous
609 HFS (3). **(B)** Normalized CA2 PN SC PSPs recorded with either 7 mM (open circles) or 16
610 mM (closed circles) Cl- in the pipette solution. A HFS (arrow at time 0) fails to induce a long-
611 lasting increase PSP amplitude when high concentration of chloride is used in the pipette
612 solution (filled circled, p = 0.4103, n = 5) but evokes normal long-term potentiation in control
613 experiments (open circles, p = 0.0047, n = 5). (C) Summary graph of experiments performed
614 using the gramicidin-perforated patch-recording configuration. HFS triggers an increase in
615 PSP amplitude (p = 0.008, n = 5) similar to the one observed using whole-cell recording
616 configuration.
617
618 **Figure 4: The increase in SC PSP amplitude in CA2 PNs is dependent on DOR**
619 **activation.**
620 **(A)** A HFS does not trigger a long-term increase of the PSP amplitude recorded in CA2 PNs
621 in the presence of 2 μM of the DOR competitive antagonist, ICI 174864 (filled circles, n = 7,
622 p = 0.135) but induces a normal long-term increase in the PSP magnitude in interleaved
623 control experiments (open circles, p = 0.005, n = 5). **(B)** Time course of normalized fPSP
624 amplitude recorded in CA2 SR showing how the application of a DOR antagonist (ICI
625 174864 or Naltrindol 0.1μM) during HFS (filled circles, p = 0.0822 and p = 0.0006 with
626 absence of ICI 74864, n = 8) prevented the induction of a lasting increase of the fPSP
627 amplitude observed in absence of drug application (open circle). Averaged PSP traces
628 corresponding to the time points before (a) and after (b) HFS performed in control conditions
629 (top) or in the presence of ICI 174864 (bottom) are shown on the right. **(C)** Application of 0.5
630 μM of a DOR-selective agonist (DPDPE) is sufficient to induce a long lasting increase in the
631 fPSP amplitude recorded in SR of CA2 in the absence (open circles, p = 0.02, n = 5) but not
632 in the presence (filled circles, p = 0.55679, n = 6) of GABAA and GABAB receptor blockers.

eNeuro Accepted Manuscript

633 Right, example fPSP traces corresponding to the time points before (a) and after (b)
634 application of DPDPE in absence (top) or in continuous presence (bottom) of the GABAA
635 and GABAB receptor blockers. Error bars show SEM in all panels.
636
637 **Figure 5: Stimulation in SR induces a heterosynaptic iLTD and increases distal**
638 **excitatory drive onto CA2 PN.**

639 **(A)** Cartoon illustrating the arrangement of the stimulating recording electrodes in SR and
640 SLM. **(B)** Average PSP amplitudes of SR (open circles) and SLM (closed circles) inputs
641 following HFS stimulation in SR. Note that both SR and SLM inputs are potentiated
642 following the HFS ($p = 0.00035$ for SR inputs, $p = 0.0017$ for SLM inputs, $n = 10$), but only
643 SR inputs show a rapid post-tetanic increase in amplitude. Top, averaged PSP traces
644 corresponding to the time points before (a) and after (b) HFS. **(C)** The increase in distally
645 evoked PSP following stimulation in SR was blocked by GABAA and GABAB receptor
646 blockers (open circles, $p = 0.52$, $n = 8$) and by the DOR antagonist naltrindol (grey circles, p
647 $= 0.31$, $n = 6$). **(D)** Average amplitude of IPSCs evoked by stimulation in SR and SLM
648 following HFS in SR. Note that both inputs express an inhibitory LTD following HFS in SR
649 ($p = 0.006$ for SR inputs, $p = 0.003$ for SLM inputs, $n = 6$).
650
651 **Figure 6: HFS in SR allows CA3 inputs to evoke action potential firing in CA2 PNs.**

652 **(A)** Traces of extracellular recordings in the CA2 pyramidal layer in response to a 20V
653 stimulation of SC inputs, before (grey traces) and after a HFS (black traces), illustrating how
654 HFS induces an increase in the population spike (PS) amplitude (negative peak). **(B)** Average
655 PS amplitude as a function of stimulation intensity before (open circle) and after (filled circle)
656 HFS (with 20V stimulation: $p = 0.01$, $n = 5$; with 30V stimulation: $p = 0.006$, $n = 5$). **(C)**

eNeuro Accepted Manuscript

657 Time course of average normalized PS amplitude recorded in CA2 pyramidal layer in
658 response to a 20V stimulation of SC inputs, showing a long lasting increase in PS amplitude
659 after HFS (p=0.01, n=5). **(D)** Traces of whole-cell current clamp recordings in a CA2
660 pyramidal cell in response to a train of stimulations (5 pulses at 100Hz) of SC inputs,
661 illustrating how APs can be evoked in CA2 neurons after (black traces) but not before HFS
662 (grey traces). **(E)** Average number of APs per train at different stimulation intensities,
663 showing how a 5 pulse train at 100Hz of the SC inputs does not trigger APs in CA2 PNs
664 before HFS (open circles) but induces APs after HFS (filled circles, with 30 V stimulation:
665 from 0 to 1.15 ± 0.3 AP per train after HFS, p = 0.012, n = 6). **(F)** A HFS increases the
666 percentage of CA2 PNs firing at least one AP during the train (with 30V stimulation: from 0%
667 to 80% of cell firing APs). **(G)** Traces of whole-cell current clamp recordings in a CA2 PN in
668 response to a stimulation train (5 pulses at 100Hz) of SC inputs in the presence of the DOR
669 antagonist ICI 174864 (2μM), illustrating how application of ICI 174864 prevents the
670 induction of APs in CA2 PNs after HFS (black traces). **(H)** Average number of APs per train
671 at different stimulation intensities in the presence of the DOR antagonist before (open circles,
672 with 0 to 30V stimulations: 0 AP per train) and after HFS (filled circles, with 30 V
673 stimulation: 0.15 ± 0.15 AP per train after HFS, p = 0.37, n = 5). **(I)** In presence of ICI
674 174864 HFS does not induce a large increase of the percentage of CA2 PNs firing at least one
675 AP during the train (with10V stimulation: 0% of cell firing APs before and after HFS; with
676 20V and 30V stimulation: from 0% to 20% of cell firing APs). Error bars show SEM in all
677 panels.

eNeuro Accepted Manuscript



eNeuro Accepted Manuscript



eNeuro Accepted Manuscript



eNeuro Accepted Manuscript



eNeuro Accepted Manuscript



eNeuro Accepted Manuscript