



Association of Partial Chromosome 3 Deletion in Uveal Melanomas With Metastasis-Free Survival

Manuel Rodrigues, Khadija Ait Rais, Flore Salviat, Nathalie Algret, Fatoumata Simaga, Raymond Barnhill, Sophie Gardrat, Vincent Servois, Pascale Mariani, Sophie Piperno-Neumann, et al.

► To cite this version:

Manuel Rodrigues, Khadija Ait Rais, Flore Salviat, Nathalie Algret, Fatoumata Simaga, et al.. Association of Partial Chromosome 3 Deletion in Uveal Melanomas With Metastasis-Free Survival. JAMA Ophthalmology, 2020, 138 (2), pp.182. 10.1001/jamaophthalmol.2019.5403 . inserm-03037871

HAL Id: inserm-03037871

<https://inserm.hal.science/inserm-03037871>

Submitted on 3 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Association of Partial Chromosome 3 Deletion in Uveal Melanomas with
Metastasis-free Survival**

Manuel Rodrigues^{1,2*}, M.D. Ph.D., Khadija Ait-Rais³, Flore Salviat⁴, M.D., Nathalie
Algret⁴, Fatoumata Simaga³, Raymond Barnhill^{5,6}, M.D., Sophie Gardrat^{1,5}, M.D.,
Vincent Servois⁷, M.D., Pascale Mariani⁸, M.D., Sophie Piperno-Neumann², M.D.,
Sergio Roman-Roman⁹, Pharm.D. Ph.D., Olivier Delattre^{1,3}, M.D. Ph.D., Nathalie
Cassoux^{6,10}, M.D. Ph.D., Alexia Savignoni⁴, M.D. Ph.D., Marc-Henri Stern^{1,3}, M.D.
Ph.D., Gaëlle Pierron³, Ph.D.

1 - Unit 830 « Cancer, heterogeneity, instability and plasticity » INSERM, Institut
Curie, PSL Research University, Paris, 75248, France

2 - Department of Medical Oncology, Institut Curie, PSL Research University, Paris,
75248, France

3- Department of Genetics, Institut Curie, PSL Research University, Paris, 75248,
France

4 - Department of Biometry, Institut Curie, PSL Research University, Paris, 75248,
France

5- Department of Biopathology, Institut Curie, PSL Research University, Paris,
75248, France

6- Faculty of Medicine, University of Paris Descartes, Paris, 75006, France

7- Department of Medical Imaging, Institut Curie, PSL Research University, Paris,
75248, France

8- Department of Surgical Oncology, Institut Curie, PSL Research University, Paris,
75248, France

9- Department of Translational Research, Institut Curie, PSL Research University,
Paris, 75248, France

10- Department of Ocular Oncology, Institut Curie, PSL Research University, Paris,
75248, France

***Correspondence to:** Dr. Manuel Rodrigues, Institut Curie 26 rue d'Ulm, 75248
Paris Cedex 05. email: manuel.rodrigues@curie.fr Tel: +33144324672 Fax:
+33153104041

Word count: 2,729

38 **Key Points**

39 **Question:** What is the association of partial chromosome 3 deletion in uveal
40 melanomas with metastasis-free survival?

41 **Findings:** In this retrospective study, partial deletions of chromosome 3
42 encompassing the *BAP1* locus were associated with a lower metastasis-free survival
43 at 60 months compared to uveal melanomas without such deletion.

44 **Meaning:** These findings suggest that uveal melanomas carrying a partial deletion of
45 chromosome 3 encompassing the *BAP1* locus have a poor prognosis.

46

Abstract

Importance

Studies on uveal melanomas (UMs) demonstrated the prognostic value of 8q gain and monosomy 3, but the prognosis of UMs with partial deletion of chromosome 3 remains to be defined.

Objective

To determine the association of partial chromosome 3 deletion in uveal melanomas with metastasis-free survival.

Design

Retrospective cohort of consecutive comparative genomic hybridization arrays from May 2006 to July 2015.

Setting

Monocentric study in a referral center.

Participants

Patients presenting with UMs with and without partial loss of chromosome 3.

Main Outcomes and Measures

Metastasis-free survival and overall survival at 60 months.

Results

Of the 1,088 consecutive comparative genomic hybridization arrays that were performed, 43 UMs (4%) carried partial deletions of chromosome 3. Median follow-up was 66 months. Metastasis-free survival at 60 months was 34% (95% confidence interval [CI], 15.8 to 71.4) for UMs carrying a deletion of the *BAP1* (*BRCA1 associated protein-1*) locus (*BAP1del*; 24 tumors) and 81% (95% CI, 64.8 to 100) for UMs without the loss of the *BAP1* locus (*BAP1 normal*; *BAP1nl*; 19 tumors; log-rank *p*-value = .001). Overall survival at 60 months was 65% (95% CI, 43.5 to 95.8) *versus*

84% (95% CI, 69.0 to 100) in the BAP1del and the BAP1nl groups, respectively (log-rank p-value < .001). In these 43 cases, metastasis-free survival at 60 months was 100% for UMs without loss of the *BAP1* locus or 8q gain, 70% (95% CI, 50.5 to 96.9) for UMs carrying one of these alterations and 13% for those carrying both (95% CI, 2.1 to 73.7; log-rank p-value < .001). Similarly, overall survival at 60 months was 100%, 81% (95% CI, 63.3 to 100) and 47% (95% CI, 23.3 to 93.6) in these three groups, respectively (log-rank p-value < .001).

Conclusions and Relevance

These findings suggest that partial deletion of chromosome 3 encompassing the *BAP1* locus is associated with poor prognosis. A cytogenetic classification of UMs could be proposed based on the status of the *BAP1* locus instead of chromosome 3, locus, while also taking chromosome 8q into account.

Introduction

Uveal melanoma (UM) is the most common primary malignant ocular tumor in adults of European ancestry¹. Despite efficient treatment, up to 50% of the patients will eventually develop metastases²⁻⁴. Reliable prognostic assessment allows a closer monitoring of high-risk patients. Pathological prognostic factors include large tumor basal diameter, thickness, ciliary body involvement, extraocular extension, epithelioid cell histology, high mitotic rate and lymphocytic infiltration⁵. The gene expression profile DecisionDx-UM (GEP; Castle Biosciences, Friendswood, TX), based on the expression level of 12 genes, is frequently used in North America to complete the prognostic assessment^{6,7}.

In the early 1990s, recurrent cytogenetic aberrations including monosomy 3 (M3), gain of 6p and 8q were identified in UM samples⁸. In 1996, M3 was empirically shown to be a robust prognostic factor⁹. Since then, genomic arrays have become routine tools to refine pathological prognosis along with the GEP. We previously refined the prognostic value of M3 and gain of 8q by defining three groups: (i) high-risk patients whose tumors present a M3 and an 8q gain with a 2-year metastasis-free interval (2y-MFI) of 37%; (ii) intermediate-risk with either a M3 or an 8q gain (2y-MFI: ~85%) and (iii) low-risk with neither M3 nor 8q gain (2y-MFI: ~100%)¹⁰.

The most common hypothesis to explain the poor prognosis of M3 tumors is the presence of one or more tumor suppressor genes (TSG) on chromosome 3. *BAP1* (*BRCA1 associated protein-1*), a TSG located on the 3p21.1 cytoband, is now established as a main actor of UM malignant transformation as it is frequently mutated in M3 tumors and germline mutations are associated with UM predisposition¹¹⁻¹⁶. However, all or most *BAP1*-mutated UMs intriguingly present a M3 (or a loss of heterozygosity of the whole chromosome 3 due an isodisomy)

110 suggesting that the role of chromosome 3 loss in UM tumorigenesis may not be
111 restricted to *BAP1* inactivation. Therefore, prognostication of UM samples with partial
112 deletions of chromosome 3, as sometimes observed in our daily practice and by
113 other authors, is problematic¹⁷. The goals of the present study were to explore these
114 UMs with partial deletions of chromosome 3, as assessed by comparative genomic
115 hybridization (array-CGH), in order to assess their prognosis and to determine the
116 minimal region of deletion associated with poor prognosis.

Materials and methods

Patients

This study was approved by our institutional ethics committee. Written informed consent for the use of tissues and data for research was signed by each patient. The study complied with the principles of the Declaration of Helsinki. All patients were referred to our institution and followed up by our physicians. Clinical diagnosis of uveal melanoma was based on the presence of typical clinical findings as previously described¹⁰. Local treatment consisted of proton beam radiotherapy, iodine 125 brachytherapy or enucleation, depending on the size and location of tumors. Tumor samples were obtained by enucleation, endoresection or fine-needle aspiration at the time of clip or plaque positioning. Liver ultrasound, liver magnetic resonance imaging or body computed tomography were performed at diagnosis and every 6 months afterwards. Diagnosis of metastasis was systematically confirmed by a biopsy.

Genomic analysis

Tumor DNA was extracted and processed as previously described¹⁰. Array-CGH was performed on three different platforms according to the period when the test was performed: bacterial artificial chromosome arrays as previously described¹⁸, NimbleGen 4×72 K arrays (Roche NimbleGen, Madison, Wisconsin, USA) and Agilent 180K CGH/LOH custom chip (Santa Clara, California, USA). Array-CGH were interpreted by three of the authors (MR, KAR, GP). Partial deletion of chromosome 3 was defined as the loss of at least one region of chromosome 3, but not the totality, whatever its size and location. Genomic positions in this article are defined in hg18 human genome assembly.

Statistical analysis

Clinical, pathological and genomic data at diagnosis and follow-up events (local and distant recurrences, second cancers, death from UM or from any other cause) were collected. The French Death Registry was consulted for patients lost to follow-up. The metastasis-free survival (MFS) at 60 months was defined as the proportion of patients alive and free of metastasis at 60 months of follow-up after local treatment of primary UM. The overall survival (OS) at 60 months was defined as the proportion of patients alive at 60 months of follow-up after local treatment of primary UM, whatever the cause of death. Survival distributions were estimated by the Kaplan–Meier method and compared using the log-rank test. All tests were bilateral and performed with a significant level of 5%. In order to identify variables associated with MFS, a Cox regression analysis of candidate prognostic factors was performed using a forward stepwise selection procedure. The added value of each variable to the Cox model was determined using a likelihood ratio test with a significant level of 5%. Statistical analysis was performed using R software V3.3 (<http://www.r-project.org/>).

Results

We prospectively re-analyzed the array-CGH profiles in 1,088 UMs which had been processed between May 2006 and July 2015, and detected 43 cases (4.0%) harboring a partial deletion of chromosome 3 (eTable 1 in the supplement). Median follow-up in these 43 cases was 66 months (range: 1.2-126.2 months). Median age was 58 years-old (range 12-79), median tumor diameter was 16 millimeters (range: 10-22) and median thickness was 10 millimeters (range: 5.3-18.2). Ciliary body and optic nerve were involved in 33% (14/43) and in 9% (4/43) of cases, respectively. Cell morphology was epithelioid or mixed in 30% of cases (13/43). Primary tumors were treated by enucleation in 42% (18/43) of cases. MFS and OS at 60 months were 61% (95% confidence interval [CI], 46 to 79.7) and 76% (95% CI, 62.8 to 92.3), respectively. A global overview of copy number profiles is provided in eFigure 1 in the supplement. Size of deletions ranged from 1.36 to 110.88 megabases.

We first explored survival data in an unsupervised manner and observed three recurrently lost regions in at least eight metastatic samples: (i) from 3pter to p22.2, (ii) from 3p22.1 to p14.2 and (iii) from 3q13.2 to q24 (Figure 1). Of these, two regions were more frequently lost in metastatic cases than in non-metastatic ones: the 3pter-p22.2 region (8/13 *versus* 6/30 cases, respectively; $p=.013$; odds ratio [OR]=6.1; 95% CI, 1.2 to 34.1) and the 3p22.1-p14.2 region, which encompasses *BAP1* (10/13 *versus* 9/30 cases, respectively; $p=.007$; OR=7.4; 95% CI, 1.5 to 51.8). These two regions were close and highly correlated between each other, as 8 out of 10 metastatic cases presenting a 3p22.1-p14.2 loss also presented a 3pter-p22.2 loss. The 3p22.1-p14.2 region carries 290 other genes beside *BAP1*, but no recurrent mutations of these 290 genes were found in public and in-house databases^{12,19,20}.

We then hypothesized that *BAP1* loss was the main driver of poor prognosis in M3. To explore this hypothesis, we compared tumors with a chromosome 3 partial deletion encompassing the *BAP1* locus (24 tumors; BAP1del) and tumors with a chromosome 3 partial deletion not encompassing the *BAP1* locus (19 tumors; BAP1nl). Tumors carrying a loss of the *BAP1* locus frequently showed large losses of the short arm of chromosome 3 (Figure 2). MFS at 60 months was 81% (95% CI, 64.8 to 100) for the BAP1nl genomic group and 34% (95% CI, 15.8 to 71.4) for the BAP1del group (Figure 3; $p=.001$). OS at 60 months was 84% (95% CI, 69.0 to 100) for the BAP1nl genomic group and 65% (95% CI, 43.5 to 95.8) for the BAP1del group ($p<.001$). The only variables associated with MFS in univariate analysis were loss of the *BAP1* locus and gain of 8q. These two variables independently contributed to MFS in multivariate analysis (Table 1).

We defined four groups depending on the *BAP1* locus (lost/not lost) and 8q (gained/not gained) statuses. Prognoses of the *BAP1* locus lost/8q normal and *BAP1* locus not lost/8q gained were similar so we merged these two groups, as in our previous classification (eFigure 2 in the supplement)¹⁰. By analogy with our previous work, we defined three prognosis groups as follows: (i) a group at low risk of metastasis without loss of the *BAP1* locus or 8q gain (9 cases), (ii) an intermediate risk group with tumors carrying either loss of the *BAP1* locus (7 cases) or 8q gain (15 cases) and (iii) a high risk group with loss of the *BAP1* locus and 8q gain (12 cases). MFS at 60 months were 100%, 70% (95% CI, 50.5 to 96.9) and 13% (95% CI, 2.1 to 73.7) for the low-, intermediate- and high-risk groups, respectively (Figure 4; $p<.001$).

205 OS at 60 months were 100%, 81% (95% CI, 63.3 to 100) and 47% (95% CI, 23.3 to
206 93.6) for the low-, intermediate- and high-risk groups, respectively ($p<.001$).

207

Discussion

In this work, we explored a relatively large series of UMs with partial deletion of chromosome 3 and showed that loss of the *BAP1* locus is likely to explain the poor prognosis of M3 UM. This result was obtained by two different approaches investigating indirectly the prognostic value of the most frequently deleted regions of chromosome 3 and then directly assessing the prognostic value of the loss of the *BAP1* locus in this series. The first consequence is to provide a potentially more accurate estimation of the prognosis of UMs presenting a partial deletion of chromosome 3. Our classification suggested efficiency in predicting metastatic outcome, identifying a group with a very good MFS with no recurrence and a group with a high risk of 92% of recurrences with a median follow-up of more than 5 years. Survival rates were close to what we observed in a previous series of UMs presenting either a M3 or a disomy 3, associated or not with 8q gain¹⁰. This hypothesis has yet to be verified in subsequent studies because direct comparison could not be done here.

Other teams are using different genomic technologies to assess UM prognosis. Fluorescence *in situ* hybridization (FISH) is widely used but it may miss the loss of the *BAP1* locus if the probe is not centered on this gene, as observed in several publications^{2,21-24}. Furthermore, FISH is often performed without chromosome 8q assessment leading to suboptimal prognosis estimation. Multiplex ligation-dependent probe amplification (MLPA) assay covering the *BAP1* locus is a good alternative to characterize recurrent genomic imbalances in UM but MLPA, as well as FISH and array-CGH, only evaluate copy number and, consequently does not identify isodisomic cases^{25,26}. GEP is a transcriptomic prognosis assay that is widely used in

United States⁷. This assay distinguishes two subsets of UMs either at low or high risk of metastasis by assessing the expression of 12 genes, including four that are located on the short arm of chromosome 3 (*EIF1B*, *LMCD1*, *ROBO1*, *SATB1*) and one on the 3q (*FXR1*). Underexpression of these genes, possibly due to M3, is associated with poor prognosis. A more accurate prediction by GEP is possible by adding the expression of *PRAME*, a gene located on an instable region of chromosome 22 exposed to duplication, which was correlated to the 8q status in the pivotal paper⁶. To our knowledge, GEP has never been specifically tested in a large series of UMs with partial chromosome 3 deletions. Furthermore, GEP has never been compared to the combined M3/8q signature in a large cohort, impeding any conclusion on the superiority of one modality on the other. BAP1 immunohistochemistry is an alternative way to assess the prognosis of UMs^{27,28}. However, immunohistochemistry for BAP1 does not correlate in all cases to the BAP1 mutational status in UM, and is therefore not a perfect surrogate²⁷.

In the present series, partial deletions of chromosome 3 were found in 4% of cases, which is comparable with some previous series²⁹⁻³¹ but lower than others^{17,32,33}. Recruitment bias may explain part of this discrepancy but it is most probably explained by the variety of technologies, as well as the different classifications that were used. Comparison of all these studies is therefore limited. Similarly, the prognosis of these tumors was not clear as a discrepancy was observed with some series associating partial loss with good prognosis^{17,29,32} while others associated it with intermediate or poor prognosis^{25,33,34}. These differences may be explained by the absence of distinction depending on the loss of the *BAP1* locus compared to other losses.

258

259 One explanation for the low MFS associated with the loss of this locus may be that
260 the loss of one *BAP1* allele contributes to the inactivation of this gene and
261 subsequent aggressiveness of the tumor. However, the minimal region of deletion we
262 found associated with the lowest MFS in our series (3p22.1-p14.2) includes 291
263 genes. Even though this region encompasses *BAP1*, it cannot be excluded that other
264 important genes are present there and that haploinsufficiency of these genes affects
265 tumorigenesis. The two alleles of a TSG are commonly inactivated in the two-hit
266 model by a combination of different mechanisms, including total or partial loss of a
267 chromosome, deleterious point mutations, short insertions/deletions, large-scale
268 insertions/deletions and promoter methylations³⁵. It is highly intriguing that, *BAP1*
269 inactivation is so frequently associated with monosomy 3 in UM, contrary to renal
270 clear-cell carcinomas and mesotheliomas, which rather carry losses of the short arm
271 of chromosome 3 only or deleterious mutations of both alleles¹⁶. Furthermore,
272 haploinsufficiency of other genes on chromosome 3, possibly on its long arm, may
273 play a role on UM tumorigenesis. This hypothesis may be of particular interest and
274 should be put in perspective with the recent discovery of *MBD4* (3q21.3) recurrent,
275 inactivating mutations in UM³⁶⁻³⁸.

276

277 There is, for now, no standard treatment in the metastatic setting, but new drugs are
278 being developed in UM³⁹. When an efficient treatment will be available, the following
279 step will consist in testing this treatment in the adjuvant setting in high-risk patients⁴⁰.
280 Accurate prognosis evaluation is essential for such trials and assays able to assess
281 the status of the *BAP1* locus and 8q status may then be required. Next-generation
282 sequencing appears to be the best option in the near future as it not only assesses

copy number, heterozygosity and mutational statuses of UMs at low cost and with a lower amount of DNA, but also allow to follow circulating tumor DNA⁴¹⁻⁴³. Moving towards the implementation of such technologies in our daily practice will allow ocular oncology to enter the modern age of precision medicine while reducing costs and refining UM prognosis.

Limitations

The conclusions of this work are limited by its retrospective nature, but prospective series are unrealistic given the rarity of such tumors. Instead, the present work provides evidence to refine the current UM genomic classification, which may help ophthalmologists to better predict the metastatic evolution of their patients. Before generalization, other series from different centers are required. Furthermore, one could argue that our series, composed of large tumors (median diameter of 16mm and median thickness of 10mm) is not reflecting the overall population of UM patients, particularly as larger UMs are known to host a greater frequency of genomic alterations, including 8q gains^{37,44}. Other centers have reported genomic studies on biopsies of smaller UMs⁴⁵. However, this procedure is not consensual and must not be undertaken in inexperienced ocular oncology centers because of potential surgical complications. Multicenter collaborative studies of small UM genomics are required to address the question of partial chromosome 3 loss frequency at this stage of primary UM development. Another limitation of this study is that the array-CGH technology is not adapted to detect chromosome 3 isodisomy, an infrequent alteration in UM, probably associated with poor prognosis. SNP-array can resolve this issue but, in the future, next-generation sequencing will probably be the privileged technology to circumvent this issue. More importantly, although the *BAP1* locus hypothesis is a

logical hypothesis, we cannot definitely affirm that *BAP1* is indeed the target of such deletions. Chromosome 3 is dense in cancer genes and the *BAP1* region, for instance, encompasses the tumor-suppressor gene *PBRM1*, which was recently found mutated in rare UMs. To confirm the *BAP1* locus hypothesis and the classification, validation series are required, ideally together with further work sequencing *BAP1* to confirm the presence of a second hit.

Conclusions

These findings suggest that partial deletion of chromosome 3 encompassing the *BAP1* locus is associated with poor prognosis. Consequently, a new cytogenetic classification of UMs is proposed, based on the status of the *BAP1* locus instead of chromosome 3. The very frequent loss of the whole chromosome 3 in UMs raises the possibility of other genes associated with UM tumorigenesis on this chromosome.

Funding/Support: This research was funded by the INCa/ITMO/AVIESAN PhD fellowship program “Formation à la recherche translationnelle” (M. Rodrigues), the Institut National de la Santé et de la Recherche Médicale (INSERM), the Ligue Nationale Contre le Cancer (Labellisation) and the Institut Curie.

Role of the Funder/Sponsor: None of the funders had any role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and the decision to submit the manuscript for publication.

Acknowledgments: We thank the patients and their family members. We thank Sylvain Dureau for his help with review.

Author Contributions: M. Rodrigues had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Rodrigues, Savignoni, Stern, Pierron

Acquisition, analysis, or interpretation of data: Rodrigues, Ait-Rais, Salviat, Algret, Simaga, Barnhill, Gardrat, Servois, Mariani, Piperno-Neumann, Roman-Roman, Delattre, Cassoux, Savignoni, Stern, Pierron.

Drafting of the manuscript: Rodrigues, Savignoni, Stern, Pierron.

Critical revision of the manuscript for important intellectual content: Rodrigues, Ait-Rais, Salviat, Algret, Simaga, Barnhill, Gardrat, Servois, Mariani, Piperno-Neumann, Roman-Roman, Delattre, Cassoux, Savignoni, Stern, Pierron

Supervision: Rodrigues, Savignoni, Stern, Pierron.

348 **Conflicts of Interest:** The authors declare no conflict of interest.

References

1. Mahendraraj K, Lau CS, Lee I, Chamberlain RS. Trends in incidence, survival, and management of uveal melanoma: a population-based study of 7,516 patients from the Surveillance, Epidemiology, and End Results database (1973-2012). *Clin Ophthalmol*. 2016;10:2113-2119.
2. Desjardins L, Levy-Gabriel C, Lumbroso-Lerouic L, et al. [Prognostic factors for malignant uveal melanoma. Retrospective study on 2,241 patients and recent contribution of monosomy-3 research]. *Journal francais d'ophtalmologie*. Sep 2006;29(7):741-749.
3. Singh AD, Turell ME, Topham AK. Uveal melanoma: trends in incidence, treatment, and survival. *Ophthalmology*. Sep 2011;118(9):1881-1885.
4. Kujala E, Makitie T, Kivela T. Very long-term prognosis of patients with malignant uveal melanoma. *Investigative ophthalmology & visual science*. Nov 2003;44(11):4651-4659.
5. Brierley JD, Gospodarowicz MK, Wittekind C. *The TNM Classification of Malignant Tumours. 8th edition*. 2016.
6. Field MG, Decatur CL, Kurtenbach S, et al. PRAME as an Independent Biomarker for Metastasis in Uveal Melanoma. *Clinical cancer research : an official journal of the American Association for Cancer Research*. Mar 1 2016;22(5):1234-1242.
7. Onken MD, Worley LA, Ehlers JP, Harbour JW. Gene expression profiling in uveal melanoma reveals two molecular classes and predicts metastatic death. *Cancer research*. Oct 15 2004;64(20):7205-7209.

- 372 8. Prescher G, Bornfeld N, Becher R. Nonrandom chromosomal abnormalities in
373 primary uveal melanoma. *Journal of the National Cancer Institute*. Nov 21
374 1990;82(22):1765-1769.
- 375 9. Prescher G, Bornfeld N, Hirche H, Horsthemke B, Jockel KH, Becher R.
376 Prognostic implications of monosomy 3 in uveal melanoma. *Lancet*. May 4
377 1996;347(9010):1222-1225.
- 378 10. Cassoux N, Rodrigues MJ, Plancher C, et al. Genome-wide profiling is a
379 clinically relevant and affordable prognostic test in posterior uveal melanoma. *The*
380 *British journal of ophthalmology*. Jun 2014;98(6):769-774.
- 381 11. Abdel-Rahman MH, Pilarski R, Cebulla CM, et al. Germline BAP1 mutation
382 predisposes to uveal melanoma, lung adenocarcinoma, meningioma, and other
383 cancers. *Journal of medical genetics*. Dec 2011;48(12):856-859.
- 384 12. Robertson AG, Shih J, Yau C, et al. Integrative Analysis Identifies Four
385 Molecular and Clinical Subsets in Uveal Melanoma. *Cancer Cell*. Jan 8
386 2018;33(1):151.
- 387 13. Field MG, Durante MA, Anbunathan H, et al. Punctuated evolution of
388 canonical genomic aberrations in uveal melanoma. *Nature communications*. Jan 9
389 2018;9(1):116.
- 390 14. Rai K, Pilarski R, Boru G, et al. Germline BAP1 alterations in familial uveal
391 melanoma. *Genes, Chromosomes & Cancer*. Feb 2017;56(2):168-174.
- 392 15. Harbour JW, Onken MD, Roberson ED, et al. Frequent mutation of BAP1 in
393 metastasizing uveal melanomas. *Science*. Dec 3 2010;330(6009):1410-1413.
- 394 16. Wiesner T, Obenaus AC, Murali R, et al. Germline mutations in BAP1
395 predispose to melanocytic tumors. *Nature genetics*. Aug 28 2011;43(10):1018-1021.

- 396 17. Abdel-Rahman MH, Christopher BN, Faramawi MF, et al. Frequency,
397 molecular pathology and potential clinical significance of partial chromosome 3
398 aberrations in uveal melanoma. *Modern pathology : an official journal of the United*
399 *States and Canadian Academy of Pathology, Inc.* Jul 2011;24(7):954-962.
- 400 18. Trolet J, Hupe P, Huon I, et al. Genomic profiling and identification of high-risk
401 uveal melanoma by array CGH analysis of primary tumors and liver metastases.
402 *Investigative ophthalmology & visual science.* Jun 2009;50(6):2572-2580.
- 403 19. Furney SJ, Pedersen M, Gentien D, et al. SF3B1 mutations are associated
404 with alternative splicing in uveal melanoma. *Cancer discovery.* Oct 2013;3(10):1122-
405 1129.
- 406 20. Johansson P, Aoude LG, Wadt K, et al. Deep sequencing of uveal melanoma
407 identifies a recurrent mutation in PLCB4. *Oncotarget.* Jan 26 2016;7(4):4624-4631.
- 408 21. Mensink HW, Vaarwater J, de Keizer RJ, et al. Chromosomal aberrations in
409 iris melanomas. *The British journal of ophthalmology.* Mar 2011;95(3):424-428.
- 410 22. Worley LA, Onken MD, Person E, et al. Transcriptomic versus chromosomal
411 prognostic markers and clinical outcome in uveal melanoma. *Clinical cancer research*
412 *: an official journal of the American Association for Cancer Research.* Mar 1
413 2007;13(5):1466-1471.
- 414 23. van Gils W, Lodder EM, Mensink HW, et al. Gene expression profiling in uveal
415 melanoma: two regions on 3p related to prognosis. *Investigative ophthalmology &*
416 *visual science.* Oct 2008;49(10):4254-4262.
- 417 24. Singh AD, Aronow ME, Sun Y, et al. Chromosome 3 status in uveal
418 melanoma: a comparison of fluorescence in situ hybridization and single-nucleotide
419 polymorphism array. *Investigative ophthalmology & visual science.* Jun 5
420 2012;53(7):3331-3339.

- 421 25. Damato B, Dopierala J, Klaasen A, van Dijk M, Sibbring J, Coupland SE.
422 Multiplex ligation-dependent probe amplification of uveal melanoma: correlation with
423 metastatic death. *Investigative ophthalmology & visual science*. Jul 2009;50(7):3048-
424 3055.
- 425 26. Larsen AC, Holst L, Kaczkowski B, et al. MicroRNA expression analysis and
426 Multiplex ligation-dependent probe amplification in metastatic and non-metastatic
427 uveal melanoma. *Acta ophthalmologica*. Sep 2014;92(6):541-549.
- 428 27. Kalirai H, Dodson A, Faqir S, Damato BE, Coupland SE. Lack of BAP1 protein
429 expression in uveal melanoma is associated with increased metastatic risk and has
430 utility in routine prognostic testing. *British journal of cancer*. Sep 23
431 2014;111(7):1373-1380.
- 432 28. van Essen TH, van Pelt SI, Versluis M, et al. Prognostic parameters in uveal
433 melanoma and their association with BAP1 expression. *Br J Ophthalmol*. Aug 21
434 2014.
- 435 29. Thomas S, Putter C, Weber S, Bornfeld N, Lohmann DR, Zeschnigk M.
436 Prognostic significance of chromosome 3 alterations determined by microsatellite
437 analysis in uveal melanoma: a long-term follow-up study. *British Journal of Cancer*.
438 Mar 13 2012;106(6):1171-1176.
- 439 30. Tschentscher F, Prescher G, Horsman DE, et al. Partial deletions of the long
440 and short arm of chromosome 3 point to two tumor suppressor genes in uveal
441 melanoma. *Cancer research*. Apr 15 2001;61(8):3439-3442.
- 442 31. Cross NA, Rennie IG, Murray AK, Sisley K. The identification of chromosome
443 abnormalities associated with the invasive phenotype of uveal melanoma in vitro. *Clin*
444 *Exp Metastasis*. 2005;22(2):107-113.

- 445 32. Shields CL, Ganguly A, Bianciotto CG, Turaka K, Tavallali A, Shields JA.
446 Prognosis of uveal melanoma in 500 cases using genetic testing of fine-needle
447 aspiration biopsy specimens. *Ophthalmology*. Feb 2011;118(2):396-401.
- 448 33. Damato B, Dopierala JA, Coupland SE. Genotypic profiling of 452 choroidal
449 melanomas with multiplex ligation-dependent probe amplification. *Clinical cancer*
450 *research : an official journal of the American Association for Cancer Research*. Dec
451 15 2010;16(24):6083-6092.
- 452 34. Ewens KG, Kanetsky PA, Richards-Yutz J, et al. Genomic profile of 320 uveal
453 melanoma cases: chromosome 8p-loss and metastatic outcome. *Invest Ophthalmol*
454 *Vis Sci*. Aug 23 2013;54(8):5721-5729.
- 455 35. Knudson AG. Two genetic hits (more or less) to cancer. *Nat Rev Cancer*. Nov
456 2001;1(2):157-162.
- 457 36. Rodrigues M, Mobuchon L, Houy A, et al. Outlier response to anti-PD1 in
458 uveal melanoma reveals germline MBD4 mutations in hypermutated tumors. *Nature*
459 *communications*. May 14 2018;9(1):1866.
- 460 37. Rodrigues M, Mobuchon L, Houy A, et al. Evolutionary routes in metastatic
461 uveal melanomas depend on MBD4 alterations. *Clin Cancer Res*. Jun 21 2019.
- 462 38. Johansson PA, Stark A, Palmer JM, et al. Prolonged stable disease in a uveal
463 melanoma patient with germline MBD4 nonsense mutation treated with
464 pembrolizumab and ipilimumab. *Immunogenetics*. May 2019;71(5-6):433-436.
- 465 39. Carvajal RD, Schwartz GK, Tezel T, Marr B, Francis JH, Nathan PD.
466 Metastatic disease from uveal melanoma: treatment options and future prospects.
467 *The British journal of ophthalmology*. Jan 2017;101(1):38-44.
- 468 40. Piperno-Neumann S, Rodrigues MJ, Servois V, et al. A randomized
469 multicenter phase 3 trial of adjuvant fotemustine versus surveillance in high risk uveal

melanoma (UM) patients (FOTEADJ). *Journal of Clinical Oncology*.
2017;35(15_suppl):9502-9502.

41. Smit KN, van Poppelen NM, Vaarwater J, et al. Combined mutation and copy-
number variation detection by targeted next-generation sequencing in uveal
melanoma. *Mod Pathol*. May 2018;31(5):763-771.

42. Afshar AR, Damato BE, Stewart JM, et al. Next-Generation Sequencing of
Uveal Melanoma for Detection of Genetic Alterations Predicting Metastasis. *Transl
Vis Sci Technol*. Mar 2019;8(2):18.

43. Matet A, Ait Rais K, Malaise D, et al. Comparative Cytogenetic Abnormalities
in Paired Choroidal Melanoma Samples Obtained Before and After Proton Beam
Irradiation by Transscleral Fine-Needle Aspiration Biopsy and Endoresection.
Cancers (Basel). Aug 14 2019;11(8).

44. Shain AH, Bagger MM, Yu R, et al. The genetic evolution of metastatic uveal
melanoma. *Nature Genetics*. Jul 2019;51(7):1123-1130.

45. Angi M, Kalirai H, Taktak A, et al. Prognostic biopsy of choroidal melanoma:
an optimised surgical and laboratory approach. *The British journal of ophthalmology*.
Aug 2017;101(8):1143-1146.

Tables

Table 1. Univariate and multivariate analyses of risk factors for metastasis.

HR (95%CI): hazard ratio (95% confidence interval); mm: millimeters; n: number of cases.

Univariate analysis				
		n	HR (95%CI)	p-value
Age	< 60 years-old	23	1	.17
	≥ 60 years-old	20	0.49 (0.17-1.4)	
Gender	male	21	1	.14
	female	22	0.46 (0.16-1.33)	
Diameter	≤ 15 mm	17	1	.31
	> 15 mm	26	1.72 (0.6-4.95)	
Thickness	≤ 10 mm	22	1	.43
	> 10 mm	21	1.49 (0.55-4)	
Tumor location	on the equator	28	1	.07
	anterior to the equator	4	2.55 (0.69-9.36)	
	posterior to the equator	10	0.36 (0.08-1.62)	
Retinal detachment	No	4	1	.06
	Yes	39	0.32 (0.09-1.11)	
Histology	spindle cells	10	1	1.00
	epithelioid/mixed	13	1 (0.28-3.55)	
BAP1 locus deletion	No	24	1	.001
	Yes	19	5.91 (1.89-18.54)	
8q gain	No	16	1	.007
	Yes	27	6.02 (1.36-26.61)	

Multivariate analysis				
		n	HR (95%CI)	p-value
BAP1 locus deletion	No	24		.001
	Yes	19	6.65 (2.09 ; 21.18)	
8q gain	No	16		.01
	Yes	27	6.88 (1.53 ; 30.86)	

Figures

Figure 1. Copy number profiles in metastatic *versus* non-metastatic cases.

Frequencies of losses at a given position are shown at the bottom. Light gray: non-metastatic cases (Met-; n=30); dark gray: metastatic cases (Met+; n=13).

Figure 2. Copy number profiles in *BAP1*del cases *versus* *BAP1*nl.

Frequencies of deletion at a given position are shown at the bottom. Light gray: *BAP1*nl cases (n=19); dark gray: *BAP1*del cases (n=24).

Figure 3. Metastasis-free and overall survivals according to the loss of the

***BAP1* locus.** Metastasis-free survival (left) and overall survival (right) curves in UMs with a partial loss of chromosome 3 encompassing the *BAP1* locus or not. *BAP1*del: deletion of the *BAP1* locus; *BAP1*nl: absence of loss of the *BAP1* locus.

Figure 4. Metastasis-free and overall survivals according to the three different

prognosis groups. Metastasis-free survival (left) and overall survival (right) curves in UMs with a partial loss of chromosome 3 according to the three different prognosis groups.





Figure 1. Rodrigues et al.






Figure 3. Rodrigues et al.





Figure 4. Rodrigues et al.