



HAL
open science

Bixafen, a succinate dehydrogenase inhibitor fungicide, causes microcephaly and motor neuron axon defects during development

Alexandre Brenet, Rahma Hassan-Abdi, Nadia Soussi-Yanicostas

► To cite this version:

Alexandre Brenet, Rahma Hassan-Abdi, Nadia Soussi-Yanicostas. Bixafen, a succinate dehydrogenase inhibitor fungicide, causes microcephaly and motor neuron axon defects during development. *Chemosphere*, 2020, pp.128781. 10.1016/j.chemosphere.2020.128781 . inserm-03058531

HAL Id: inserm-03058531

<https://inserm.hal.science/inserm-03058531>

Submitted on 11 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Contents lists available at ScienceDirect

Chemosphere

journal homepage: www.elsevier.com/locate/chemosphere

Short Communication

Bixafen, a succinate dehydrogenase inhibitor fungicide, causes microcephaly and motor neuron axon defects during development


Alexandre Brenet¹, Rahma Hassan-Abdi¹, Nadia Soussi-Yanicostas^{*}

Université de Paris, NeuroDiderot, Inserm, F-75019, Paris, France

HIGHLIGHTS

- Zebrafish embryos exposed to the succinate dehydrogenase inhibitor (SDHI) bixafen show CNS development defects.
- Brain development is disturbed in embryos exposed to SDHI, bixafen, leading to microcephaly.
- Spinal cord axon outgrowth and branching are adversely affected in embryos exposed to SDHI bixafen.
- The developing CNS is vulnerable to bixafen exposure.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 1 July 2020

Received in revised form

7 October 2020

Accepted 25 October 2020

Available online xxx

Handling Editor: A. Gies

Keywords:

Succinate dehydrogenase inhibitors (SDHIs)

Fungicide

Zebrafish

Neurodevelopment

In vivo imaging

ABSTRACT

Succinate dehydrogenase inhibitors (SDHIs), the most widely used fungicides in agriculture today, act by blocking succinate dehydrogenase (SDH), an essential and evolutionarily conserved component of mitochondrial respiratory chain. Recent results showed that several SDHIs used as fungicides not only inhibit the SDH activity of target fungi but also block this activity in human cells in *in vitro* models, revealing a lack of specificity and thus a possible health risk for exposed organisms, including humans. Despite the frequent detection of SDHIs in the environment and on harvested products and their increasing use in modern agriculture, their potential toxic effects *in vivo*, especially on neurodevelopment, are still under-evaluated. Here we assessed the neurotoxicity of bixafen, one of the latest-generation SDHIs, which had never been tested during neurodevelopment. For this purpose, we used a well-known vertebrate model for toxicity testing, namely zebrafish transparent embryos, and live imaging using transgenic lines labelling the brain and spinal cord. Here we show that bixafen causes microcephaly and defects on motor neuron axon outgrowth and their branching during development. Our findings show that the central nervous system is highly sensitive to bixafen, thus demonstrating *in vivo* that bixafen is neurotoxic in vertebrates and causes neurodevelopmental defects. This work adds to our knowledge of the toxic effect of SDHIs on neurodevelopment and may help us take appropriate precautions to ensure protection against the neurotoxicity of these substances.

© 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Bixafen, a methyl-pyrazole carboxamide, is a fungicide widely used on cereal and rapeseed crops. It was initially approved and marketed in 2011 by Bayer (Ravichandra 2018). Its high efficiency and rapid penetration resulted in a significant rise in usage in

* Corresponding author.

E-mail address: nadia.soussi@inserm.fr (N. Soussi-Yanicostas).

¹ Co-first authors.




Fig. 1. Bixafen exposure causes microcephaly. (A) Experimental setup used to analyze the body morphological parameters and brain size of zebrafish embryos exposed to 0.2 or 0.5 μM bixafen, from 6 hpf onward up to 36 or 66 hpf (brain analysis), or 96 hpf (morphological analysis). (B–D) Lateral and dorsal views of 96 hpf larvae treated with 0.1% DMSO (B) or 0.2 μM (C) or 0.5 μM bixafen (D). (E) Scheme depicting the different morphological parameters measured: body length (black double-arrow), head size (purple area), eye diameter (red double-arrow) and eye-otolith distance (green double-arrow). (F–I) Body length (F), head size (G), eye diameter (H) and eye-otolith distance (I), of 96 hpf larvae exposed to 0.1% DMSO (n = 14), or 0.2 μM (n = 15) or 0.5 μM (n = 12) bixafen. (J–O) Dorsal views of the brain of 36 (J–L) or 66 hpf transgenic Tg[huc:G/U:RFP] embryos (M–O),

Europe and the US. Currently, nearly 30 different products are sold on the French market alone that contain bixafen as sole active substance or mixed with other fungicides. Bixafen belongs to the succinate dehydrogenase inhibitor (SDHI) family, the fungicides most widely used in agriculture to fight a broad range of fungal diseases (Lalève et al., 2014; Zhang et al., 2019). Bixafen, which is derived from carboxin and categorized as a latest-generation SDHIs, acts through inhibition of mitochondrial respiration chain complex II, also known as succinate dehydrogenase (SDH). However, the inhibitory effect of SDHIs is not restricted to fungi: they also block the mitochondrial respiration chain in many other species, including human cells (Bénit et al., 2019). Additionally, the latest-generation SDHIs have been shown to also inhibit mitochondrial respiration complex III (Bénit et al., 2019), suggesting more critical effects of these new fungicides. SDH activity is essential for proper mitochondrial respiration and metabolism, and defects in its activity impair the cellular metabolome and functions (Bénit et al., 2014). Many human diseases are related to SDH deficiency, including neurodegenerative disorders, such as Leigh syndrome, an early-onset neurodegenerative disease characterized by developmental delay, ataxia and seizures (Birch-Machin et al., 2000; Bourgeron 2015; Van Coster et al., 2003; Finster 2008; Horváth et al., 2006; Parfait et al., 2000), and cancers, such as familial paraganglioma syndrome, infantile leukoencephalopathy, and neuroblastoma (Ghezzi et al., 2009; Martin et al., 2007; Perry et al., 2006; Ricketts et al., 2009; Stratakis and Carney 2009; Timmers et al., 2009). The Norwegian scientific committee for food safety (VKM) has assessed the health and environmental risk of Aviator Xpro EC225, a fungicide containing bixafen, and concluded that the effects of bixafen in animals should be considered relevant for humans (VKM Report 2014). Also, M44, a bixafen metabolite, which has the potential to contaminate groundwater, causes abnormalities in the rabbit foetus, suggesting that similar effects in humans cannot be excluded (VKM Report 2015). However, despite the increasing use of SDHIs in modern agriculture and their frequent detection in the environment and on harvested products (Abad-Fuentes et al., 2015; Añasco et al. 2010; Tanabe and Kawata 2009; Tsuda et al., 2009; Vu et al., 2016), the potential toxic effects of these substances, especially on neurodevelopment, have so far been poorly evaluated.

To fill this gap and assess the neurotoxicity of these latest-generation SDHI fungicides *in vivo*, we tested the potential toxic effect of bixafen on zebrafish embryos. We observed that embryos exposed to low or medium bixafen concentrations, LOAEL and LC30, developed a series of defects, including microcephaly and disorganized motor neuron axons and branches. This study demonstrated the toxicity of bixafen on CNS development and gives insight into the environmental risk of this SDHI on the nervous system in vertebrates.

2. Results

2.1. Toxicity of bixafen *in vivo* in zebrafish embryos

To assess the potential toxicity of bixafen, a SDHI belonging to the pyrazole group of fungicides, zebrafish embryos were exposed to increasing concentrations of bixafen ranging from 0.1 μM to 4 μM , from 6 h post-fertilization (hpf) onward up to 96 hpf (Supplementary Fig. 1A). As no differences in survival rates or

embryo morphology could be detected between the 0.1% DMSO and fish water control groups (data not shown), embryos incubated in 0.1% DMSO were used as controls. Exposure of zebrafish embryos to 0.5 μM bixafen and above induced gross morphological defects and significantly increased mortality, compared with that seen in controls (Supplementary Fig. 1). Moreover, we observed a dose-dependent effect, with increasing concentrations of bixafen inducing increased mortality rates and increased phenotypic defects in surviving embryos (Supplementary Fig. 1B₁-K).

Results indicated that the median lethal concentration of bixafen was 2.7 μM for wild-type zebrafish embryos (AB line) after 96 h of incubation, i.e. 96 hpf LC50, and closely similar values were observed with the two other zebrafish lines used in this study, the transgenic Tg[Olig2:eGFP] and Tg[elav13:Gal4/5xUAS:RFP] double transgenic line (Akerboom et al., 2012; Asakawa et al., 2008), hereafter referred to as Tg[HuC:G/U:RFP] (Supplementary Fig. 1K).

To investigate the potential toxicity of bixafen, and particularly the effect of this SDHI on the morphology of the brain and spinal cord, we chose 0.2 μM , the lowest-observed-adverse-effect-level (LOAEL), and 0.5 μM , the 96 hpf LC30, two concentrations that induce mild and moderate phenotypic defects in 96 hpf embryos, respectively, except for small heart oedema in embryos exposed to 0.2 μM bixafen (Fig. 1C), and large heart oedema and curved tail in embryos exposed to 0.5 μM bixafen (Fig. 1D). We next carefully examined the morphological parameters of embryos exposed to 0.2 and 0.5 μM bixafen, namely body length, head size, eye diameter and eye-otolith distance. Results indicated that following exposure to either 0.2 or 0.5 μM bixafen, surviving embryos show reduced body length and eye diameter, and decreased head size and eye-otolith distance, compared with that observed in age-matched controls (Fig. 1E-I).

2.2. Brain defects in embryos exposed to bixafen

Because several investigations have reported that exposure to SDHIs may induce neurotoxicity (Wang et al., 2020; Yao et al., 2018), we took advantage of the Tg[HuC:G/U:RFP] transgenic line, in which all post-mitotic brain neurons are labelled by the fluorescent RFP protein, to precisely examine the morphology of the brain of 36 and 66 hpf embryos, following exposure to 0.2 and 0.5 μM bixafen. Live imaging of the brain of 36 and 66 hpf embryos, clearly indicated that exposure to both 0.2 and 0.5 μM bixafen, induced smaller brain size (Fig. 1J-O). Precise quantification of brain volumes (Imaris software, Bitplane), indicated that 36 hpf embryos exposed to 0.2 ($3.07 \pm 0.13 \times 10^6 \mu\text{m}^3$) or 0.5 μM bixafen ($1.60 \pm 0.13 \times 10^6 \mu\text{m}^3$), displayed a significantly smaller brain compared with that seen in age-matched controls ($3.57 \pm 0.07 \times 10^6 \mu\text{m}^3$) (Fig. 1P). Similarly, 66 hpf embryos exposed to 0.2 ($7.45 \pm 0.24 \times 10^6 \mu\text{m}^3$) or 0.5 μM bixafen ($3.64 \pm 0.15 \times 10^6 \mu\text{m}^3$), also displayed small brains compared to age-matched controls ($8.51 \pm 0.25 \times 10^6 \mu\text{m}^3$) (Fig. 1Q). Our data thus showed that exposure of zebrafish embryos to low-dose bixafen corresponding to LOAEL induced microcephaly, a phenotype exacerbated in individuals incubated in 0.5 μM bixafen.

2.3. Impaired mobility and spinal motor neuron axon outgrowth defects in embryos exposed to bixafen

The small size of the brain observed in embryos exposed to

treated with 0.1% DMSO (J, M), or 0.2 (K, N) or 0.5 μM bixafen (L, O), and showing the microcephaly of embryos exposed to bixafen low/medium doses. (P-Q) Measurements of brain volume of 36 (P) or 66 hpf (Q) embryos exposed to 0.1% DMSO ($n_{36\text{hpf}} = 8$; $n_{66\text{hpf}} = 5$), or 0.2 μM ($n_{36\text{hpf}} = 6$; $n_{66\text{hpf}} = 5$) or 0.5 μM bixafen ($n_{36\text{hpf}} = 8$; $n_{66\text{hpf}} = 4$), further illustrate the microcephaly of embryos exposed to bixafen at 96 hpf LC30 and LOAEL concentrations. ****, $p < 0.0001$; ***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$; n.s., non-significant. Scale bar: (B-D) = 0.5 mm; (J-O) = 100 μm . Abbreviations: Tel, telencephalon; OT, optic tectum; Cer, cerebellum; SC, spinal cord. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)




Fig. 2. Bixafen exposure induces motor deficit and motor neuron axon defects. (A) Experimental setup used to analyze the motor activity and patterning of caudal primary motor neuron (CaPMN) axons and branches, in zebrafish embryos exposed to 0.2 or 0.5 μM bixafen, from 6 hpf onward up to 36 or 66 hpf (CaPMN axon morphology analysis), or 96 hpf (motor activity analysis). (B) Tracking plots illustrating the distance traveled by 96 hpf larvae treated with 0.1% DMSO ($n = 85$), or 0.2 ($n = 72$) or 0.5 μM bixafen ($n = 77$), during a 25 min time period. (C) Quantification of the distance traveled by 96 hpf larvae exposed to 0.1% DMSO, or 0.2 or 0.5 μM bixafen, during a 25 min time period. (D–I) Three-dimensional reconstruction of the organization of CaPMN axons and branches in 36 (D–F) or 66 hpf transgenic Tg[Olig2:eGFP] embryos (G–I), exposed to 0.1% DMSO (D, G),

bixafen suggests a neurotoxic effect of this SDHI. To further investigate this issue and precisely assess the consequences of bixafen exposure on brain functioning, we studied and compared the locomotor activity of 0.2 and 0.5 μM bixafen-exposed and control embryos by measuring the distance traveled by these individuals over a 25 min period, using a ZebraBox tracking automate (View-Point®). Results indicated that embryos treated with 0.2 or 0.5 μM bixafen, displayed reduced mobility, as shown by the distance traveled over a 25 min period 77.3 ± 5.7 mm ($N = 4$, $n = 72$) and 52.7 ± 4.4 mm ($N = 4$, $n = 77$), respectively, compared to the distance traveled by controls (97.5 ± 4.9 mm, $N = 4$, $n = 85$) (Fig. 2B–C). Interestingly, the hypoactive behavior of embryos exposed to 0.2 μM bixafen was markedly worsened in those exposed to 0.5 μM , which were almost motionless.

Because bixafen exposure induces brain defects, we next determined whether the locomotor deficits observed in bixafen-exposed embryos was a consequence of defects in spinal cord motor neurons or their axons. To address this issue, we next analyzed the morphology of the spinal cord in 36 and 66 hpf embryos exposed to 0.2 and 0.5 μM bixafen, using the Tg[Olig2:eGFP] transgenic line (Shin et al., 2003), in which all motor neuron progenitors and their axons are labelled by eGFP. In particular, we focused on caudal primary motor neurons (CaPMN) and their axons, which are the most accessible.

Live imaging indicated that in 36 hpf controls, primary motor neuron axons extended ventrally to contact target muscles, forming a highly regular pattern (Fig. 2D). By contrast, motor neuron axons were markedly shortened in 36 hpf embryos treated with 0.2 bixafen (Fig. 2E), a phenotype markedly exacerbated in age-matched embryos exposed to 0.5 μM , bixafen (Fig. 2F). Three-dimensional reconstruction of CaPMN and their axons in 66 hpf control embryos revealed a highly stereotyped pattern with axons extending ventrally and branches projecting dorsally and laterally along the dorsal myoseptum to form a regular pattern in the space between the notochord and the myotome (Fig. 2G, G'). By contrast, following exposure to 0.2 μM bixafen, 66 hpf embryos displayed significantly shorter axons (Fig. 2H, H'), a phenotype significantly worsened in embryos exposed to 0.5 μM bixafen (Fig. 2I, I'), which also showed a marked disorganization of motor neuron axons and their branches, which extended in all directions.

Measurements of the length of CaPMN axons confirmed that bixafen exposure induced a marked shortening of these axons. Whereas in control embryos, the length of CaPMN axons was 167.9 ± 2.4 μm and 368.3 ± 9.3 μm , at 36 and 66 hpf, respectively, in embryos exposed to 0.2 μM , these values were reduced to 148.6 ± 3.1 μm at 36 hpf and 296.4 ± 9.0 μm at 66 hpf. Moreover, these defects were exacerbated in embryos exposed to 0.5 μM bixafen, with a mean CaPMN axon length of 61.8 ± 3.7 μm at 36 hpf and 223.7 ± 9.6 μm at 66 hpf (Fig. 2J and K). Thus, bixafen is neurotoxic and impairs the outgrowth and patterning of spinal motor axons and their branches.

3. Discussion

It is well-known that the developing CNS in vertebrates is especially sensitive and vulnerable to toxic chemicals, including phytosanitary agents, which are increasingly present in the environment. More specifically, in recent years, bixafen, a latest-generation succinate dehydrogenase inhibitor (SDHI), has become

one of the most widely-used fungicides worldwide. However, the toxicity of bixafen has been under-appraised *in vivo* in vertebrate models so far, and especially its neurotoxicity for the developing CNS, an especially serious issue because of the complete lack of species-specificity of this latest-generation SDHI (Bénit et al., 2019). In the present work, we used zebrafish embryos as an *in vivo* model and demonstrate that bixafen exposure during embryogenesis induces a series of CNS defects.

In close agreement with previous investigations (Li et al., 2020; Wu et al., 2018), we first observed that zebrafish embryos exposed to bixafen showed clear developmental delays, as shown by morphological examination of embryos exposed to 0.5 μM bixafen and above, but also by a decrease in the eye-otolith distance and hypopigmentation, both observed in embryos incubated in 0.2 and 0.5 μM bixafen, corresponding to LOEL and 96 hpf LC30, respectively. Moreover, at the brain level, we observed that these two same concentrations induced brain defects and microcephaly, an observation in close agreement with the markedly reduced expression of the neuronal *neuroD* gene observed in zebrafish embryos treated with bixafen (Li et al., 2020). Microcephaly has been repeatedly observed in embryos exposed to many pesticides, including Maxim® XL (Svartz et al., 2016) and glyphosate (Paganelli et al., 2010), emphasizing the high vulnerability of the brain to these chemicals, and our study shows that SDHI bixafen also induces microcephaly, even at low doses.

In addition, using an automatic device that quantifies the speed and distance traveled by embryos, we also observed that 96 hpf embryos exposed to 0.2 or 0.5 M bixafen displayed reduced motility or almost complete paralysis, respectively. Quantification of the swimming behavior clearly showed motor deficits in embryos exposed to 0.2 μM bixafen which worsens in embryos exposed to 0.5 μM (Fig. 2C), that likely results from neuron abnormalities, not developmental delay. As it is known that the motor neuron is a major cell type regulating swimming behavior in zebrafish (Brustein et al., 2003), the motor defects seen in bixafen-exposed embryos prompted us to investigate the differentiation of motor neuron axons and branches in embryos exposed to 0.2 and 0.5 μM bixafen. In control embryos, motor neurons extend their axons in a highly stereotyped manner during development. In particular, caudal primary motor neuron (CaPMN) axons exit the spinal cord and extend toward the ventral muscle. Interestingly, in embryos exposed to bixafen, we observed a reduction in axon outgrowth and defects in their branches, even in those exposed to 0.2 μM bixafen corresponding to LOEL, indicating that the development of motor neuron axons is highly sensitive to bixafen. Although the toxic effects of bixafen on the CNS are probably due at least in part to developmental delay, the disorganization of motor neuron axons and branches clearly reveals a neurotoxicity of bixafen in zebrafish embryos.

Bixafen is a methyl-pyrazole, one of the SDHI fungicides used to manage plant diseases and inhibit the respiration of pathogenic fungi by blocking succinate complex II in the mitochondrial respiratory chain. SDH is an enzyme involved in both oxidative phosphorylation and the tricarboxylic acid cycle, two processes that generate energy. This enzyme has been shown to be irreplaceable in mitochondrial and cell metabolism and also to be highly conserved among all fungi, plants and animal species (Rehfus et al., 2016; Veloukas et al. 2013; Yamashita and Fraaije 2018). Any adverse change or inhibition of SDH activity can therefore lead to

0.2 (E, H) or 0.5 μM bixafen (F, I), showing the disorganization of CaPMN axons and branches, following exposure to 0.2 and, even more, 0.5 μM bixafen (white arrow). (G–I') Higher magnifications of the morphology of CaPMN axons and branches, in embryos exposed to 0.1% DMSO (G'), or 0.2 (H') or 0.5 μM bixafen (I'). (J, K) Quantification of the length of the primary CaPMN axons in 36 (J) and 66 hpf embryos (K), treated with DMSO ($n_{36\text{hpf}} = 13$; $n_{66\text{hpf}} = 7$), or 0.2 ($n_{36\text{hpf}} = 8$; $n_{66\text{hpf}} = 6$) or 0.5 μM bixafen ($n_{36\text{hpf}} = 22$; $n_{66\text{hpf}} = 9$), showing significantly shorter axons following exposure to bixafen at LOEL (0.2 μM) and, even more, 96 hpf LC30 (0.5 μM). ****, $p < 0.0001$; ***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$. Scale bar: (D–I) = 100 μm ; (G'–I') = 50 μm .

many diseases, including those that have been linked to mitochondrial dysfunction (Birch-Machin et al., 2000; Bourgeron 2015; Van Coster et al., 2003; Finsterer 2008; Horváth et al., 2006; Parfait et al., 2000).

Several epidemiological studies published in the last two decades suggest harmful effects of pesticides on human health (Merhi et al., 2007; Weichenthal et al. 2010). Pesticide poisoning is a serious health problem that disproportionately affects infants and children (Rauh et al., 2012). Pesticides are known to cause millions of acute poisoning cases per year (Bertolote et al., 2006). Human exposure to pesticides can occur environmentally, through consumption in food and water (van den Berg et al., 2012). Bixafen is considered to be highly persistent in the environment (EFSA 2012). The half-lives of bixafen were 105 days in loam, 316 days in sandy loam, 1235 days in slit loam, and 82 days in water (EFSA 2012). A number of studies show that prenatal and early childhood exposure to some pesticides is associated with neurodevelopmental and cognitive disorders (Muñoz-Quezada et al., 2013; Ross et al., 2013). Aside from SDH inhibition, very little is known about the consequences of SDHI exposure on human health, and it is not known how bixafen affects the development of the CNS. As it is the case for other SDHIs, the toxic effect of bixafen observed in this study may be linked to the basic disruption of mitochondrial respiration chain complex II activity and subsequent accumulation of reactive oxygen species and oxidative DNA damage. Results showed that a high concentration of bixafen induces reactive oxygen species production and subsequent DNA damage in human cell lines (Graillot et al., 2012). However, a recent study also demonstrated that bixafen exposure did not cause oxidative stress in zebrafish (Li et al., 2020), suggesting that induction of oxidative stress may not be the main mechanism underlying bixafen-induced neurotoxicity in zebrafish embryos. Thus the brain and spinal cord defects observed in embryos exposed to low-medium bixafen concentrations may be associated with the metabolic abnormalities reported in SDHI-exposed animals (Graillot et al., 2012; Qian et al., 2018; Wu et al., 2018; Yang et al., 2018).

In summary, our study provides new evidence of bixafen toxicity on neurodevelopment in a vertebrate model, and may help us take appropriate precautions to ensure protection against the neurotoxic effects of these substances.

4. Materials and methods

4.1. Ethics statement

All the animal experiments described in the present study were conducted at the French National Institute of Health and Medical Research (INSERM) UMR 1141 in Paris in accordance with European Union guidelines for the handling of laboratory animals (http://ec.europa.eu/environment/chemicals/lab_animals/home_en.htm). They were approved by the Direction Départementale de la Protection des Populations de Paris and the French Animal Ethics Committee under reference No. 2012-15/676-0069.

4.2. Zebrafish lines and maintenance

Zebrafish were maintained at 26.5 °C in 14 h light and 10 h dark cycles. Embryos were collected by natural spawning, and 0.003% 1-phenyl-2-thiourea (PTU) was added at 1 dpf (day post-fertilization) for embryos which were imaged *in vivo* to avoid pigmentation. The following transgenic lines were used: Tg[elav13:Gal4]^{zf349} (Akerboom et al., 2012), Tg[5xUAS:RFP]^{nkuasrfp1a} (Asakawa et al., 2008) and Tg[Olig2:eGFP] (Shin et al., 2003).

4.3. Chemicals

Dimethyl sulfoxide (DMSO) hybrid-max sterile (Sigma, D2650) was diluted in fish water to a final concentration of 0.1% (v/v). Bixafen, N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1-methylpyrazole-4-carboxamide (Sigma-Aldrich, St. Louis, MO, USA, code: 32581, CAS-No: 581809-46-3, Lot# BCBS9732V, purity >98%) was dissolved at a stock concentration of 400 mM in pure DMSO (100%) and conserved at -20 °C in aliquots until use.

4.4. Toxicity assay

Wild-type (AB line) and transgenic lines, Tg[HuC:G/U:RFP] or Tg [Olig2:eGFP] embryos were collected after spawning and washed twice with clean fish water. Six hpf embryos were selected, distributed randomly in 6-well microplates, with 30 embryos per well, following OECD TG 236 fish acute toxicity test (OECD 2013). Each well was filled with 6 mL of solution for each condition, with 12 conditions in all: fish water, 0.1% DMSO, 0.1 μM, 0.2 μM, 0.3 μM, 0.5 μM, 1 μM, 1.5 μM, 2 μM, 2.5 μM, 3 μM and 4 μM bixafen in 0.1% DMSO. Each day, dead embryos were selected under OECD guidelines (coagulation, no heartbeat or no somite) and discarded.

4.5. Locomotor activity

96 hpf larvae were distributed individually in a 96 well plate in 200 μl of fish water. The plate was then placed in the recording chamber of the zebrafish system for a 30 min habituation period before 25 min recording of larvae locomotion (Brenet et al., 2019). The results indicate the distance traveled by the larvae during the 25 min recording period.

4.6. Embryo imaging

36 hpf or 66 hpf embryos were anesthetized with 112 μg/mL 3-aminobenzoic acid ethyl ester (tricaine, Sigma), immobilized in 1% low melting-point agarose at the center of a 35 mm glass-bottomed dish (Corning), and covered with fish water containing 112 μg/mL tricaine. For embryo morphology analysis, bright field images were captured using a stereomicroscope (Zeiss). Live imaging of transgenic lines Tg[HuC:G/U:RFP] and Tg[Olig2:eGFP] was carried out using a Leica SP8 laser scanning confocal microscope equipped with a Leica 20×/0.75 multi-immersion objective.

4.7. Image analysis

Body length, head size, eye diameter and eye-otolith distance measurements were determined from bright field embryo images using the ruler tool in ImageJ. Brain volume and spinal neuron CaPMN axon length three-dimensional reconstructions and quantifications were performed using Imaris Measurement Pro (Bitplane Inc.).

4.8. Statistics

All statistics were obtained using Prism5 (GraphPad) and assessed by a one-way ANOVA tests followed by a Tukey post-tests. All data are represented as means ± SEM.

Author contributions

A.B. and R.H.A. performed the experiments and the analysis, designed the figures, and wrote the original draft. N.S.Y. supervised the project and wrote the manuscript.

Funding

This work was supported by Institut National de la Santé et la Recherche Médicale (INSERM), the National Center for Scientific Research (CNRS), and the French National Research Agency (ANR-16-CE18-0010). Funding sources had no involvement in study design, collection, analysis or interpretation of data, or decision to publish.

Declaration of competing interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgments

We thank Christiane Romain (Inserm UMR 1141), Olivier Bar (Inserm UMR 1141) for their technical assistance.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.chemosphere.2020.128781>.

References

- Abad-Fuentes, Antonio, Ceballos-Alcantarilla, Eric, Mercader, Josep V., Agulló, Consuelo, Abad-Somovilla, Antonio, Francesc, A., Esteve-Turrillas, 2015. "Determination of succinate-dehydrogenase-inhibitor fungicide residues in fruits and vegetables by liquid chromatography–tandem mass spectrometry. *Anal. Bioanal. Chem.* 407 (14).
- Akerboom, J., Chen, T.W., Wardill, T.J., Tian, L., Marvin, J.S., Mutlu, S., Calderon, N.C., Esposti, F., Borghuis, B.G., Sun, X.R., Gordus, A., Orger, M.B., Portugues, R., Engert, F., Macklin, J.J., Filosa, A., Aggarwal, A., Kerr, R.A., Takagi, R., Kracun, S., Shigetomi, E., Khakh, B.S., Baier, H., Lagnado, L., Wang, S.S.H., Bargmann, C.I., Kimmel, B.E., Jayaraman, V., Svoboda, K., Kim, D.S., Schreiner, E.R., Looger, L.L., 2012. Optimization of a GCaMP calcium indicator for neural activity imaging. *J. Neurosci.* 32 (40), 13819–13840.
- Añaso, Nathaniel C., Koyama, Jiro, Uno, Seiichi, 2010. Pesticide residues in coastal waters affected by rice paddy effluents temporarily stored in a wastewater reservoir in southern Japan. *Arch. Environ. Contam. Toxicol.* 58 (2), 352–360.
- Asakawa, Kazuhide, Suster, Maximiliano L., Mizusawa, Kanta, Nagayoshi, Saori, Kotani, Tomoya, Urasaki, Akihiro, Kishimoto, Yasuyuki, Hibi, Masahiko, Kawakami, Koichi, 2008. Genetic dissection of neural circuits by Tol2 transposon-mediated Gal4 gene and enhancer trapping in zebrafish. *Proc. Natl. Acad. Sci. U.S.A.* 105 (4), 1255–1260.
- Bénit, Paule, Kahn, Agathe, Chretien, Dominique, Bortoli, Sylvie, Huc, Laurence, Schiff, Manuel, Gimenez-Roqueplo, Anne Paule, Favier, Judith, Gressens, Pierre, Rak, Malgorzata, Rustin, Pierre, 2019. Evolutionarily conserved susceptibility of the mitochondrial respiratory chain to SDHI pesticides and its consequence on the impact of SDHIs on human cultured cells. *PLoS One* 14 (11).
- Bénit, Paule, Letouzé, Eric, Rak, Malgorzata, Aubry, Laetitia, Burnichon, Nelly, Favier, Judith, Gimenez-Roqueplo, Anne Paule, Rustin, Pierre, 2014. Undiscovered task for an old team: succinate, fumarate and other Krebs cycle acids in metabolic remodeling. *Biochim. Biophys. Acta Bioenerg.* 1837 (8), 1330–1337.
- van den Berg, F., Tiktak, A., Heuvelink, G.B.M., Burgers, S.L.G.E., Brus, D.J., de Vries, F., Stolte, J., Kroes, J.G., 2012. Propagation of uncertainties in soil and pesticide properties to pesticide leaching. *J. Environ. Qual.* 41 (1), 253–261.
- Bertolote, José M., Fleischmann, A., Eddleston, M., Gunnell, D., 2006. Deaths from pesticide poisoning: a global response. *Br. J. Psychiatry* 189 (SEP), 201–203.
- Birch-Machin, Mark A., Taylor, Robert W., Cochran, Bruce, Ackrell, Brian A.C., Turnbull, Douglass M., 2000. Late-onset optic atrophy, ataxia, and myopathy associated with a mutation of a complex II gene. *Ann. Neurol.* 48 (3), 330–335.
- Bourgeron, Thomas, 2015. From the genetic architecture to synaptic plasticity in autism spectrum disorder. *Nat. Rev. Neurosci.* 16 (9), 551–563.
- Brenet, Alexandre, Hassan-Abdi, Rahma, Somkhit, Julie, Yanicostas, Constantin, Soussi-Yanicostas, Nadia, 2019. Defective excitatory/inhibitory synaptic balance and increased neuron apoptosis in a zebrafish model of dravet syndrome. *Cells* 8 (10), 1199.
- Brustein, Edna, Louis Saint-Amant, Buss, Robert R., Chong, Mabel, McDearmid, Jonathan R., Drapeau, Pierre, 2003. "Steps during the development of the zebrafish locomotor network." In: *Journal of Physiology Paris*, vol. 97, pp. 77–86.
- Van Coster, Rudy, Seneca, S., Smet, J., Van Hecke, R., Gerlo, E., Devreese, B., Van Beeumen, J., Leroy, J.G., De Meirleir, L., Lissens, W., 2003. Homozygous Gly555Glu mutation in the nuclear-encoded 70 KDa flavoprotein gene causes instability of the respiratory chain complex II. *Am. J. Med. Genet.* 120A (1), 13–18.
- EFSA, 2012. Conclusion on the peer review of the pesticide risk assessment of the active substance bifenox. *EFSA J.* 10 (11), 2917.
- Finsterer, Josef, 2008. Cognitive decline as a manifestation of mitochondrial disorders (mitochondrial dementia). *J. Neurol. Sci.* 272 (1–2), 20–33.
- Ghezzi, Daniele, Goffrini, Paola, Uziel, Graziella, Horvath, Rita, Klopstock, Thomas, Lochmüller, Hanns, D'Adamo, Pio, Gasparini, Paolo, Strom, Tim M., Prokisch, Holger, Invernizzi, Federica, Ferrero, Ileana, Zeviani, Massimo, 2009. SDHAF1, encoding a LYR complex-II specific assembly factor, is mutated in SDH-defective infantile leukoencephalopathy. *Nat. Genet.* 41 (6), 654–656.
- Graillot, Vanessa, Tomasetig, Jean, Cravedi, Jean, Audebert, Marc, 2012. Evidence of the in vitro genotoxicity of methyl-pyrazole pesticides in human cells. *Mutat. Res. Genet. Toxicol. Environ. Mutagen* 748 (1–2), 8–16.
- Horváth, Rita, Abicht, A., Holinski-Feder, E., Laner, A., Gempel, K., Prokisch, H., Lochmüller, H., Klopstock, T., Jaksch, M., 2006. Leigh syndrome caused by mutations in the flavoprotein (fp) subunit of succinate dehydrogenase (SDHA). *J. Neurol. Neurosurg. Psychiatr.* 77 (1), 74–76.
- Lalève, Anaïs, Gamet, Stéphanie, Anne, Sophie Walker, Debieu, Danièle, Toquin, Valérie, Fillinger, Sabine, 2014. Site-directed mutagenesis of the P225, N230 and H272 residues of succinate dehydrogenase subunit B from botrytis cinerea highlights different roles in enzyme activity and inhibitor binding. *Environ. Microbiol.* 16 (7), 2253–2266.
- Li, Wenhua, Yuan, Mingrui, Wu, Yaqing, Liu, Xuan, 2020. Bifenox exposure induces developmental toxicity in zebrafish (*Danio rerio*) embryos. *Environ. Res.* 189 (June), 109923.
- Martin, Caroline, Dubouchaud, Hervé, Mosoni, Laurent, Michel Chardigny, Jean, Oudot, Alexandra, Fontaine, Eric, Vergely, Catherine, Keriell, Christiane, Rochette, Luc, Lervere, Xavier, Demaison, Luc, 2007. Abnormalities of mitochondrial functioning can partly explain the metabolic disorders encountered in sarcopenic gastrocnemius. *Aging Cell* 6 (2), 165–177.
- Merhi, M., Raynal, H., Cahuzac, E., Vinson, F., Cravedi, J.P., Gamet-Payastre, L., 2007. Occupational exposure to pesticides and risk of hematopoietic cancers: meta-analysis of case-control studies. *Cancer Causes Control* 18 (10), 1209–1226.
- Muñoz-Quezada, Teresa, María, Lucero, Boris A., Barr, Dana B., Steenland, Kyle, Levy, Karen, Barry Ryan, P., Iglesias, Veronica, Alvarado, Sergio, Concha, Carlos, Rojas, Evelyn, Vega, Catalina, 2013. Neurodevelopmental effects in children associated with exposure to organophosphate pesticides: a systematic review. *Neurotoxicology* 39, 158–168.
- OECD, 2013. "Test No. 236: Fish Embryo Acute Toxicity (FET) Test." Guidelines for the Testing of Chemicals.
- Paganelli, Alejandra, Gnazzo, Victoria, Acosta, Helena, López, Silvia L., Carrasco, Andrés E., 2010. Glyphosate-based herbicides produce teratogenic effects on vertebrates by impairing retinoic acid signaling. *Chem. Res. Toxicol.* 23 (10), 1586–1595.
- Parfait, Béatrice, Chretien, Dominique, Agnès, Rötig, Marsac, Cécile, Arnold, Munnich, Rustin, Pierre, 2000. Compound heterozygous mutations in the flavoprotein gene of the respiratory chain complex II in a patient with Leigh syndrome. *Hum. Genet.* 106 (2), 236–243.
- Perry, Colin G., Young, William J., McWhinney, Sarah R., Bei, Thalia, Stergiopoulos, Sotirios, Knudson, Ryan A., Ketterling, Rhett P., Eng, Charis, Stratakis, Constantine A., Aidan Carney, J., 2006. Functioning paraganglioma and gastrointestinal stromal tumor of the jejunum in three women: syndrome or coincidence. *Am. J. Surg. Pathol.* 30 (1), 42–49.
- Qian, Le, Cui, Feng, Yang, Yang, Liu, Yuan, Qi, Suzhen, Wang, Chengju, 2018. Mechanisms of developmental toxicity in zebrafish embryos (*Danio rerio*) induced by boscalid. *Sci. Total Environ.* 634, 478–487.
- Rauh, Virginia A., Perera, Frederica P., Megan, K. Horton, Whyatt, Robin M., Bansal, Ravi, Hao, Xuejun, Liu, Jun, Barr, Dana Boyd, Slotkin, Theodore A., Peterson, Bradley S., 2012. Brain anomalies in children exposed prenatally to a common organophosphate pesticide. *Proc. Natl. Acad. Sci. U.S.A.* 109 (20), 7871–7876.
- Ravichandra, N.G., 2018. Agrochemicals in Plant Disease Management.
- Rehfs, Alexandra, Miessner, Simone, Achenbach, Janosch, Strobel, Dieter, Bryson, Rosie, Gerd Stammler, 2016. Emergence of succinate dehydrogenase inhibitor resistance of *Pyrenopeziza teres* in Europe. *Pest Manag. Sci.* 72 (10), 1977–1988.
- Ricketts, Christopher, Zeegers, Maurice P., Jan, Lubinski, Maher, Eamonn R., 2009. Analysis of germline variants in CDH1, IGFBP3, MMP1, MMP3, STK15 and VEGF in familial and sporadic renal cell carcinoma. *PLoS One* 4 (6).
- Ross, Sarah Mac Kenzie, McManus, I.C., Harrison, Virginia, Mason, Oliver, 2013. Neurobehavioral problems following low-level exposure to organophosphate pesticides: a systematic and meta-analytic review. *Crit. Rev. Toxicol.* 43 (1), 21–44.
- Shin, Jimann, Park, Hae Chul, Topczewska, Jolanta M., Madwsley, David J., Appel, Bruce, 2003. Neural cell fate analysis in zebrafish using Olig2 BAC transgenics. *Methods Cell Sci.* 25 (1–2), 7–14.
- Stratakis, C.A., Carney, J.A., 2009. "The triad of paragangliomas, gastric stromal tumours and pulmonary chondromas (Carney triad), and the dyad of paragangliomas and gastric stromal sarcomas (Carney-Stratakis syndrome): molecular genetics and clinical implications." In: *Journal of Internal Medicine*, vol. 266, pp. 43–52 (J Intern Med).
- Svartz, Gabriela, Meijide, Fernando, Coll, Cristina Pérez, 2016. Effects of a fungicide formulation on embryo-larval development, metamorphosis, and

- gonadogenesis of the south American toad *Rhinella arenarum*. *Environ. Toxicol. Pharmacol.* 45, 1–7.
- Tanabe, Akiko, Kawata, Kuniaki, 2009. Daily variation of pesticides in surface water of a small river flowing through paddy field area. *Bull. Environ. Contam. Toxicol.* 82 (6), 705–710.
- Timmers, Henri J.L. M., Gimenez-Roqueplo, Anne Paule, Mannelli, Massimo, Pacak, Karel, 2009. Clinical aspects of SDHx-related pheochromocytoma and paraganglioma. *Endocr. Relat. Canc.* 16 (2), 391–400.
- Tsuda, T., Nakamura, T., Inoue, A., Tanaka, K., 2009. Pesticides in water, fish and shellfish from littoral area of lake biwa. *Bull. Environ. Contam. Toxicol.* 82 (6), 716–721.
- Veloukas, Thomas, Markoglou, Anastasios N., Karaoglanidis, George S., 2013. Differential effect of SdhB gene mutations on the sensitivity to SDHI fungicides in *Botrytis cinerea*. *Plant Dis.* 97 (1), 118–122.
- VKM Report, 2014. Health Risk Assessment of the Fungicide Aviator Xpro EC 225 - Vitenskapskomiteen for Mat Og Miljø, pp. 1–19.
- VKM Report, 2015. Risk Assessment of the Metabolite M44 of Bixafen, One of the Active Substances in Aviator Xpro EC 225 - Vitenskapskomiteen for Mat Og Miljø, pp. 1–20.
- Vu, Hung T., Keough, Michael J., Long, Sara M., Pettigrove, Vincent J., 2016. Effects of the boscalid fungicide Filan® on the marine amphipod *Allorchestes compressa* at environmentally relevant concentrations. *Environ. Toxicol. Chem.* 35 (5), 1130–1137.
- Wang, Honglei, Meng, Zhen, Liu, Fasheng, Zhou, Liqun, Su, Meile, Meng, Yunlong, Zhang, Shouhua, Liao, Xinjun, Cao, Zigang, Lu, Huiqiang, 2020. Characterization of boscalid-induced oxidative stress and neurodevelopmental toxicity in zebrafish embryos. *Chemosphere* 238.
- Weichenthal, Scott, Moase, Connie, Chan, Peter, 2010. A review of pesticide exposure and cancer incidence in the agricultural health study cohort. *Environ. Health Perspect.* 118 (8), 1117–1125.
- Wu, Siyu, Lei, Lili, Liu, Mengting, Song, Yang, Lu, Shibo, Li, Dan, Shi, Huahong, Kathleen, M., Raley-Susman, He, Defu, 2018. Single and Mixture Toxicity of Strobilurin and SDHI Fungicides to *Xenopus Tropicalis* Embryos.
- Yamashita, Masao, Fraaije, Bart, 2018. Non-target site SDHI resistance is present as standing genetic variation in field Populations of *Zymoseptoria tritici*. *Pest Manag. Sci.* 74 (3), 672–681.
- Yang, Yang, Dong, Fengshou, Liu, Xingang, Xu, Jun, Wu, Xiaohu, Qi, Suzhen, Liu, Wenxian, Zheng, Yongquan, 2018. Thifluzamide affects lipid metabolism in zebrafish (*Danio reio*). *Sci. Total Environ.* 633, 1227–1236.
- Yao, Hongzhou, Yu, Jianping, Zhou, Ying, Xiang, Qingqing, Xu, Chao, 2018. The Embryonic Developmental Effect of Sedaxane on Zebrafish (*Danio Rerio*).
- Zhang, A., Yue, Ying, Yang, Y., Yang, Jian, Tao, Ke, Jin, Hong, Hou, Taiping, 2019. Discovery of N-(4-Fluoro-2-(Phenylamino)Phenyl)-Pyrazole-4-Carboxamides as potential succinate dehydrogenase inhibitors. *Pestic. Biochem. Physiol.* 158, 175–184.