

# Brain changes associated with sleep disruption in cognitively unimpaired older adults: a short review of neuroimaging studies

Claire André, Alice Laniepce, Gaël Chételat, Géraldine Rauchs

#### ▶ To cite this version:

Claire André, Alice Laniepce, Gaël Chételat, Géraldine Rauchs. Brain changes associated with sleep disruption in cognitively unimpaired older adults: a short review of neuroimaging studies. Ageing Research Reviews - ARR, 2021, pp.101252. 10.1016/j.arr.2020.101252. inserm-03124348

### HAL Id: inserm-03124348 https://inserm.hal.science/inserm-03124348v1

Submitted on 28 Jan 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Title page

2	
3	Brain changes associated with sleep disruption in cognitively
4	unimpaired older adults: a short review of neuroimaging studies.
5	
6	Claire André <sup>1,2</sup> , Alice Laniepce <sup>1</sup> , Gaël Chételat <sup>2</sup> , Géraldine Rauchs <sup>1</sup> *
7	
8	<sup>1</sup> Normandie Univ, UNICAEN, PSL Université, EPHE, INSERM, U1077, CHU de Caen, GIF
9	Cyceron, NIMH « Neuropsychologie et Imagerie de la Mémoire Humaine », Caen, France.
10	<sup>2</sup> Normandie Univ, UNICAEN, INSERM, U1237, PhIND "Physiopathology and Imaging of
11	Neurological Disorders", Institut Blood and Brain @ Caen-Normandie, Cyceron, Caen
12	France.
13	* Corresponding author.
14	
15	
16	
17	Corresponding author:
18	Géraldine Rauchs, PhD
19	Inserm-EPHE-UNICAEN U1077 NIMH,
20	GIP Cyceron,
21	Bd Henri Becquerel, BP 5229,
22	14074 CAEN cedex 5,
23	France
24	geraldine.rauchs@inserm.fr
25	

Running title: Ageing, sleep and brain integrity.
Keywords: Sleep, Ageing, Alzheimer's disease, Neuroimaging, Amyloid, Tau.
Word count: 3011 words

#### 34 **Highlights:**

- Sleep disruption is increasingly considered as a risk factor for Alzheimer's disease.
- Poor sleep is associated with diffuse frontal, temporal and parietal gray matter atrophy.
- Alzheimer's disease biomarkers are associated with several sleep parameters.
- However, the specificity, topography and causality of these links are unclear.

**Abstract:** Ageing is characterized by a progressive decline of sleep quality. Sleep difficulties are increasingly recognized as a risk factor for Alzheimer's disease (AD), and have been associated with cognitive decline. However, the brain substrates underlying this association remain unclear. In this review, our objective was to provide a comprehensive overview of the relationships between sleep changes and brain structural, functional and molecular integrity, including amyloid and tau pathologies in cognitively unimpaired older adults. We especially discuss the topography and causality of these associations, as well as the potential underlying mechanisms. Taken together, current findings converge to a link between several sleep parameters, amyloid and tau levels in the CSF, and neurodegeneration in diffuse frontal, temporal and parietal areas. However, the existing literature remains heterogeneous, and the specific sleep changes associated with early AD pathological changes, in terms of topography and neuroimaging modality, is not clearly established yet. Notably, if slow wave sleep disruption seems to be related to frontal amyloid deposition, the brain correlates of sleepdisordered breathing and REM sleep disruption remains unclear. Moreover, sleep parameters associated with tau- and FDG-PET imaging are largely unexplored. Lastly, whether sleep disruption is a cause or a consequence of brain alterations remains an open question.

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

#### 1. Introduction

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

As the worldwide population is ageing, preventing cognitive decline and the development of dementia represents an important challenge. Intense research aims at identifying modifiable lifestyle factors that might help to promote healthy ageing. Among them, sleep is receiving particular attention, as about 50% of older adults complain of poor sleep (Foley et al., 1995). Age-related changes consist, at the circadian level, in advanced sleep timing and reduced amplitude of the circadian rhythms (Duffy et al., 2015; Kondratova and Kondratov, 2012). Sleep changes include longer sleep latency, decreased sleep duration, and increased number and duration of nocturnal awakenings, resulting in greater sleep fragmentation and lower sleep efficiency (Li et al., 2018; Ohayon et al., 2004). Remarkably, the amount of slow wave sleep (SWS), the deepest non-rapid eye movement (NREM) sleep stage, linearly decreases across the adult lifespan, contrasting with the increase of lighter NREM sleep (especially stage N1). Substantial changes in NREM sleep oscillations such as delta waves (Carrier et al., 2001; Landolt et al., 1996; Schwarz et al., 2017) and sleep spindles, key features of N2 sleep (Crowley et al., 2002; Schwarz et al., 2017) are also reported. Rapid-eye movement (REM) sleep time is reduced with age, albeit to a lesser extent and later in life than SWS (Floyd et al., 2007; Ohayon et al., 2004). Finally, 30 to 80% of older adults suffer from sleep-disordered breathing (SDB; Senaratna et al., 2017), a respiratory disorder defined by recurrent upper airway collapse during sleep, resulting in intermittent hypoxia episodes and sleep fragmentation, which may trigger neurodegenerative processes. Sleep is essential for an optimal daytime cognitive functioning, notably for attention, executive functioning, memory, and synaptic plasticity (Buzsáki, 1996; Killgore, 2010; Lowe et al., 2017; Tononi and Cirelli, 2006). In older adults, growing evidence show that sleep disturbances may represent a risk factor for cognitive decline and dementia. Indeed, longitudinal studies indicate that poor self-reported sleep quality, short but also long sleep duration (i.e., <7 or >8 hours of sleep), longer sleep latency, lower sleep efficiency, greater sleep fragmentation, decreased REM sleep, excessive daytime sleepiness and SDB are associated with cognitive decline and incident mild cognitive impairment or dementia diagnosis (Bubu et al., 2017; Diem et al., 2016; Foley et al., 2001; Gabelle et al., 2017; Leng et al., 2017; Lim et al., 2013a; Pase et al., 2017; Song et al., 2015; Virta et al., 2013). As Alzheimer's disease (AD) pathology develops several years before the first clinical symptoms (Jack et al., 2018), it is crucial to better understand the associations between sleep disturbances and brain integrity (including AD biomarkers) in older adults who are still cognitively unimpaired. Thus, our objective was to summarize existing data about these associations, with an emphasis on their topography and causality, and discuss the potential underlying mechanisms. We focused the present short review on papers published during the last decade (i.e., between 2010 and November 2020), involving cognitively unimpaired older adults, with no history or current diagnosis of major neurological or psychiatric diseases, or any other major medical condition.

#### 2. Sleep quality and gray matter volume changes

Cross-sectional studies show that lower grey matter (GM) volume within frontal areas is one of the brain changes most consistently associated with sleep quality in cognitively unimpaired older adults (**Table 1**). Sleep parameters associated with frontal atrophy include self-reported poor sleep quality (Sexton et al., 2014) and inadequate sleep duration (Lo et al., 2014; Westwood et al., 2017), early-morning awakenings (Stoffers et al., 2012), excessive daytime sleepiness (Killgore et al., 2012), greater sleep fragmentation obtained from actigraphy (Lim et al., 2016), and decreased slow wave activity (Dubé et al., 2015; Latreille et al., 2019; Mander et al., 2013).

In addition, reduced GM volume in lateral and medial temporal regions (including the hippocampus and the amygdala), the thalamus, and parietal cortex, have been associated with poor self-reported sleep quality (Alperin et al., 2019; Liu et al., 2018), inadequate sleep duration (Lo et al., 2014; Westwood et al., 2017) and excessive daytime sleepiness (Carvalho et al., 2017). Branger et al. (2016) also found an association between a higher number of self-reported nocturnal awakenings and lower GM volume in the insula. Importantly, these findings are supported by studies using actigraphy or polysomnography. Indeed, SWS integrity has been associated with GM volume in parietal and insular cortices (Dubé et al., 2015), and reduced GM volume in medial temporal areas has been related to greater sleep-wake rhythm fragmentation (Van Someren et al., 2019). GM volume changes associated with SDB in older adults vary importantly, some studies showing atrophy (Huang et al., 2019; Shi et al., 2017; Tahmasian et al., 2016; Weng et al., 2014) in frontal, temporal and parietal areas, while others rather report hypertrophy (André et al., 2020; Baril et al., 2017; Cross et al., 2018; Rosenzweig et al., 2013) in the same brain areas. A few longitudinal studies reported that self-estimated poor sleep quality (Fjell et al., 2019; Sexton et al., 2014) and short sleep duration (Lo et al., 2014; Spira et al., 2016) are associated with an increased rate of atrophy within frontal, temporal (including the hippocampus) and parietal areas. However, long self-reported sleep duration (> 7h) has also been related to a higher rate of frontal GM atrophy (Spira et al., 2016).

125

126

127

128

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

#### 3. Sleep, brain glucose metabolism and perfusion changes

Only a few studies have explored the associations between sleep quality and resting-state glucose metabolism using <sup>18</sup>F-fluorodeoxyglucose (FDG) Positron Emission Tomography

(PET), or perfusion, using Arterial Spin Labeling (ASL) or Single Photon Emission Computed Tomography (SPECT) imaging (see **Table 1**).

In cognitively unimpaired older adults, if self-reported sleep measures were not associated to glucose metabolism (Branger et al., 2016), greater sleep fragmentation obtained from actigraphy was related to lower glucose metabolism within the ventromedial prefrontal cortex and the hippocampi (André et al., 2019). In addition, as for GM volume, SDB has been related to both lower (Baril et al., 2020, 2015; Innes et al., 2015; Kim et al., 2017; Nie et al., 2017) and greater (André et al., 2020; Baril et al., 2015; Nie et al., 2017) metabolism and/or perfusion mainly in frontal, temporal and parietal areas, including the precuneus and posterior cingulate cortex (**Table 1**).

#### 4. Sleep and amyloid pathology

In animal models of Alzheimer's disease and in humans, amyloid-ß peptide (Aß) levels fluctuate following a circadian pattern, increasing with wakefulness and decreasing during NREM sleep (Kang et al., 2009). Whether this effect is mainly attributable to decreased metabolic activity during SWS, and/or increased metabolites clearance through the "glymphatic system" (Tarasoff-Conway et al., 2015a; Xie et al., 2013) is still debated. Nevertheless, the circadian fluctuations of Aß levels in the cerebrospinal fluid (CSF) are attenuated with ageing and with increased Aß deposition (Huang et al., 2012; Roh et al., 2012). In cognitively unimpaired older adults, various sleep parameters have been associated with greater global Aß levels measured in the CSF or using PET, including poorer self-reported sleep quality, both cross-sectionally (Sprecher et al., 2017) and longitudinally (Fjell et al., 2018), longer subjective (Brown et al., 2016) and objective (Ettore et al., 2019) sleep latency, both

insufficient and long self-reported sleep duration (Spira et al., 2013; Xu et al., 2020), lower sleep efficiency (Ettore et al., 2019; Ju et al., 2013; Molano et al., 2017), increased sleep fragmentation (Ettore et al., 2019; Lucey et al., 2019; Wilckens et al., 2018), excessive daytime sleepiness (Xu et al., 2020), and altered slow wave activity (SWA) (Ju et al., 2017; Varga et al., 2016; Winer et al., 2019) (Table 1). Furthermore, Mander et al., (2015) showed that amyloid burden in medial prefrontal areas disrupts SWA, negatively affecting sleep-dependent memory consolidation. In addition, Winer et al. (2020) showed that decreased SWA and sleep efficiency significantly predicted the subsequent AB accumulation over several years. Studies using regional approaches have revealed that greater amyloid deposition in frontal and/or parietal areas, including the precuneus and posterior cingulate cortex, is associated with lower self-reported sleep quality (Sprecher et al., 2017), longer self-reported sleep latency (Branger et al., 2016), lower subjective sleep adequacy (Sprecher et al., 2015), corresponding to insufficient sleep quality and duration, and excessive daytime sleepiness, both crosssectionally (Sprecher et al., 2015) and longitudinally (Carvalho et al., 2018). Short sleep duration has been associated with greater amyloid deposition in the precuneus (Spira et al., 2013), but this result failed replication by Gabelle et al. (2019). Lastly, SBD has also been related to higher amyloid levels measured in the CSF or using PET imaging (both cross-sectionally and longitudinally) (Bu et al., 2015; Bubu et al., 2019; Jackson et al., 2020; Kong et al., 2020; Liguori et al., 2017; Sharma et al., 2018; Ylä-Herttuala et al., 2020), and specifically in the posterior cingulate cortex and the precuneus (André et al., 2020; Ylä-Herttuala

174

175

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

#### 5. Sleep and tau pathology

et al., 2020; Yun et al., 2017).

Besides Aß deposition, sleep disruption may also exacerbate tau pathology in older adults without cognitive deficits, the second pathophysiological hallmark of AD (Holth et al., 2019). Self-estimated poor sleep quality and excessive daytime sleepiness have been associated with higher t-tau and p-tau/Aβ42 ratios (Sprecher et al., 2017). In addition, Fjell et al. (2018) showed that CSF tau levels predict poor sleep quality in amyloid-positive older adults. Moreover, CSF tau levels increased after one night of sleep deprivation (Holth et al., 2019), and are associated with greater sleep fragmentation (Lim et al., 2013b) and lower sleep efficiency (Ju et al., 2017). Interestingly, in this latter study, the association between sleep and tau was mainly driven by increased neuronal activity during sleep. Moreover, patients with SDB also exhibit increased CSF tau levels, both cross-sectionally (Bu et al., 2015; Kong et al., 2020; Liguori et al., 2017) and longitudinally (Bubu et al., 2019). Recent studies using tau-PET imaging have provided mixed results (Table 1). Lucey et al. (2019) reported that both amyloid and tau pathologies are associated with decreased SWA, with a stronger link for tau. In contrast, Winer et al. (2019) did not report any direct association between tau in the medial temporal lobe and prefrontal SWA, but rather a relationship between medial temporal tau burden and altered coupling between slow waves and sleep spindles. However, research using tau-PET imaging is still in their infancy, and further studies are needed to unravel the associations between tau pathology and sleep quality.

194

195

196

197

198

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

#### 6. Discussion

Poor sleep quality in cognitively unimpaired older adults is associated with diffuse and heterogeneous structural, functional and molecular brain changes in frontal, temporal and parietal areas. While some of these brain substrates are consistent with early pathological

changes observed in AD, other sleep-associated brain changes seem less suggestive of AD, 199 200 both in terms of topography and nature of brain alterations involved. Indeed, frontal amyloid burden is associated with self-reported sleep difficulties (Branger et al., 201 2016; Sprecher et al., 2017, 2015). Moreover, SWS disruption appears to be robustly associated 202 203 with amyloid pathology (Ju et al., 2017; Varga et al., 2016), notably in medial prefrontal areas (Mander et al., 2015). Interestingly, AD is defined by the presence of amyloid deposition (Jack 204 et al., 2018) spreading from frontal areas, before the appearance of cognitive deficits. 205 Moreover, self-reported sleep difficulties (Alperin et al., 2019; Carvalho et al., 2017; Fjell et al., 206 2019) and greater fragmentation of the sleep/wake rhythm (Van Someren et al., 2019) have been 207 linked to medial temporal lobe (MTL) atrophy. Of note, SDB has been related to both reduced 208 (Huang et al., 2019; Tahmasian et al., 2016; Weng et al., 2014) and greater (Cross et al., 2018; 209 210 Rosenzweig et al., 2013) MTL volume. It seems crucial to better characterize sleep parameters 211 associated with MTL atrophy, as medial temporal areas are known to be affected by tau pathology and atrophied since AD pre-dementia stages (Braak and Braak, 1991; Villemagne and 212 213 Chételat, 2016). In addition, beyond the MTL, tau pathology also affects the brainstem, which is involved in sleep-wake regulation and REM sleep generation (Horner and Peever, 2017). 214 215 REM sleep is reduced in AD patients (Brayet et al., 2016; Hassainia et al., 1997; Hita-Yañez et al., 2012; Montplaisir et al., 1995) and this alteration is predictive of cognitive decline in older 216 adults (Pase et al., 2017). Liguori et al. (2020) have recently demonstrated that REM sleep 217 reduction is associated with increased CSF tau levels, in a cohort of cognitively healthy older 218 adults and patients with subjective cognitive decline, mild cognitive impairment and AD. 219 220 However, due to the relatively recent emergence of tau-PET neuroimaging, the relationships between REM sleep changes and tau pathology in the brainstem and medial temporal areas 221 222 remains to be clarified, specifically in cognitively normal older populations.

Lastly, the associations between sleep changes and glucose metabolism have been much less investigated. SDB has been associated with decreased metabolism and/or perfusion in posterior parietal areas (Baril et al., 2015; Yaouhi et al., 2009), but increases have also been reported (André et al., 2020). Taken together, it is unclear whether sleep quality is associated with the marked hypometabolism in posterior cingulate and precuneus areas observed in AD and mild cognitive impairment patients (Chételat et al., 2003; Minoshima et al., 1997; Schroeter et al., 2009).

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

223

224

225

226

227

228

229

Besides, sleep modifications have been linked to other brain changes which seem less directly suggestive of early AD pathological processes, and could rather reflect normal ageing processes. Indeed, GM atrophy in frontal and parietal areas is robustly associated with SWS disruption, in particular decreased slow wave activity (Dubé et al., 2015; Latreille et al., 2019; Mander et al., 2013). These results are consistent with the fact that these regions are involved in sleep physiology and the generation of slow waves (Massimini et al., 2004; Murphy et al., 2009). Moreover, greater sleep fragmentation is related to lower frontal GM volume (Lim et al., 2016) and fronto-hippocampal metabolism (André et al., 2019), two areas critical for sleepdependent memory consolidation (Buzsáki, 1996; Maingret et al., 2016). The brain correlates of SDB appear however still conflicting, with both positive and negative associations with frontal GM volume and metabolism, MTL metabolism, and parietal GM volume (see Table 1). The pattern of SDB-related neuronal hyperactivity (i.e., greater GM volume, metabolism and perfusion) reported in some studies could reflect acute and early reactive processes, likely due to neuroinflammation in response to hypoxia. This neuronal hyperactivity may also reflect compensatory mechanisms due to greater individual brain reserve, which may help to maintain cognitive performance in the normal range by increasing brain activity. Importantly, these acute changes may only be temporary and may trigger neurodegeneration (i.e., GM

atrophy and hypometabolism) and the development of cognitive deficits over time. 248 249 Longitudinal studies combining several neuroimaging modalities on large cohorts with various levels of SDB severity will be necessary to confirm this hypothesis. 250 Frontal, temporal and posterior parietal regions thus constitute common brain substrates 251 252 between sleep physiology and AD pathology, but a critical question is whether sleep disturbances are causal and/or consecutive to brain alterations. First, AD pathology located in 253 254 frontal, medial temporal and brainstem areas, involved in sleep rhythms, could disrupt sleep 255 quality, eventually impairing cognitive performance. Supporting this hypothesis, atrophy and 256 amyloid burden within medial prefrontal areas have been shown to disrupt SWS (Mander et 257 al., 2015, 2013), ultimately impairing sleep-dependent memory consolidation. Moreover, sleep 258 disruption may also mediate or moderate the association between brain alterations and cognitive performance. Indeed, poor self-estimated sleep quality (Molano et al., 2017) and 259 260 increased nocturnal wakefulness (Wilckens et al., 2018) moderate the association between amyloid burden and memory performance. Lastly, greater sleep fragmentation mediates the 261 association between fronto-hippocampal hypometabolism and poorer executive functioning 262 (André et al., 2019). 263 However, it is likely that the links between sleep and brain alterations are bidirectional (Ju et 264 al., 2014). Indeed, the consequence of disrupted sleep is to increase the amount of wakefulness 265 266 during sleep and to decrease the amount of sleep per se. Subsequently, greater neuronal 267 activity and reduced brain clearance function may result in greater amyloid and tau release (Fig. 1). If increased activity and reduced clearance are likely to co-exist, their respective 268 contributions are unclear. A pilot study in humans showed that these links are mainly driven 269 by increased Aβ secretion, and not by decreased clearance (Lucey et al., 2018). However, the 270 271 study of brain clearance mechanisms is a recent area of research. In mice, clearance through

the blood brain barrier and interstitial fluid bulk-flow seem to be both implicated in amyloid

clearance and related to sleep quality (Rasmussen et al., 2018; Tarasoff-Conway et al., 2015b; Xie et al., 2013). Notably, the glymphatic system hypothesis proposes that arterial pulsations drive waste clearance through a directional convective flow of CSF, from periarterial areas in the deep brain parenchyma to the perivenous spaces. This process implicates water channels, like aquaporin-4, located in glial cells (liff et al., 2013, 2012; Nedergaard, 2013), and is facilitated during sleep (Xie et al., 2013). Nevertheless, several aspects of this model are currently debated (Mestre et al., 2020), and most findings in mice are pending replication in humans, as non-invasive neuroimaging tools of clearance mechanisms are still under development. Moreover, the associations with AD pathology remain to be confirmed. In healthy young subjects, a promising recent study demonstrated that slow waves during NREM sleep are followed by hemodynamic oscillations, which are in turn coherent with large waves of CSF (Fultz et al., 2019).

#### 7. Conclusion

Age-related sleep disruption is associated with brain changes mainly in frontal, temporal and parietal areas, some of them being suggestive of early AD pathological changes. These links may explain why sleep disruption is related to lower cognitive performance and steeper cognitive decline. However, the specific aspects of sleep disruption involved in these associations, the impact on tau pathology, and the associations with disrupted brain clearance mechanisms during sleep remain to be further explored. In addition, if we only discussed the associations between sleep disruption and regional GM integrity and functioning, sleep-related changes in structural and functional connectivity specifically in cognitively unimpaired older adults are a promising area of research. From a clinical perspective, it is crucial to screen for sleep problems in older adults, as they may exacerbate AD

pathophysiological mechanisms and may represent a modifiable lifestyle factor contributing 297 298 to healthy ageing. 299 8. Competing interests 300 301 None declared. 302 9. References 303 Alperin, N., Wiltshire, J., Lee, S.H., Ramos, A.R., Hernandez-Cardenache, R., Rundek, T., Curiel Cid, R., 304 305 Loewenstein, D., 2019. Effect of sleep quality on amnestic mild cognitive impairment vulnerable 306 brain regions in cognitively normal elderly individuals. Sleep 42. 307 https://doi.org/10.1093/sleep/zsy254 308 André, C., Rehel, S., Kuhn, E., Landeau, B., Moulinet, I., Touron, E., Ourry, V., Le Du, G., Mézenge, F., 309 Tomadesso, C., de Flores, R., Bejanin, A., Sherif, S., Delcroix, N., Manrique, A., Abbas, A., 310 Marchant, N.L., Lutz, A., Klimecki, O.M., Collette, F., Arenaza-Urquijo, E.M., Poisnel, G., Vivien, D., Bertran, F., de la Sayette, V., Chételat, G., Rauchs, G., Medit-Ageing Research Group, 2020. 311 312 Association of Sleep-Disordered Breathing With Alzheimer Disease Biomarkers in Community-313 Dwelling Older Adults: A Secondary Analysis of a Randomized Clinical Trial. JAMA neurology e200311. https://doi.org/10.1001/jamaneurol.2020.0311 314 315 André, C., Tomadesso, C., de Flores, R., Branger, P., Rehel, S., Mézenge, F., Landeau, B., Sayette, V. de 316 la, Eustache, F., Chételat, G., Rauchs, G., 2019. Brain and cognitive correlates of sleep 317 fragmentation in elderly subjects with and without cognitive deficits. Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring 11, 142-150. 318 319 https://doi.org/10.1016/j.dadm.2018.12.009 320 Baril, A.-A., Gagnon, K., Arbour, C., Soucy, J.-P., Montplaisir, J., Gagnon, J.-F., Gosselin, N., 2015. 321 Regional Cerebral Blood Flow during Wakeful Rest in Older Subjects with Mild to Severe 322 Obstructive Sleep Apnea. Sleep 38, 1439–49. https://doi.org/10.5665/sleep.4986 323 Baril, A.A., Gagnon, K., Brayet, P., Montplaisir, J., Carrier, J., Soucy, J.P., Lafond, C., Blais, H., d'Aragon, 324 C., Gagnon, J.F., Gosselin, N., 2020. Obstructive sleep apnea during REM sleep and daytime cerebral functioning: A regional cerebral blood flow study using high-resolution SPECT. Journal 325 326 of Cerebral Blood Flow and Metabolism 40, 1230-1241. 327 https://doi.org/10.1177/0271678X18814106 328 Baril, A.-A., Gagnon, K., Brayet, P., Montplaisir, J., De Beaumont, L., Carrier, J., Lafond, C., L'Heureux, 329 F., Gagnon, J.-F., Gosselin, N., 2017. Gray Matter Hypertrophy and Thickening with Obstructive Sleep Apnea in Middle-aged and Older Adults. American journal of respiratory and critical care 330 331 medicine 195, 1509–1518. https://doi.org/10.1164/rccm.201606-12710C

332 333	Braak, H., Braak, E., 1991. Neuropathological stageing of Alzheimer-related changes. Acta neuropathologica 82, 239–59. https://doi.org/10.1007/BF00308809
334 335 336 337	Branger, P., Arenaza-Urquijo, E.M., Tomadesso, C., Mézenge, F., André, C., de Flores, R., Mutlu, J., de La Sayette, V., Eustache, F., Chételat, G., Rauchs, G., 2016. Relationships between sleep quality and brain volume, metabolism, and amyloid deposition in late adulthood. Neurobiology of aging 41, 107–114. https://doi.org/10.1016/j.neurobiologing.2016.02.009
338 339 340	Brayet, P., Petit, D., Frauscher, B., Gagnon, J.F., Gosselin, N., Gagnon, K., Rouleau, I., Montplaisir, J., 2016. Quantitative EEG of Rapid-Eye-Movement Sleep: A Marker of Amnestic Mild Cognitive Impairment. Clinical EEG and Neuroscience. https://doi.org/10.1177/1550059415603050
341 342 343 344	Brown, B.M., Rainey-Smith, S.R., Villemagne, V.L., Weinborn, M., Bucks, R.S., Sohrabi, H.R., Laws, S.M., Taddei, K., Macaulay, S.L., Ames, D., Fowler, C., Maruff, P., Masters, C.L., Rowe, C.C., Martins, R.N., AIBL Research Group, 2016. The Relationship between Sleep Quality and Brain Amyloid Burden. Sleep 39, 1063–1068. https://doi.org/10.5665/sleep.5756
345 346 347	Bu, XL., Liu, YH., Wang, QH., Jiao, SS., Zeng, F., Yao, XQ., Gao, D., Chen, JC., Wang, YJ., 2015. Serum amyloid-beta levels are increased in patients with obstructive sleep apnea syndrome. Scientific Reports 5, 13917. https://doi.org/10.1038/srep13917
348 349 350 351	Bubu, O.M., Brannick, M., Mortimer, J., Umasabor-Bubu, O., Sebastião, Y. V, Wen, Y., Schwartz, S., Borenstein, A.R., Wu, Y., Morgan, D., Anderson, W.M., 2017. Sleep, Cognitive impairment, and Alzheimer's disease: A Systematic Review and Meta-Analysis. Sleep 40. https://doi.org/10.1093/sleep/zsw032
352 353 354 355 356 357	Bubu, O.M., Pirraglia, E., Andrade, A.G., Sharma, R.A., Gimenez-Badia, S., Umasabor-Bubu, O.Q., Hogan, M.M., Shim, A.M., Mukhtar, F., Sharma, N., Mbah, A.K., Seixas, A.A., Kam, K., Zizi, F., Borenstein, A.R., Mortimer, J.A., Kip, K.E., Morgan, D., Rosenzweig, I., Ayappa, I., Rapoport, D.M., Jean-Louis, G., Varga, A.W., Osorio, R.S., Alzheimer's Disease Neuroimaging Initiative, 2019. Obstructive sleep apnea and longitudinal Alzheimer's disease biomarker changes. Sleep 42. https://doi.org/10.1093/sleep/zsz048
358 359	Buzsáki, G., 1996. The Hippocampo-Neocortical Dialogue. Cerebral Cortex 6, 81–92. https://doi.org/10.1093/cercor/6.2.81
360 361 362	Carrier, J., Land, S., Buysse, D.J., Kupfer, D.J., Monk, T.H., 2001. The effects of age and gender on sleep EEG power spectral density in the middle years of life (ages 20-60 years old). Psychophysiology 38, 232–42. https://doi.org/10.1111/1469-8986.3820232
363 364 365 366 367	Carvalho, D.Z., St. Louis, E.K., Boeve, B.F., Mielke, M.M., Przybelski, S.A., Knopman, D.S., Machulda, M.M., Roberts, R.O., Geda, Y.E., Petersen, R.C., Jack, C.R., Vemuri, P., 2017. Excessive daytime sleepiness and fatigue may indicate accelerated brain aging in cognitively normal late middle-aged and older adults. Sleep Medicine 32, 236–243. https://doi.org/10.1016/j.sleep.2016.08.023
368 369 370 371 372	Carvalho, D.Z., St Louis, E.K., Knopman, D.S., Boeve, B.F., Lowe, V.J., Roberts, R.O., Mielke, M.M., Przybelski, S.A., Machulda, M.M., Petersen, R.C., Jack, C.R., Vemuri, P., 2018. Association of Excessive Daytime Sleepiness With Longitudinal β-Amyloid Accumulation in Elderly Persons Without Dementia. JAMA Neurology 75, 672–680. https://doi.org/10.1001/jamaneurol.2018.0049

373 374 375	Chételat, G., Desgranges, B., de la Sayette, V., Viader, F., Eustache, F., Baron, JC., 2003. Mild cognitive impairment: Can FDG-PET predict who is to rapidly convert to Alzheimer's disease? Neurology 60, 1374–7. https://doi.org/10.1212/01.wnl.0000055847.17752.e6
376 377 378 379	Cross, N.E., Memarian, N., Duffy, S.L., Paquola, C., LaMonica, H., D'Rozario, A., Lewis, S.J.G., Hickie, I.B., Grunstein, R.R., Naismith, S.L., 2018. Structural brain correlates of obstructive sleep apnoea in older adults at risk for dementia. European Respiratory Journal 52, 1800740. https://doi.org/10.1183/13993003.00740-2018
380 381 382 383	Crowley, K., Trinder, J., Kim, Y., Carrington, M., Colrain, I.M., 2002. The effects of normal aging on sleep spindle and K-complex production. Clinical neurophysiology: official journal of the International Federation of Clinical Neurophysiology 113, 1615–22. https://doi.org/10.1016/S1388-2457(02)00237-7
384 385 386 387	Diem, S.J., Blackwell, T.L., Stone, K.L., Yaffe, K., Tranah, G., Cauley, J.A., Ancoli-Israel, S., Redline, S., Spira, A.P., Hillier, T.A., Ensrud, K.E., 2016. Measures of Sleep–Wake Patterns and Risk of Mild Cognitive Impairment or Dementia in Older Women. The American Journal of Geriatric Psychiatry 24, 248–258. https://doi.org/10.1016/j.jagp.2015.12.002
388 389 390	Dubé, J., Lafortune, M., Bedetti, C., Bouchard, M., Gagnon, J.F., Doyon, J., Evans, A.C., Lina, JM., Carrier, J., 2015. Cortical Thinning Explains Changes in Sleep Slow Waves during Adulthood. Journal of Neuroscience 35, 7795–7807. https://doi.org/10.1523/JNEUROSCI.3956-14.2015
391 392	Duffy, J.F., Zitting, KM., Chinoy, E.D., 2015. Aging and Circadian Rhythms. Sleep Medicine Clinics 10, 423–434. https://doi.org/10.1016/j.jsmc.2015.08.002
393 394 395 396	Ettore, E., Bakardjian, H., Solé, M., Levy Nogueira, M., Habert, MO., Gabelle, A., Dubois, B., Robert, P., David, R., 2019. Relationships between objectives sleep parameters and brain amyloid load in subjects at risk to Alzheimer's disease: the INSIGHT-preAD Study. Sleep. https://doi.org/10.1093/sleep/zsz137
397 398 399 400	Fjell, A.M., Idland, AV., Sala-Llonch, R., Watne, L.O., Borza, T., Brækhus, A., Lona, T., Zetterberg, H., Blennow, K., Wyller, T.B., Walhovd, K.B., 2018. Neuroinflammation and Tau Interact with Amyloid in Predicting Sleep Problems in Aging Independently of Atrophy. Cerebral cortex (New York, N.Y.: 1991) 28, 2775–2785. https://doi.org/10.1093/cercor/bhx157
401 402 403 404 405 406	Fjell, A.M., Sørensen, Ø., Amlien, I.K., Bartrés-Faz, D., Bros, D.M., Buchmann, N., Demuth, I., Drevon, C.A., Düzel, S., Ebmeier, K.P., Idland, AV., Kietzmann, T.C., Kievit, R., Kühn, S., Lindenberger, U., Mowinckel, A.M., Nyberg, L., Price, D., Sexton, C.E., Solé-Padullés, C., Pudas, S., Sederevicius, D., Suri, S., Wagner, G., Watne, L.O., Westerhausen, R., Zsoldos, E., Walhovd, K.B., 2019. Self-reported sleep relates to hippocampal atrophy across the adult lifespan - results from the Lifebrain consortium. Sleep. https://doi.org/10.1093/sleep/zsz280
407 408	Floyd, J.A., Janisse, J.J., Jenuwine, E.S., Ager, J.W., 2007. Changes in REM-sleep percentage over the adult lifespan. Sleep 30, 829–36. https://doi.org/10.1093/sleep/30.7.829
409	Foley, D., Monjan, A., Masaki, K., Ross, W., Havlik, R., White, L., Launer, L., 2001. Daytime sleepiness

is associated with 3-year incident dementia and cognitive decline in older Japanese-American

men. Journal of the American Geriatrics Society 49, 1628–32. https://doi.org/10.1111/j.1532-

410

411 412

5415.2001.49271.x

413 414 415	Foley, D.J., Monjan, A.A., Brown, S.L., Simonsick, E.M., Wallace, R.B., Blazer, D.G., 1995. Sleep complaints among elderly persons: an epidemiologic study of three communities. Sleep 18, 425–32. https://doi.org/10.1093/sleep/18.6.425
416 417 418	Fultz, N.E., Bonmassar, G., Setsompop, K., Stickgold, R.A., Rosen, B.R., Polimeni, J.R., Lewis, L.D., 2019. Coupled electrophysiological, hemodynamic, and cerebrospinal fluid oscillations in human sleep. Science (New York, N.Y.) 366, 628–631. https://doi.org/10.1126/science.aax5440
419 420 421 422 423	Gabelle, A., Gutierrez, L.A., Jaussent, I., Ben Bouallegue, F., De Verbizier, D., Navucet, S., Grasselli, C., Bennys, K., Marelli, C., David, R., Mariano-Goulart, D., Andrieu, S., Vellas, B., Payoux, P., Berr, C., Dauvilliers, Y., 2019. Absence of Relationship Between Self-Reported Sleep Measures and Amyloid Load in Elderly Subjects. Frontiers in Neurology 10, 989. https://doi.org/10.3389/fneur.2019.00989
424 425 426 427	Gabelle, A., Gutierrez, LA., Jaussent, I., Navucet, S., Grasselli, C., Bennys, K., Marelli, C., David, R., Andrieu, S., Berr, C., Vellas, B., Dauvilliers, Y., 2017. Excessive Sleepiness and Longer Nighttime in Bed Increase the Risk of Cognitive Decline in Frail Elderly Subjects: The MAPT-Sleep Study. Frontiers in Aging Neuroscience 9, 312. https://doi.org/10.3389/fnagi.2017.00312
428 429 430	Hassainia, F., Petit, D., Nielsen, T., Gauthier, S., Montplaisir, J., 1997. Quantitative EEG and Statistical Mapping of Wakefulness and REM Sleep in the Evaluation of Mild to Moderate Alzheimer's Disease. European Neurology 37, 219–224. https://doi.org/10.1159/000117446
431 432 433	Hita-Yañez, E., Atienza, M., Gil-Neciga, E., Cantero, J.L., 2012. Disturbed sleep patterns in elders with mild cognitive impairment: the role of memory decline and ApoE ε4 genotype. Current Alzheimer research 9, 290–7. https://doi.org/10.2174/156720512800107609
434 435 436 437	Holth, J.K., Fritschi, S.K., Wang, C., Pedersen, N.P., Cirrito, J.R., Mahan, T.E., Finn, M.B., Manis, M., Geerling, J.C., Fuller, P.M., Lucey, B.P., Holtzman, D.M., 2019. The sleep-wake cycle regulates brain interstitial fluid tau in mice and CSF tau in humans. Science (New York, N.Y.) 363, 880–884. https://doi.org/10.1126/science.aav2546
438 439	Horner, R.L., Peever, J.H., 2017. Brain Circuitry Controlling Sleep and Wakefulness. CONTINUUM Lifelong Learning in Neurology. https://doi.org/10.1212/CON.0000000000000495
440 441 442	Huang, X., Tang, S., Lyu, X., Yang, C., Chen, X., 2019. Structural and functional brain alterations in obstructive sleep apnea: a multimodal meta-analysis. Sleep Medicine 54, 195–204. https://doi.org/10.1016/j.sleep.2018.09.025
443 444 445 446	<ul> <li>Huang, Y., Potter, R., Sigurdson, W., Santacruz, A., Shih, S., Ju, YE., Kasten, T., Morris, J.C., Mintun, M., Duntley, S., Bateman, R.J., 2012. Effects of Age and Amyloid Deposition on Aβ Dynamics in the Human Central Nervous System. Archives of Neurology 69, 51. https://doi.org/10.1001/archneurol.2011.235</li> </ul>

- Iliff, J.J., Wang, M., Liao, Y., Plogg, B.A., Peng, W., Gundersen, G.A., Benveniste, H., Vates, G.E.,
   Deane, R., Goldman, S.A., Nagelhus, E.A., Nedergaard, M., 2012. A Paravascular Pathway
   Facilitates CSF Flow Through the Brain Parenchyma and the Clearance of Interstitial Solutes,
   Including Amyloid. Science Translational Medicine 4, 147ra111-147ra111.
- 451 https://doi.org/10.1126/scitranslmed.3003748
- 452 Iliff, J.J., Wang, M., Zeppenfeld, D.M., Venkataraman, A., Plog, B.A., Liao, Y., Deane, R., Nedergaard, 453 M., 2013. Cerebral Arterial Pulsation Drives Paravascular CSF-Interstitial Fluid Exchange in the

- 454 Murine Brain. Journal of Neuroscience 33, 18190–18199.
- 455 https://doi.org/10.1523/JNEUROSCI.1592-13.2013
- Innes, C.R.H., Kelly, P.T., Hlavac, M., Melzer, T.R., Jones, R.D., 2015. Decreased Regional Cerebral
- 457 Perfusion in Moderate-Severe Obstructive Sleep Apnoea during Wakefulness. Sleep 38, 699–
- 458 706. https://doi.org/10.5665/sleep.4658
- Jack, C.R., Bennett, D.A., Blennow, K., Carrillo, M.C., Dunn, B., Haeberlein, S.B., Holtzman, D.M.,
- Jagust, W., Jessen, F., Karlawish, J., Liu, E., Molinuevo, J.L., Montine, T., Phelps, C., Rankin, K.P.,
- 461 Rowe, C.C., Scheltens, P., Siemers, E., Snyder, H.M., Sperling, R., Contributors, 2018. NIA-AA
- 462 Research Framework: Toward a biological definition of Alzheimer's disease. Alzheimer's &
- dementia: the journal of the Alzheimer's Association 14, 535–562.
- 464 https://doi.org/10.1016/j.jalz.2018.02.018
- 465 Jackson, M.L., Cavuoto, M., Schembri, R., Doré, V., Villemagne, V.L., Barnes, M., O'Donoghue, F.J.,
- Rowe, C.C., Robinson, S.R., 2020. Severe Obstructive Sleep Apnea Is Associated with Higher
- 467 Brain Amyloid Burden: A Preliminary PET Imaging Study. Journal of Alzheimer's Disease 78, 1–7.
- 468 https://doi.org/10.3233/jad-200571
- Ju, Y.-E.S., Lucey, B.P., Holtzman, D.M., 2014. Sleep and Alzheimer disease pathology--a bidirectional
- 470 relationship. Nature Reviews Neurology 10, 115–9. https://doi.org/10.1038/nrneurol.2013.269
- Ju, Y.-E.S., McLeland, J.S., Toedebusch, C.D., Xiong, C., Fagan, A.M., Duntley, S.P., Morris, J.C.,
- 472 Holtzman, D.M., 2013. Sleep Quality and Preclinical Alzheimer Disease. JAMA Neurology 70,
- 473 587. https://doi.org/10.1001/jamaneurol.2013.2334
- Ju, Y.-E.S., Ooms, S.J., Sutphen, C., Macauley, S.L., Zangrilli, M.A., Jerome, G., Fagan, A.M., Mignot, E.,
- 475 Zempel, J.M., Claassen, J.A.H.R., Holtzman, D.M., 2017. Slow wave sleep disruption increases
- 476 cerebrospinal fluid amyloid-β levels. Brain 140, 2104–2111.
- 477 https://doi.org/10.1093/brain/awx148
- 478 Kang, J., Lim, M.M., Bateman, R.J., Lee, J.J., Smyth, L.P., Cirrito, J.R., Fujiki, N., Nishino, S., Holtzman,
- 479 D.M., Louis, S., 2009. Amyloid-β Dynamics are Regulated by Orexin and the Sleep-Wake Cycle.
- 480 Science 326, 1005–1007. https://doi.org/10.1126/science.1180962.Amyloid-
- 481 Killgore, W.D.S., 2010. Effects of sleep deprivation on cognition, in: Progress in Brain Research. pp.
- 482 105–129. https://doi.org/10.1016/B978-0-444-53702-7.00007-5
- 483 Killgore, W.D.S., Schwab, Z.J., Kipman, M., DelDonno, S.R., Weber, M., 2012. Voxel-based
- 484 morphometric gray matter correlates of daytime sleepiness. Neuroscience Letters 518, 10–13.
- 485 https://doi.org/10.1016/j.neulet.2012.04.029
- 486 Kim, J.S., Seo, J.H., Kang, M.-R., Seong, M.J., Lee, W.G., Joo, E.Y., Hong, S.B., 2017. Effect of
- continuous positive airway pressure on regional cerebral blood flow in patients with severe
- obstructive sleep apnea syndrome. Sleep Medicine 32, 122–128.
- 489 https://doi.org/10.1016/j.sleep.2016.03.010
- 490 Kondratova, A.A., Kondratov, R. V, 2012. The circadian clock and pathology of the ageing brain.,
- 491 Nature Reviews Neuroscience. https://doi.org/10.1038/nrn3208
- Kong, W., Zheng, Y., Xu, W., Gu, H., Wu, J., 2020. Biomarkers of Alzheimer's disease in severe
- obstructive sleep apnea–hypopnea syndrome in the Chinese population. European Archives of
- 494 Oto-Rhino-Laryngology. https://doi.org/10.1007/s00405-020-05948-2

- Landolt, H.P., Dijk, D.J., Achermann, P., Borbély, A.A., 1996. Effect of age on the sleep EEG: Slow-
- 496 wave activity and spindle frequency activity in young and middle-aged men. Brain Research
- 497 738, 205–12. https://doi.org/10.1016/S0006-8993(96)00770-6
- Latreille, V., Gaubert, M., Dubé, J., Lina, J.-M., Gagnon, J.-F., Carrier, J., 2019. Age-related cortical
- signatures of human sleep electroencephalography. Neurobiology of Aging 76, 106–114.
- 500 https://doi.org/10.1016/j.neurobiolaging.2018.12.012
- Leng, Y., McEvoy, C.T., Allen, I.E., Yaffe, K., 2017. Association of Sleep-Disordered Breathing With
- 502 Cognitive Function and Risk of Cognitive Impairment: A Systematic Review and Meta-analysis.
- 503 JAMA neurology 74, 1237–1245. https://doi.org/10.1001/jamaneurol.2017.2180
- Li, J., Vitiello, M. V, Gooneratne, N.S., 2018. Sleep in Normal Aging. Sleep medicine clinics 13, 1–11.
- 505 https://doi.org/10.1016/j.jsmc.2017.09.001
- Liguori, C., Mercuri, N.B., Izzi, F., Romigi, A., Cordella, A., Sancesario, G., Placidi, F., 2017. Obstructive
- 507 Sleep Apnea is Associated With Early but Possibly Modifiable Alzheimer's Disease Biomarkers
- 508 Changes. Sleep 40. https://doi.org/10.1093/sleep/zsx011
- Liguori, C., Placidi, F., Izzi, F., Spanetta, M., Mercuri, N.B., Di Pucchio, A., 2020. Sleep dysregulation,
- memory impairment, and CSF biomarkers during different levels of neurocognitive functioning
- in Alzheimer's disease course. Alzheimer's research & therapy 12, 5.
- 512 https://doi.org/10.1186/s13195-019-0571-3
- Lim, A.S.P., Fleischman, D.A., Dawe, R.J., Yu, L., Arfanakis, K., Buchman, A.S., Bennett, D.A., 2016.
- Regional Neocortical Gray Matter Structure and Sleep Fragmentation in Older Adults. Sleep 39,
- 515 227–235. https://doi.org/10.5665/sleep.5354
- 516 Lim, A.S.P., Kowgier, M., Yu, L., Buchman, A.S., Bennett, D.A., 2013a. Sleep Fragmentation and the
- 517 Risk of Incident Alzheimer's Disease and Cognitive Decline in Older Persons. Sleep 36, 1027–
- 518 1032. https://doi.org/10.5665/sleep.2802
- Lim, A.S.P., Yu, L., Kowgier, M., Schneider, J.A., Buchman, A.S., Bennett, D.A., 2013b. Modification of
- 520 the Relationship of the Apolipoprotein E ε4 Allele to the Risk of Alzheimer Disease and
- Neurofibrillary Tangle Density by Sleep. JAMA Neurology 70, 1544.
- 522 https://doi.org/10.1001/jamaneurol.2013.4215
- 523 Liu, Y.-R., Fan, D.-Q., Gui, W.-J., Long, Z.-L., Lei, X., Yu, J., 2018. Sleep-related brain atrophy and
- disrupted functional connectivity in older adults. Behavioural Brain Research 347, 292–299.
- 525 https://doi.org/10.1016/j.bbr.2018.03.032
- Lo, J.C., Loh, K.K., Zheng, H., Sim, S.K.Y., Chee, M.W.L., 2014. Sleep Duration and Age-Related
- 527 Changes in Brain Structure and Cognitive Performance. Sleep 37, 821–821.
- 528 https://doi.org/10.5665/sleep.3832
- 529 Lowe, C.J., Safati, A., Hall, P.A., 2017. The neurocognitive consequences of sleep restriction: A meta-
- analytic review. Neuroscience & Biobehavioral Reviews 80, 586–604.
- 531 https://doi.org/10.1016/j.neubiorev.2017.07.010
- Lucey, B.P., Hicks, T.J., McLeland, J.S., Toedebusch, C.D., Boyd, J., Elbert, D.L., Patterson, B.W., Baty,
- J., Morris, J.C., Ovod, V., Mawuenyega, K.G., Bateman, R.J., 2018. Effect of sleep on overnight
- cerebrospinal fluid amyloid β kinetics. Annals of Neurology 83, 197–204.
- 535 https://doi.org/10.1002/ana.25117

536 537 538 539	A.M., McCue, L., Xiong, C., Morris, J.C., Benzinger, T.L.S., Holtzman, D.M., 2019. Reduced non–rapid eye movement sleep is associated with tau pathology in early Alzheimer's disease. Science Translational Medicine 11, eaau6550. https://doi.org/10.1126/scitranslmed.aau6550
540 541 542	Maingret, N., Girardeau, G., Todorova, R., Goutierre, M., Zugaro, M., 2016. Hippocampo-cortical coupling mediates memory consolidation during sleep. Nature Neuroscience 19, 959–964. https://doi.org/10.1038/nn.4304
543 544 545 546	Mander, B.A., Marks, S.M., Vogel, J.W., Rao, V., Lu, B., Saletin, J.M., Ancoli-Israel, S., Jagust, W.J., Walker, M.P., 2015. β-amyloid disrupts human NREM slow waves and related hippocampus-dependent memory consolidation. Nature neuroscience 18, 1051–7. https://doi.org/10.1038/nn.4035
547 548 549	Mander, B.A., Rao, V., Lu, B., Saletin, J.M., Lindquist, J.R., Ancoli-Israel, S., Jagust, W., Walker, M.P., 2013. Prefrontal atrophy, disrupted NREM slow waves and impaired hippocampal-dependent memory in aging. Nature Neuroscience 16, 357–364. https://doi.org/10.1038/nn.3324
550 551 552	Massimini, M., Huber, R., Ferrarelli, F., Hill, S., Tononi, G., 2004. The Sleep Slow Oscillation as a Traveling Wave. Journal of Neuroscience 24, 6862–6870. https://doi.org/10.1523/JNEUROSCI.1318-04.2004
553 554	Mestre, H., Mori, Y., Nedergaard, M., 2020. The Brain's Glymphatic System: Current Controversies. Trends in Neurosciences 43, 458–466. https://doi.org/10.1016/j.tins.2020.04.003
555 556 557	Minoshima, S., Giordani, B., Berent, S., Frey, K.A., Foster, N.L., Kuhl, D.E., 1997. Metabolic reduction in the posterior cingulate cortex in very early Alzheimer's disease. Annals of Neurology 42, 85–94. https://doi.org/10.1002/ana.410420114
558 559 560	Molano, J.R. V, Roe, C.M., Ju, YE.S., 2017. The interaction of sleep and amyloid deposition on cognitive performance. Journal of Sleep Research 26, 288–292. https://doi.org/10.1111/jsr.12474
561 562 563	Montplaisir, J., Petit, D., Lorrain, D., Gauthier, S., Nielsen, T., 1995. Sleep in Alzheimer's disease: further considerations on the role of brainstem and forebrain cholinergic populations in sleepwake mechanisms. Sleep 18, 145–8. https://doi.org/10.1093/sleep/18.3.145
564 565 566	Murphy, M., Riedner, B.A., Huber, R., Massimini, M., Ferrarelli, F., Tononi, G., 2009. Source modeling sleep slow waves. Proceedings of the National Academy of Sciences 106, 1608–1613. https://doi.org/10.1073/pnas.0807933106
567 568	Nedergaard, M., 2013. Garbage Truck of the Brain. Science (New York, N.Y.) 340, 1529–1530. https://doi.org/10.1126/science.1240514
569 570 571	Nie, S., Peng, DC., Gong, HH., Li, HJ., Chen, LT., Ye, CL., 2017. Resting cerebral blood flow alteration in severe obstructive sleep apnoea: an arterial spin labelling perfusion fMRI study. Sleep and Breathing 21, 487–495. https://doi.org/10.1007/s11325-017-1474-9
572 573 574	Ohayon, M.M., Carskadon, M.A., Guilleminault, C., Vitiello, M. V, 2004. Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. Sleep 27, 1255–73. https://doi.org/10.1093/sleep/27.7.1255


- Pase, M.P., Himali, J.J., Grima, N.A., Beiser, A.S., Satizabal, C.L., Aparicio, H.J., Thomas, R.J., Gottlieb,
   D.J., Auerbach, S.H., Seshadri, S., 2017. Sleep architecture and the risk of incident dementia in
   the community. Neurology 89, 1244–1250. https://doi.org/10.1212/WNL.00000000000004373
- Rasmussen, M.K., Mestre, H., Nedergaard, M., 2018. The glymphatic pathway in neurological disorders. The Lancet. Neurology 17, 1016–1024. https://doi.org/10.1016/S1474-
- 580 4422(18)30318-1
- Roh, J.H., Huang, Y., Bero, A.W., Kasten, T., Stewart, F.R., Bateman, R.J., Holtzman, D.M., 2012.
- Disruption of the sleep-wake cycle and diurnal fluctuation of  $\beta$ -amyloid in mice with Alzheimer's
- disease pathology. Science Translational Medicine 4, 150ra122.
- 584 https://doi.org/10.1126/scitranslmed.3004291
- Rosenzweig, I., Kempton, M.J., Crum, W.R., Glasser, M., Milosevic, M., Beniczky, S., Corfield, D.R.,
  Williams, S.C., Morrell, M.J., 2013. Hippocampal Hypertrophy and Sleep Apnea: A Role for the
  Ischemic Preconditioning? PLoS ONE 8, e83173. https://doi.org/10.1371/journal.pone.0083173
- Schroeter, M.L., Stein, T., Maslowski, N., Neumann, J., 2009. Neural correlates of Alzheimer's disease
   and mild cognitive impairment: A systematic and quantitative meta-analysis involving 1351
   patients. NeuroImage 47, 1196–1206. https://doi.org/10.1016/j.neuroimage.2009.05.037
- Schwarz, J.F.A., Åkerstedt, T., Lindberg, E., Gruber, G., Fischer, H., Theorell-Haglöw, J., 2017. Age
   affects sleep microstructure more than sleep macrostructure. Journal of Sleep Research 26,
   277–287. https://doi.org/10.1111/jsr.12478
- Senaratna, C. V., Perret, J.L., Lodge, C.J., Lowe, A.J., Campbell, B.E., Matheson, M.C., Hamilton, G.S.,
   Dharmage, S.C., 2017. Prevalence of obstructive sleep apnea in the general population: A
   systematic review. Sleep Medicine Reviews 34, 70–81.
   https://doi.org/10.1016/j.smrv.2016.07.002
- Sexton, C.E., Storsve, A.B., Walhovd, K.B., Johansen-Berg, H., Fjell, A.M., 2014. Poor sleep quality is associated with increased cortical atrophy in community-dwelling adults. Neurology 83, 967–73. https://doi.org/10.1212/WNL.00000000000774
- Sharma, R.A., Varga, A.W., Bubu, O.M., Pirraglia, E., Kam, K., Parekh, A., Wohlleber, M., Miller, M.D.,
  Andrade, A., Lewis, C., Tweardy, S., Buj, M., Yau, P.L., Sadda, R., Mosconi, L., Li, Y., Butler, T.,
  Glodzik, L., Fieremans, E., Babb, J.S., Blennow, K., Zetterberg, H., Lu, S.E., Badia, S.G., Romero,
  S., Rosenzweig, I., Gosselin, N., Jean-Louis, G., Rapoport, D.M., de Leon, M.J., Ayappa, I., Osorio,
  R.S., 2018. Obstructive Sleep Apnea Severity Affects Amyloid Burden in Cognitively Normal
- 606 Elderly. A Longitudinal Study. American Journal of Respiratory and Critical Care Medicine 197,
- 933–943. https://doi.org/10.1164/rccm.201704-07040C
- Shi, Y., Chen, L., Chen, T., Li, L., Dai, J., Lui, S., Huang, X., Sweeney, J.A., Gong, Q., 2017. A Meta analysis of Voxel-based Brain Morphometry Studies in Obstructive Sleep Apnea. Scientific
   Reports 7, 10095. https://doi.org/10.1038/s41598-017-09319-6
- Song, Y., Blackwell, T., Yaffe, K., Ancoli-Israel, S., Redline, S., Stone, K.L., Osteoporotic Fractures in
   Men (MrOS) Study Group, 2015. Relationships Between Sleep Stages and Changes in Cognitive
   Function in Older Men: The MrOS Sleep Study. Sleep 38, 411–421.
- 614 https://doi.org/10.5665/sleep.4500
- Spira, A.P., Gamaldo, A.A., An, Y., Wu, M.N., Simonsick, E.M., Bilgel, M., Zhou, Y., Wong, D.F.,
  Ferrucci, L., Resnick, S.M., 2013. Self-reported sleep and β-amyloid deposition in community-

617 618	dwelling older adults. JAMA neurology 70, 1537–43. https://doi.org/10.1001/jamaneurol.2013.4258
619 620 621	Spira, A.P., Gonzalez, C.E., Venkatraman, V.K., Wu, M.N., Pacheco, J., Simonsick, E.M., Ferrucci, L., Resnick, S.M., 2016. Sleep Duration and Subsequent Cortical Thinning in Cognitively Normal Older Adults. Sleep 39, 1121–1128. https://doi.org/10.5665/sleep.5768
622 623 624 625	Sprecher, K.E., Bendlin, B.B., Racine, A.M., Okonkwo, O.C., Christian, B.T., Koscik, R.L., Sager, M.A., Asthana, S., Johnson, S.C., Benca, R.M., 2015. Amyloid burden is associated with self-reported sleep in nondemented late middle-aged adults. Neurobiology of Aging 36, 2568–2576. https://doi.org/10.1016/j.neurobiolaging.2015.05.004
626 627 628 629	Sprecher, K.E., Koscik, R.L., Carlsson, C.M., Zetterberg, H., Blennow, K., Okonkwo, O.C., Sager, M.A., Asthana, S., Johnson, S.C., Benca, R.M., Bendlin, B.B., 2017. Poor sleep is associated with CSF biomarkers of amyloid pathology in cognitively normal adults. Neurology 89, 445–453. https://doi.org/10.1212/WNL.00000000000001171
630 631 632 633	Stoffers, D., Moens, S., Benjamins, J., van Tol, MJ., Penninx, B.W.J.H., Veltman, D.J., Van der Wee, N.J.A., Van Someren, E.J.W., 2012. Orbitofrontal gray matter relates to early morning awakening: a neural correlate of insomnia complaints? Frontiers in neurology 3, 105. https://doi.org/10.3389/fneur.2012.00105
634 635 636 637	Tahmasian, M., Rosenzweig, I., Eickhoff, S.B., Sepehry, A.A., Laird, A.R., Fox, P.T., Morrell, M.J., Khazaie, H., Eickhoff, C.R., 2016. Structural and functional neural adaptations in obstructive sleep apnea: An activation likelihood estimation meta-analysis. Neuroscience & Biobehavioral Reviews 65, 142–156. https://doi.org/10.1016/j.neubiorev.2016.03.026
638 639 640 641	Tarasoff-Conway, J.M., Carare, R.O., Osorio, R.S., Glodzik, L., Butler, T., Fieremans, E., Axel, L., Rusinek, H., Nicholson, C., Zlokovic, B. V., Frangione, B., Blennow, K., Ménard, J., Zetterberg, H., Wisniewski, T., de Leon, M.J., 2015a. Clearance systems in the brain—implications for Alzheimer disease. Nature Reviews Neurology 11, 457–470. https://doi.org/10.1038/nrneurol.2015.119
642 643 644 645 646	Tarasoff-Conway, J.M., Carare, R.O., Osorio, R.S., Glodzik, L., Butler, T., Fieremans, E., Axel, L., Rusinek, H., Nicholson, C., Zlokovic, B. V., Frangione, B., Blennow, K., Ménard, J., Zetterberg, H., Wisniewski, T., de Leon, M.J., 2015b. Clearance systems in the brain—implications for Alzheimer disease. Nature Reviews Neurology 11, 457–470. https://doi.org/10.1038/nrneurol.2015.119
647 648	Tononi, G., Cirelli, C., 2006. Sleep function and synaptic homeostasis. Sleep Medicine Reviews 10, 49–62. https://doi.org/10.1016/j.smrv.2005.05.002
649 650 651 652 653	Van Someren, E.J.W., Oosterman, J.M., Van Harten, B., Vogels, R.L., Gouw, A.A., Weinstein, H.C., Poggesi, A., Scheltens, P., Scherder, E.J.A., 2019. Medial temporal lobe atrophy relates more strongly to sleep-wake rhythm fragmentation than to age or any other known risk. Neurobiology of learning and memory 160, 132–138. https://doi.org/10.1016/j.nlm.2018.05.017
654 655 656 657	Varga, A.W., Wohlleber, M.E., Giménez, S., Romero, S., Alonso, J.F., Ducca, E.L., Kam, K., Lewis, C., Tanzi, E.B., Tweardy, S., Kishi, A., Parekh, A., Fischer, E., Gumb, T., Alcolea, D., Fortea, J., Lleó, A., Blennow, K., Zetterberg, H., Mosconi, L., Glodzik, L., Pirraglia, E., Burschtin, O.E., de Leon, M.J., Rapoport, D.M., Lu, S., Ayappa, I., Osorio, R.S., 2016. Reduced Slow-Wave Sleep Is Associated

658 659	with High Cerebrospinal Fluid A $\beta$ 42 Levels in Cognitively Normal Elderly. Sleep 39, 2041–2048. https://doi.org/10.5665/sleep.6240
660 661	Villemagne, V.L., Chételat, G., 2016. Neuroimaging biomarkers in Alzheimer's disease and other dementias. Ageing Research Reviews 30, 4–16. https://doi.org/10.1016/j.arr.2016.01.004
662 663 664	Virta, J.J., Heikkilä, K., Perola, M., Koskenvuo, M., Räihä, I., Rinne, J.O., Kaprio, J., 2013. Midlife sleep characteristics associated with late life cognitive function. Sleep 36, 1533–41, 1541A. https://doi.org/10.5665/sleep.3052
665 666 667	Weng, HH., Tsai, Yuan-Hsiung, Chen, CF., Lin, YC., Yang, CT., Tsai, Ying-Huang, Yang, CY., 2014 Mapping Gray Matter Reductions in Obstructive Sleep Apnea: An Activation Likelihood Estimation Meta-Analysis. Sleep 37, 167–175. https://doi.org/10.5665/sleep.3330
668 669 670	Westwood, A.J., Beiser, A., Jain, N., Himali, J.J., DeCarli, C., Auerbach, S.H., Pase, M.P., Seshadri, S., 2017. Prolonged sleep duration as a marker of early neurodegeneration predicting incident dementia. Neurology 88, 1172–1179. https://doi.org/10.1212/WNL.000000000003732
671 672 673 674	Wilckens, K.A., Tudorascu, D.L., Snitz, B.E., Price, J.C., Aizenstein, H.J., Lopez, O.L., Erickson, K.I., Lopresti, B.J., Laymon, C.M., Minhas, D., Mathis, C.A., Buysse, D.J., Klunk, W.E., Cohen, A.D., 2018. Sleep moderates the relationship between amyloid beta and memory recall. Neurobiology of Aging 71, 142–148. https://doi.org/10.1016/j.neurobiolaging.2018.07.011
675 676 677 678	Winer, J.R., Mander, B.A., Helfrich, R.F., Maass, A., Harrison, T.M., Baker, S.L., Knight, R.T., Jagust, W.J., Walker, M.P., 2019. Sleep as a potential biomarker of tau and β-amyloid burden in the human brain. The Journal of neuroscience: the official journal of the Society for Neuroscience 0503–19. https://doi.org/10.1523/JNEUROSCI.0503-19.2019
679 680 681	Winer, J.R., Mander, B.A., Kumar, S., Reed, M., Baker, S.L., Jagust, W.J., Walker, M.P., 2020. Sleep Disturbance Forecasts β-Amyloid Accumulation across Subsequent Years. Current Biology 30, 4291-4298.e3. https://doi.org/10.1016/j.cub.2020.08.017
682 683 684 685	Xie, L., Kang, H., Xu, Q., Chen, M.J., Liao, Y., Thiyagarajan, M., O'Donnell, J., Christensen, D.J., Nicholson, C., Iliff, J.J., Takano, T., Deane, R., Nedergaard, M., 2013. Sleep Drives Metabolite Clearance from the Adult Brain. Science (New York, N.Y.) 342, 373–377. https://doi.org/10.1126/science.1241224
686 687 688 689	Xu, W., Tan, L., Su, B.J., Yu, H., Bi, Y.L., Yue, X.F., Dong, Q., Yu, J.T., 2020. Sleep characteristics and cerebrospinal fluid biomarkers of Alzheimer's disease pathology in cognitively intact older adults: The CABLE study. Alzheimer's and Dementia 16, 1146–1152. https://doi.org/10.1002/alz.12117
690 691 692	Yaouhi, K., Bertran, F., Clochon, P., Mézenge, F., Denise, P., Foret, J., Eustache, F., Desgranges, B., 2009. A combined neuropsychological and brain imaging study of obstructive sleep apnea. Journal of Sleep Research 18, 36–48. https://doi.org/10.1111/j.1365-2869.2008.00705.x
693 694 695 696 697	Ylä-Herttuala, S., Hakulinen, M., Poutiainen, P., Laitinen, T.M., Koivisto, A.M., Remes, A.M., Hallikainen, M., Lehtola, JM., Saari, T., Korhonen, V., Könönen, M., Vanninen, R., Mussalo, H., Laitinen, T., Mervaala, E., 2020. Severe Obstructive Sleep Apnea and Increased Cortical Amyloid-β Deposition. Journal of Alzheimer's disease: JAD. https://doi.org/10.3233/JAD-200736

698	Yun, CH., Lee, HY., Lee, S.K., Kim, H., Seo, H.S., Bang, S.A., Kim, S.E., Greve, D.N., Au, R., Shin, C.,
699	Thomas, R.J., 2017. Amyloid Burden in Obstructive Sleep Apnea. Journal of Alzheimer's Disease
700	59, 21–29. https://doi.org/10.3233/JAD-161047
701	
702	

#### **Figure 1**



<u>Figure 1.</u> Summary of the associations between sleep, brain and cognitive integrity in older adults.

<u>Abbreviations:</u> PFC: prefrontal cortex; MTL: medial temporal lobe; PCC: posterior cingulate cortex; REM-S: rapid eye movement sleep; SD: sleep deprivation; SDB: sleep-disordered breathing; SF: sleep fragmentation; SWS: slow wave sleep.

Studies show that poor sleep quality in older adults is associated with decreased gray matter volume in frontal, temporal and parietal areas, increased global, frontal, and parietal amyloid levels, as well as increased global tau levels. These brain areas are both sensitive to ageing and Alzheimer's disease, and some of them (notably frontal areas) are involved in the generation and maintenance of sleep oscillations. Their alteration contributes to the aggravation of amyloid and tau pathologies and may explain why poor sleep quality is associated with an increased risk of cognitive decline.

Sleep parameter	Method	Grey matter volume	Perfusion and glucose metabolism <sup>1</sup>	Amyloid pathology <sup>2</sup>	Tau pathology <sup>3</sup>
Sleep deprivation	Objective			Global levels (Ooms 2014)	Global levels (Holth 2019)
☑ Global sleep quality	Self- reported	Frontal (Sexton 2014), hippocampus (Alperin 2018, Liu 2018), thalamus (Liu 2018), parietal and amygdala (Alperin 2018).  Longitudinal: frontal, temporal and parietal (Sexton 2014), hippocampus (Fjell 2019)		Global levels (Sprecher 2017), frontal & parietal (Sprecher 2017)  Longitudinal: global levels (Fjell 2018)	Global levels (Sprecher 2017)  Longitudinal: global levels (Fjell 2018)
<b>↗</b> Excessive daytime sleepiness	Self- reported	MTL, frontal, parietal and occipital (Carvalho 2017)		Global levels (Xu 2020), frontal & parietal (Sprecher 2015).  Longitudinal: global levels (Spira 2018), anterior and posterior cingulate & precuneus (Carvalho 2018)	Global levels (Sprecher 2017)
	Self-			Global levels (Brown 2016), frontal	
<b>尽</b> Sleep latency	reported			(Branger 2016)	
	Objective			Global levels (Ettore 2019)	
☑ Sleep efficiency	Objective			Global levels (Ju 2013, Molano 2017, Ettore 2019) Longitudinal: global levels (Winer 2020)	Global levels (Ju 2017)
Short sleep duration	Self- reported	Global atrophy (Lo 2014).  Longitudinal: fronto-temporal (Spira 2016: <7h)		Global levels (Spira 2013, Xu 2020: ≤6h) & precuneus (Spira 2013: ≤6h)	
Long sleep duration	Self- reported	Global atrophy (Westwood 2017: ≥9h), occipital (Spira 2016: >7h). Longitudinal: frontal (Spira 2016: >7h)		Global levels (Xu 2020: ≥6h)	
7 Nocturnal awakenings & sleep	Self- reported	OFC (Stoffers 2012, early awakenings), insula (Branger 2016, nocturnal awakenings)			
fragmentation	Objective	Frontal (Lim 2016)	Frontal & hippocampus (André 2019)	Global levels (Wilckens 2018, Lucey 2019, Ettore 2019)	Longitudinal: global levels (Lim 2013)
Sleep-wake rhythm fragmentation	Objective	MTL (Van Someren 2018)			
ש Slow wave sleep ∠	Objective	Frontal (Mander 2013, Dubé 2015, Latreille 2019), parietal & insula (Dubé 2015)	PFC (Wilckens 2016)	Global levels (Ju 2017, Varga 2016, Winer 2019), PFC (Mander 2015) Longitudinal: global levels (Winer 2020)	Global levels (Lucey 2019, Liguori 2020)
☑ Sleep spindles	Objective				Global levels (Kam 2019: spindles density); MTL (Winer 2019: spindles/slow waves coupling)

<b>凶</b> REM sleep latency	Objective			Global levels (Lucey 2019)	
	Self- reported			Global levels (Elias 2018)	
SDB	Objective	Frontal (凶: Shi 2017, Huang 2019;  7: Baril 2017), MTL (ଧ: Weng 2014, Huang 2019, Tahmasian 2016; 7: Cross 2018, Rosenzweig 2013), temporo-parietal (凶: Weng 2014, Shi 2017), parieto-occipital (万: Baril 2017, Cross 2018, André 2020), insula (凶: Tahmasian 2016)	Parieto-occipital (凶: Yaouhi 2009, Baril 2015, Joo 2007, Kim 2017; ↗: André 2020), frontal (凶: Yaouhi 2009, Innes 2015, Kim 2017, Shiota 2014, Nie 2017, Baril 2020; ↗: Nie 2017), temporal & MTL (凶: Innes 2015, Kim 2017, Joo 2017, Nie 2017, Baril 2020), thalamus (凶: Innes 2015), insula (凶: Baril 2020; ↗: Baril 2015), basal ganglia & limbic regions (↗: Baril 2015)	Global levels (Osorio 2014, Bu 2015, Liguori 2017, Jackson 2020, Ylä- Herttuala 2020, Kong 2020), posterior cingulate, precuneus & temporal (Yun 2017, André 2020, Ylä-Herttuala 2020). Longitudinal: global levels (Sharma 2018, Bubu 2019)	Global levels (Ju 2016, Osorio 2014, Bu 2015, Liguori 2017, Motamedi 2018, Kong 2020).  Longitudinal: global levels (Bubu 2019)

- 721 <u>Table 1.</u> Summary of the associations between sleep parameters and brain integrity in older adults.
- Abbreviations: ASL: Arterial Spin Labeling, CSF: cerebrospinal fluid, FDG: <sup>18</sup>F-fluorodeoxyglucose, MTL: medial temporal lobe, OFC: orbitofrontal cortex,
- PET: Positrons Emission Tomography, PFC: prefrontal cortex, SDB: sleep-disordered breathing, SPECT: Single Photon Emission Computed Tomography.
- When meta-analyses (in bold) were available, original studies showing the same results were not mentioned. Objective sleep measures encompass actigraphy
- and polysomnography measures. Longitudinal studies are indicated in italics.
- 726 <sup>1</sup> Brain perfusion was measured using ASL (Innes et al., 2015; Nie et al., 2017) or SPECT imaging (Joo et al., 2007; Shiota et al., 2014; Baril et al., 2015,
- 727 2020; Kim et al., 2017), and brain glucose metabolism using FDG-PET (Yaouhi et al., 2009; Wilckens et al., 2016; André et al., 2019, 2020).
- 728 Amyloid levels were measured in the CSF (Ju et al., 2013; Ooms et al., 2014; Osorio et al., 2014; Varga et al., 2016; Liguori et al., 2017; Sprecher et al.,
- 729 2017; Molano et al., 2017; Ju et al., 2017; Fjell et al., 2018; Sharma et al., 2018; Bubu et al., 2019; Liguori et al., 2020; Kong 2020; Xu 2020), the blood (Bu
- 730 et al., 2015) or using PET imaging (Spira et al., 2013; Mander et al., 2015; Sprecher et al., 2015; Branger et al., 2016; Brown et al., 2016; Yun et al., 2017;
- 731 Elias et al., 2018; Wilckens et al., 2018; Spira et al., 2018; Carvalho et al., 2018; Lucey et al., 2019; Ettore et al., 2019; Winer et al., 2019, 2020; Bubu et al.,
- 732 2019, André et al., 2020; Jackson et al., 2020; Ylä-Herttuala et al., 2020).

<sup>3</sup> Tau levels were measured in the CSF (Osorio et al., 2014; Ju et al., 2016; Liguori et al., 2017; Sprecher et al., 2017; Ju et al., 2017; Fjell et al., 2018; Bubu et al., 2019; Holth et al., 2019; Kam et al., 2019; Liguori et al., 2020; Kong 2020), the blood (Bu et al., 2015; Motamedi et al., 2018), using PET imaging (Lucey et al., 2019; Winer et al., 2019), or at autopsy (Lim et al., 2013b).