



HAL
open science

Association between occupational exposure to irritant agents and a distinct asthma endotype in adults

Miora Valérie Andrianjafimasy, Mickaël Febrissy, Farid Zerimech, Brigitte Dananché, Hans Kromhout, Régis Matran, Mohamed Nadif, Dominique Oberson-Geneste, Catherine Quinot, Vivi Schlünssen, et al.

► To cite this version:

Miora Valérie Andrianjafimasy, Mickaël Febrissy, Farid Zerimech, Brigitte Dananché, Hans Kromhout, et al.. Association between occupational exposure to irritant agents and a distinct asthma endotype in adults. Occupational and Environmental Medicine, 2021, pp.oemed-2020-107065. 10.1136/oemed-2020-107065 . inserm-03331184

HAL Id: inserm-03331184

<https://inserm.hal.science/inserm-03331184>

Submitted on 1 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Title: Association between occupational exposure to irritant agents and a distinct asthma endotype in adults

Authors:

Miora Andrianjafimasy¹, Mickaël Febrissy², Farid Zerimech^{3,4,5}, Brigitte Dananché⁶, Hans Kromhout⁷, Régis Matran^{3,4,5}, Mohamed Nadif², Dominique Oberson-Geneste⁸, Catherine Quinot¹, Vivi Schlünssen^{9,10}, Valérie Siroux¹¹, Jan Paul Zock¹², Nicole Le Moual¹, Rachel Nadif¹, Oriane Dumas¹

Affiliations:

1. Université Paris-Saclay, UVSQ, Univ. Paris-Sud, Inserm, Équipe d'Épidémiologie respiratoire intégrative, CESP, 94807, Villejuif, France.
2. LIPADE, Université de Paris Descartes, Paris, France
3. Univ. Lille, ULR 4483 – IMPECS, F-59000 Lille, France
4. CHU Lille, F-59000 Lille, France
5. Institut Pasteur de Lille, F-59000 Lille, France
6. Industrial hygiene consulting, Courlaoux, France
7. Utrecht University, Division of Environmental Epidemiology, Institute for Risk Assessment Sciences, Utrecht, Netherlands
8. Toxibio-consultant, Lescar, France
9. Aarhus University, Department of Public Health, Environment, Occupation and Health, Danish Ramazzini Centre, Aarhus, Denmark
10. National Research Center for the Working Environment, Copenhagen, Denmark
11. Univ. Grenoble Alpes, Inserm, CNRS, Team of environmental epidemiology applied to Reproduction and Respiratory health, IAB, Grenoble, France.
12. Institute for Global Health (ISGlobal) and Universitat Pompeu Fabra (UPF), Barcelona, Spain

Corresponding author:

Orianne Dumas

Inserm, CESP, Équipe d'Épidémiologie respiratoire intégrative

16 avenue Paul Vaillant Couturier

94807 Villejuif

France

Phone number: 33 (0) 145 59 53 57

Email: orianne.dumas@inserm.fr

Contributorship statement:

The authors' contributions to the study were as follows: AM, RN and OD were involved in the conception, hypotheses delineation and design of the analysis strategy of the study; RM, VS, NLM and RN participated in the acquisition of the data; HK, JPZ, NLM and OD were involved in the update of the occupational asthma-specific job exposure matrix (OAsJEM); BD and DOG (experts), CQ and NLM were involved in the occupational exposure assessment in the EGEA survey; AM, MF, MN, and RN were involved in the identification of respiratory endotypes; AM and RN analysed the data; AM, RN and OD wrote the paper; MF, FZ, BD, HK, RM, MN, DOG, CQ, VS, VS, JPZ, and NLM reviewed the paper and revised it critically; All authors approved the final version of the manuscript.

Funding:

This work was funded by the Fonds AGIR pour les maladies chroniques, the French Agency for Food, Environmental and Occupational Health & Safety (ANSES/AFSSET, EST- 09–15, PNR-EST 2017), Merck Sharp & Dohme (MSD), the National hospital program of clinical research (PHRC-National 2012, EvAdA), the National Research Agency – Health Environment, Health-Work Program (ANR CES-2009), and the Région Hauts de France.

Competing interest:

No, there are no competing interests for any author.

Data availability:

Due to third party restrictions, EGEA data are not publicly available. Please see the following URL for more information: <https://egeanet.vjf.inserm.fr/index.php/en/contacts-en>.

Interested researchers should contact egea.cohorte@inserm.fr with further questions regarding data access.

Ethics approval statements: Human participants

Ethical approval was obtained from the relevant institutional review board committees (Cochin Port-Royal Hospital and Necker-Enfants Malades Hospital, Paris). Participants signed a written informed consent.

Acknowledgements:

EGEA cooperative group. Coordination: V Siroux (epidemiology, PI since 2013); F Demenais (genetics); I Pin (clinical aspects); R Nadif (biology); F Kauffmann (PI 1992-2012). Respiratory epidemiology: Inserm ex-U 700, Paris: M Korobaeff (Egea1), F Neukirch (Egea1); Inserm ex-U 707, Paris: I Annesi-Maesano (Egea1-2); Inserm U 1018, Villejuif: O Dumas, F Kauffmann, N Le Moual, R Nadif, MP Oryszczyn (Egea1-2), R Varraso; Inserm U 1209 Grenoble: J Lepeule, V Siroux.

Genetics: Inserm ex-U 393, Paris: J Feingold; Inserm UMR 1124, Paris: E Bouzigon, MH Dizier, F Demenais; CNG, Evry: I Gut (now CNAG, Barcelona, Spain), M Lathrop (now Univ McGill, Montreal, Canada). Clinical centers: Grenoble: I Pin, C Pison; Lyon: D Ecochard (Egea1), F Gormand, Y Pacheco; Marseille: D Charpin (Egea1), D Vervloet (Egea1-2);

Montpellier: J Bousquet; Paris Cochin: A Lockhart (Egea1), R Matran (now in Lille); Paris Necker: E Paty (Egea1-2), P Scheinmann (Egea1-2); Paris-Trousseau: A Grimfeld (Egea1-2), J Just. Data management and quality: Inserm ex-U155, Paris: J Hochez (Egea1); Inserm U 1018, Villejuif: N Le Moual, L Orsi; Inserm ex-U780, Villejuif: C Ravault (Egea1-2); Inserm ex-U794, Evry: N Chateigner (Egea1-2); Inserm UMR 1124, Paris: H Mohamdi; Inserm U1209, Grenoble: A Boudier, J Quentin (Egea1-2). The authors thank all those who participated to the setting of the study and on the various aspects of the examinations involved: interviewers, technicians for lung function testing and skin prick tests, blood sampling, IgE determinations, coders, those involved in quality control, data and sample management and all those who supervised the study in all centers. The authors are grateful to the three CIC-Inserm of Necker, Grenoble and Marseille who supported the study and in which participants were examined. They are also grateful to the biobanks in Lille (CIC Inserm), and at Annemasse (Etablissement français du sang) where biological samples are stored. They are indebted to all the individuals who participated, without whom the study would not have been possible.

We thank Theodore Lytras, Eva Andersson, Linnéa Lillienberg and Geza Benke for their contribution to the update of the asthma-specific job-exposure matrix (OAsJEM).

Abstract (259 words)

Aim: The biological mechanisms of work-related asthma induced by irritants remain unclear. We investigated the associations between occupational exposure to irritants and respiratory endotypes previously identified among never asthmatics (NA) and current asthmatics (CA) integrating clinical characteristics and biomarkers related to oxidative stress and inflammation.

Methods: We used cross-sectional data from 999 adults (mean 45 years old, 46% men) from the case-control and familial Epidemiological study on the Genetics and Environments of Asthma (EGEA) study. Five respiratory endotypes have been identified using a cluster-based approach: NA1 (n=463) asymptomatic, NA2 (n=169) with respiratory symptoms, CA1 (n=50) with active treated adult-onset asthma, poor lung function, high blood neutrophil counts and high Fluorescent Oxidation Products level, CA2 (n=203) with mild middle-age asthma, rhinitis and low Immunoglobulin E level, and CA3 (n=114) with inactive/mild untreated allergic childhood-onset asthma. Occupational exposure to irritants during the current or last held job was assessed by the updated Occupational Asthma-specific Job-Exposure Matrix (levels of exposure: no/medium/high). Associations between irritants and each respiratory endotype (NA1 asymptomatic as reference) were studied using logistic regressions adjusted for age, sex and smoking status.

Results: Prevalence of high occupational exposure to irritants was 7% in NA1, 6% in NA2, 16% in CA1, 7% in CA2 and 10% in CA3. High exposure to irritants was associated with CA1 (adjusted Odds Ratio aOR, [95% confidence interval]: 2.7 [1.0-7.3]). Exposure to irritants was not significantly associated with other endotypes (aOR range: 0.8-1.5).

Conclusion: Occupational exposure to irritants was associated with a distinct respiratory endotype suggesting oxidative stress and neutrophilic inflammation as potential associated biological mechanisms.

What this paper adds

What is already known about this subject?

- Epidemiological studies have suggested a role of repeated, chronic occupational exposure to low/moderate levels of irritant agents in causing asthma.
- Biological mechanisms by which exposure to irritant agents affect respiratory health remain unclear.

What are the new findings?

- High occupational exposure to irritants was associated with a respiratory endotype characterized by active treated adult-onset asthma, poor lung function, high blood neutrophil counts and high Fluorescent Oxidation Products level, a biomarker of damages related to oxidative stress.
- No other endotype was associated with irritants.

How might this impact on policy or clinical practice in the foreseeable future?

- Results provide additional knowledge on the biological mechanisms of irritant-induced asthma, suggesting oxidative stress and neutrophilic inflammation as potential mechanisms.
- If replicated, these findings may help improving the recognition and management of irritant-induced work-related asthma.

Introduction

Asthma is a heterogeneous chronic inflammatory disease encompassing several phenotypes that may have various risk factors including occupational exposures.[1,2] More than 500 workplace sensitizing or irritant agents have been identified as possible risk factors for asthma.[2,3] Work-related asthma is thus considered as a good model to study asthma in general.[4] Irritant-induced asthma remains poorly understood. Occupational asthma induced by irritants was first described as sudden onset of asthma after a single high peak of exposure to irritants.[5] In the last decade, several epidemiological studies have suggested a role of repeated, chronic exposure to low/moderate levels of irritant agents in causing asthma.[2,5,6] However, the biological mechanisms by which exposure to irritant agents affect respiratory health remain unclear.[3,5–7]

One of the biological mechanisms that could underlie irritant-induced asthma is oxidative stress that reflects the imbalance between reactive oxygen species (ROS) and antioxidant defenses in favor of the former.[2,8,9] Neutrophilic inflammation [2,3] has also been suggested. However, few studies have examined the role of these pathways in irritant-induced asthma, especially in human.[7,10]

Asthma heterogeneity has classically been approached by investigating different phenotypes, defined as a set of observable clinical characteristics.[11] Endotypes, *i.e.* disease subtypes characterized by a distinct functional or pathobiological mechanism [12] would allow to better understand the biological mechanisms associated with irritant-induced asthma.[11] In the Epidemiological study on the Genetics and Environments of Asthma (EGEA), we recently identified five respiratory endotypes using cluster analysis jointly integrating asthma clinical characteristics and biomarkers related to oxidative stress and inflammation.[13] In particular, we identified among asthmatics an endotype characterized by poor lung function, respiratory symptoms, high level of fluorescent oxidation products (FIOPs, a biomarker of damages related

to oxidative stress), and high blood neutrophil counts. We hypothesized occupational exposure to irritants may be associated to this specific endotype.

The updated Occupational Asthma-specific Job Exposure Matrix (OAsJEM),[14] of the former asthma-specific job exposure matrix,[15] has recently been published with improved assessment of occupational exposure, specifically for irritant agents. Taking advantage of these novel exposure data and the identification of respiratory endotypes in EGEA, we investigated the associations between occupational exposure to irritants and respiratory endotypes.

Methods

The EGEA study

EGEA is a French cohort started in 1990s with two follow-ups over 20 years. The first EGEA survey (EGEA1) included cases with asthma recruited in five chest clinics, their first-degree relatives and population-based controls (n=2047). A first follow-up of the participants was completed in 2003-2007 (EGEA2), including 1602 subjects with complete examination, almost exclusively adults (98%). At each survey, participants answered a standardized questionnaire on asthma, occupational history and environmental exposures. The protocol and participants' characteristics have been described previously,[16] and details are provided in online supplement. The EGEA collection was certified ISO 9001 (2006-2018) and is referenced in the Biobank network.[17] Ethical approval was obtained from the relevant institutional review board committees (Cochin Port-Royal Hospital and Necker-Enfants Malades Hospital, Paris). Participants signed a written informed consent.

At EGEA2, participants with “ever asthma” were those recruited as cases at EGEA1, or family members or controls who answered positively to one of two standardized questions. Among participants with ever asthma, current asthma was defined by respiratory symptoms, asthma attacks or treatment in the past 12 months (see online supplement).

Selection of the study population

The present cross-sectional study was based on data collected at the first follow-up (EGEA2). Participants less than 16 years old (n=31), those with ever asthma but without current asthma (n=125), or with missing data for any of the clinical and biological characteristics used to define endotypes (n=378) were excluded from the analyses. We also excluded participants who had never worked (n=60) or with missing data for occupational history (n=4). Finally, because our analysis focused on exposure to irritants, we excluded participants exposed only to occupational

sensitizers in the current or last held job (n=5, all only exposed to high molecular weight agents). In total, 999 participants were selected (online supplementary Figure E1).

Occupational exposure to irritant agents

Complete occupational history was collected at EGEA2 with information on job, industry and tasks. Jobs were coded according to the International Standard Classification of Occupation 1988 (ISCO-88) by an experienced coder. Occupational exposures to 30 agents at risk for asthma classified in seven groups were estimated using the OAsJEM (<http://oasjem.vjf.inserm.fr/index-en.htm>).[14] To improve the exposure estimate, the OAsJEM assessment was completed by an expert reassessment step for selected job codes defined *a priori* and regrouping jobs with heterogeneous tasks and/or industries. If the job description provided sufficient information, two experts (BD, DOG) reevaluated exposure levels case-by-case, independently of each other and blinded of asthma status. In case of disagreement between the experts, the final decision was taken by consensus (BD, DOG, CQ, NLM).

In the present study, we considered exposures to 19 agents known or suspected to cause asthma through irritant mechanisms, including chronic low-to-moderate level of exposures. Within this large group of irritant agents, two partly overlapping subgroups were further identified: highly reactive chemicals (eight agents) and biocides (five agents). In addition, we grouped three specific agents in a subgroup “cleaning products / disinfectants” (online supplementary Table E1). Among the 19 irritant agents, nine agents were also classified as low molecular weight (LMW) sensitizers, because for these nine agents both mechanisms have been suggested [14] (online supplementary Table E1).

The OAsJEM classified exposures to each irritant agent into three classes: “high” for high probability of exposure and moderate-to-high intensity, “medium” for low-to-moderate

probability or low intensity of exposure, and “no” for unlikely to be exposed. When analyzing groups of agents, the maximum exposure level among all agents in the group was considered. In all analyses, the reference group included participants classified as “non-exposed” to all of the 30 agents of the OAsJEM.

Definition of respiratory endotypes

Five respiratory endotypes were previously identified by cluster analysis [13] and are described in Table 1: two among never asthmatics (NA) and three among current asthmatics (CA), after taking into account a total of 23 variables for never asthmatics and 28 variables for current asthmatics. Cluster analysis jointly considered personal (age, sex, smoking status and BMI), clinical/functional (age of asthma onset, respiratory symptoms, asthma treatments, dyspnoea, skin prick tests positivity for at least one of 12 aeroallergens, current rhinitis, ever eczema, asthma attacks, hospital or emergency admission, forced expiratory volume in 1 second (FEV1) and forced vital capacity (FVC), and biological (neutrophil and eosinophil counts, total IgE and level of Fluorescent oxidation products) characteristics. More details on variables and cluster analysis used to define respiratory endotypes are provided in online supplement. A detailed description of all characteristics included in the cluster analysis is shown in online supplementary Table E2.

Statistical analyses

Chi-squared tests and Fisher exact tests were used to study the association between occupational exposure to irritant agents and each endotype in univariate models. Logistic regression models using generalized estimated equation (GEE) were performed to take into account familial dependence, and adjusted for age (continuous), sex and smoking status (non-smokers, ex- and current smokers). In the main analyses, irritant agents were estimated by OAsJEM completed

by the expert reassessment step. As clinical characteristics integrated in the definition of endotypes have been defined in the last 12 months and three endotypes have been defined among current asthmatics, exposures of interest were evaluated for the current or last held jobs. In a secondary analysis, we investigated the association between lifetime occupational exposure (instead of current exposure) to irritants and respiratory endotypes. In all models, the endotype of asymptomatic never asthmatics (NA1) was used as reference group for the outcome and “no exposure” to any OAsJEM agent was used as reference group for occupational exposure.

Three sensitivity analyses were carried out. First, we studied association between irritant agents estimated by OAsJEM only *i.e* without the expert reassessment step. Second, due to the partial overlap between the subgroups of irritants and LMW agents, we studied the associations between occupational exposures to LMW agents and endotypes. Third, the main analysis does not adjust for BMI as obesity has been described as an asthma comorbidity and asthma in obese patients may correspond to a specific asthma endotype.[18] However, as high BMI has also been described as a risk factor for asthma,[18] BMI was added to the other confounders in a sensitivity analysis.

Statistical analyses were performed using SAS statistical software (version 9.4; SAS Institute, Cary, NC, USA). A p-value of <0.05 was considered statistically significant.

Results

Characteristics of the study population

Table 2 shows the personal and clinical characteristics of the 999 participants and according to respiratory endotypes. The mean age of all participants was 45 years, 54% were women, 22% were current smokers and 11% were obese. Among endotypes in never asthmatics, participants in NA1 (n=463) had an average age of 47 years, 55% were women. In NA2 (n=169), 65% were women, 30% were current smokers and 35% were overweight. Among endotypes in current asthmatics, participants in CA1 (n=50) had an average age of 53 years, 12% were current smokers, 30% were overweight and 26% were obese. In CA2 (n=203), 33% were ex-smokers, and 33% were overweight. Participants in CA3 (n=114) were younger with an average age of 32 years, 62% were men and 33% were current smokers.

Occupational exposure to irritant agents in each respiratory endotype is described in Table E3 in online supplement. Among all participants, 72% had no exposure to irritants, 21% had medium exposure and 8% had high exposure. Regarding subgroups of irritants 4% of all participants had high exposure to highly reactive chemicals or biocides and 3% had high exposure to cleaning and disinfecting products. CA1 had the highest percentage of participants with high exposure to any irritant agents as well as each subgroup of irritants. The job titles of the eight participants in the CA1 endotype with high occupational exposure to irritants are described in Table E4 (online supplement). After the expert reassessment step, the number of participants classified with high or medium exposure slightly decreased while the number of participants classified as unexposed slightly increased (online supplementary Table E5).

Associations between occupational exposure to any irritant agents and respiratory endotypes

We observed a positive and significant association between high exposure to irritant agents and CA1 (vs. NA1, adjusted odds-ratio (aOR) [95% CI] = 2.70 [1.00 - 7.33], Table 3). No significant association was observed between medium exposure to irritants and CA1. We did not observe any significant association between any irritant agents and the other endotypes. Similar results were observed with exposure to irritants estimated without the expert reassessment step (aOR [95% CI] = 2.79 [1.19 - 6.56] for CA1 vs. NA1, online supplementary Table E6).

When we further adjusted for BMI, results remained similar with ORs > 2.5 for the association between irritants and CA1 (online supplementary Table E7). We did not observe a significant association between lifetime occupational exposure to irritants and CA1 nor the other endotypes (Table 4).

Associations between occupational exposure to subgroups of irritants and CA1 endotype

Despite the small sample size, we observed an association between high exposure to biocides and CA1 (aOR [95%CI] = 3.13 [1.00 - 9.83], online supplementary Table E8). Despite ORs higher than 2, there was no significant association between occupational exposure to highly reactive chemicals and cleaning products/disinfectants and CA1. Similar results were observed for subgroups of irritants estimated without the expert reassessment step. No significant association was observed between occupational exposure to LMW agents and CA1 (data not shown).

Discussion

The present study investigated the associations between occupational exposure to irritant agents and respiratory endotypes identified by a cluster analysis among never-asthmatics and among current asthmatics separately. We observed a significant association between current high exposure to irritant agents and an endotype predominantly characterized by adult-onset asthma, poor lung function, respiratory symptoms, high blood fluorescent oxidation product level and high neutrophil counts.

In our study, most of the asthmatics were recruited in chest clinics as asthma cases, with careful procedures set up to include true asthmatics, and others were mostly recruited as first-degree relatives of asthmatic cases, leading to the recruitment of participants with a wide range of asthma severity and control. The availability of key biomarkers related both to asthma and to inflammatory or oxidative stress pathways allowed identifying specific respiratory endotypes.[13] The characteristics included in the cluster analysis reflect as comprehensively as possible the participants' demographic, clinical and biological characteristics. We excluded variables missing for many participants such as the measurement of the exhaled fraction of nitric oxide (FeNO) which could have been useful to define the clusters. The EGEA study provides a relatively large sample of individuals with detailed characteristics on endotype (n=999). However, analyses by endotype resulted in smaller groups of individuals, in particular for the main endotype of interest (CA1, n=50), which is a limitation. Thus, although our findings are consistent with our *a priori* hypothesis of an association between occupational exposure to irritants and endotype CA1, they should be interpreted with caution and need to be confirmed in other studies. We used the recently updated OAsJEM [14] to improve exposure assessment. Indeed, the number of estimated agents has increased, ranging from 22 agents estimated by the previous asthma-specific JEM [15] to 30 agents at risk for asthma by the new

OAsJEM. These 30 agents have been classified in large groups distinguishing sensitizers from irritants which is of interest for studying their associations with distinct respiratory endotypes. We investigated the association between several subgroups of irritants (biocides, highly reactive chemicals and cleaning products / disinfectants) and the endotype CA1. Despite the significant association for high exposure to any irritants, we did not observe significant association with specific subgroups. The analyses of subgroups of irritants were limited by the small sample size and should therefore be interpreted with caution. Nonetheless, they were consistent with those for the large group of irritants with ORs of the same order of magnitude (all >2.0). These subgroups of irritants were associated with asthma in previous studies. Indeed, associations between occupational exposure to pesticides,[19,20] classified in the biocides group in our study, and to cleaning products/disinfectants [10,21,22] and asthma have been reported. Beside irritant mechanisms, other biological mechanisms could be involved: cleaning products/disinfectants are mixtures of substances that can be irritants or sensitizers [23] and pesticides have been associated with allergic asthma in some studies.[19] Studies with a larger sample size would be needed to precise the relationships for these subgroups and specific agents.

The association between current occupational exposure to irritants and CA1, and not with lifetime exposure might be related to the endotype definition. Indeed, since respiratory endotypes included clinical characteristics assessed over the last 12 months, reflecting the current activity of the disease, stronger associations were expected with current exposure rather than with past exposure. However, larger studies would be needed to study more accurately the temporal relationship between occupational exposures and endotypes. In addition, although CA1 included mostly participants with adult-onset asthma, we could not differentiate occupational asthma and work-exacerbated asthma, and the observed association may reflect both types of work-related asthma. We did not observe an association between occupational

exposure to LMW agents and CA1 nor the other endotypes. These results are consistent with the hypothesis of different biological mechanism for LMW agents, possibly involving a specific immune response, with or without specific IgE production.[24,25]

The OAsJEM evaluated chronic exposure to low/moderate levels of irritants, considered as non-accidental, and not peaks of exposure to irritants. The exposures estimated by the OAsJEM were completed by an expert reassessment step. Both the OAsJEM and the reassessment step have been developed by favoring specificity for high level of exposure rather than sensitivity.[14,26,27] This trade-off between sensitivity and specificity generally increases the positive predictive ability of a JEM.[26] However, it also reduces the number of subjects classified as exposed, thus possibly decreasing statistical power to detect associations. Nonetheless, our results were almost similar whatever the exposure estimation. Although the possibility of unmeasured confounding can never be ruled out, our analyses were adjusted for the confounders (age, sex, smoking status) usually used in the literature to study the relationships between occupational exposure and asthma, and further adjustment for BMI in a sensitivity analysis led to similar results. Analyses were not adjusted for education level or socioeconomic status as adding these variables may result in overadjustment in models investigating health effects of occupational exposures, and lead to biased results.[28] Finally, previous analyses in this cohort have suggested a healthy worker effect,[29] which may have led to underestimated or null associations. However, it is notable that we observed a positive association between current exposure and endotype CA1 despite this potential bias.

To our knowledge, this study is the first to investigate the association between occupational exposure to irritants and respiratory endotypes identified by cluster analysis integrating personal, clinical, functional and biological characteristics. Among 2030 healthcare workers,

Su *et al.* identified five asthma clusters using hierarchical clustering and differentially associated with five clusters of irritant exposures.[30] However, the asthma clusters were identified without integrating biological markers and exposure was restricted to cleaning products, making direct comparison with our results difficult. Other studies conducted in clinical settings performed cluster analysis of occupational asthma integrating biological markers (FeNO,[31] blood level of neutrophils and eosinophils [32]), and studied occupational exposures to specific agents (diisocyanates) or groups of HMW or LMW agents,[31,32] but did not include patients with irritant-induced asthma.

Statistical clustering approaches are useful to identify disease endotypes based on a large number of clinical and biological characteristics in the context of highly heterogeneous diseases such as asthma. However, addressing the stability of the resulting clusters and their clinical interpretation is always challenging. Thus, comparison with studies using *a priori* definitions of asthma phenotypes/endotypes is important. Our results are consistent with previous studies which examined individually some of the asthma characteristics included in the definition of endotypes in relation to occupational exposure to irritants. The endotype CA1 was predominantly characterized by adult-onset asthma, poor lung function and respiratory symptoms. Previous studies have shown associations between cleaning products with respiratory symptoms and lower lung function among asthmatics [22] or lung function decline independently of asthma.[21,33] Exposure to vapors, dust, gas and fumes, agents generally considered as irritants, has been associated with poor lung function or lung function decline in many studies.[33,34]

CA1 was also characterized by a high level of FLOPs, a biomarker of damages related to oxidative stress. In the EGEA study,[9] we previously reported that among men without asthma, occupational exposure to irritants and chemicals was associated with higher level of plasma FLOPs. A few other studies investigated the association between biomarkers related to oxidative

stress and occupational asthma induced by irritants. Corradi *et al.* [7] showed that exhaled breath condensate (EBC) H₂O₂ level, was significantly higher in hospital cleaners as compared to controls. Casimirri *et al.* [35] showed that EBC malondialdehyde level, was higher among cleaners compared to non-exposed controls. In contrast, Vizcaya *et al.* [10] did not observe an association between occupational exposure to cleaning products and 8-isoprostanes, another biomarker of lipid peroxidation. Although these studies differ in terms of biological compartment and type of biological marker (reactive oxygen species, lipid peroxidation), overall results including ours, support the role of oxidative stress as one potential mechanism by which occupational exposure to irritants may affect asthma.

Participants in CA1 were also characterized by a high level of neutrophil counts. To our knowledge, few studies investigated the role of non-eosinophilic inflammation in irritant-induced asthma.[36,37] In the Lifelines cohort, occupational exposures to chemicals and pesticides were associated with a lower count of neutrophils at baseline, but no association was found in the longitudinal analysis.[38] In contrast, in EGEA, Matulonga *et al.* [36] reported an association between frequent use of bleach for home cleaning and high neutrophil counts in women with current asthma. In a murine model of irritant-induced asthma, McGovern *et al.* [37] showed that neutrophils were significantly increased after exposure of mice to chlorine gas, a well-known irritant agent. Our results were consistent with the latter findings, suggesting a role of neutrophilic inflammation, in addition to oxidative stress, as potential mechanisms by which occupational exposure to irritants may affect asthma. This hypothesis appears especially relevant since these two mechanisms are closely related.[7] In addition, the absence of association between irritants and the endotype characterized by allergic sensitization and high IgE level (CA3) observed in our study is consistent with the hypothesis of a non-allergic/non-immunologic mechanism of irritant-induced asthma.[2]

Conclusion

Overall, our study shows that occupational exposure to irritants, including chronic low-to-moderate level of exposure, is related to an asthma endotype characterized by poor lung function, respiratory symptoms, high level of fluorescent oxidation products and neutrophil counts. Our results provide additional knowledge on the biological mechanisms of irritant-induced asthma, suggesting oxidative stress and neutrophilic inflammation as potential mechanisms. They support the usefulness of distinct asthma endotypes when studying risk factors for asthma. If replicated, our findings may help improving the recognition and management of irritant-induced work-related asthma. Other studies, in an epidemiological or clinical setting, and possibly including additional relevant pathways for irritant-induced asthma, such as neurogenic inflammation involving transient receptor potential (TRP) channels [39] would be useful to further characterize asthma endotypes related to irritant exposures.

Word counts: 3515 words

References

- 1 Beasley R, Semprini A, Mitchell EA. Risk factors for asthma: is prevention possible? *Lancet* 2015;**386**:1075–85.
- 2 Dumas O, Le Moual N. Do chronic workplace irritant exposures cause asthma? *Curr Opin Allergy Clin Immunol* 2016;**16**:75–85.
- 3 Tarlo SM, Lemiere C. Occupational asthma. *N Engl J Med* 2014;**370**:640–9.
- 4 Malo JL, Tarlo SM, Sastre J, *et al.* An Official American Thoracic Society Workshop Report: Presentations and discussion of the fifth Jack Pepys Workshop on Asthma in the Workplace: Comparisons between asthma in the workplace and non-work-related asthma. *Ann Am Thorac Soc* 2015;**12**:S99–110.
- 5 Vandenplas O, Wiszniewska M, Raulf M, *et al.* EAACI position paper: Irritant-induced asthma. *Allergy* 2014;**69**:1141–53.
- 6 Dumas O, Laurent E, Bousquet J, *et al.* Occupational irritants and asthma: An Estonian cross-sectional study of 34 000 adults. *Eur Respir J* 2014;**44**:647–56.
- 7 Corradi M, Gergelova P, Di Pilato E, *et al.* Effect of exposure to detergents and other chemicals on biomarkers of pulmonary response in exhaled breath from hospital cleaners: A pilot study. *Int Arch Occup Environ Health* 2012;**85**:389–96.
- 8 Czerska M, Zieliński M, Gromadzińska J. Isoprostanes - A novel major group of oxidative stress markers. *Int J Occup Med Environ Health* 2016;**29**:179–90.
- 9 Dumas O, Matran R, Zerimech F, *et al.* Occupational exposures and fluorescent oxidation products in 723 adults of the EGEA study. *Eur Respir J* 2015;**46**:258–61.
- 10 Vizcaya D, Mirabelli MC, Orriols R, *et al.* Functional and biological characteristics of asthma in cleaning workers. *Respir Med* 2013;**107**:673–83.
- 11 Howard R, Rattray M, Prosperi M, *et al.* Distinguishing Asthma Phenotypes Using Machine Learning Approaches. *Curr Allergy Asthma Rep* 2015;**15**:38.
- 12 Kaur R, Chupp G. Phenotypes and endotypes of adult asthma: Moving toward precision medicine. *J Allergy Clin Immunol* 2019;**144**:1–12.
- 13 Nadif R, Febrissy M, Andrianjafimasy MV, *et al.* Endotypes identified by cluster analysis in asthmatics and non-asthmatics and their clinical characteristics at follow-up: the case-control EGEA study. *BMJ Open Resp Res* 2020;**7**:632.
- 14 Le Moual N, Zock J-P, Dumas O, *et al.* Update of an occupational asthma-specific job exposure matrix to assess exposure to 30 specific agents. *Occup Env Med*

- 2018;**75**:507–14.
- 15 Kennedy SM, Le Moual N, Choudat D, *et al.* Development of an asthma specific job exposure matrix and its application in the epidemiological study of genetics and environment in asthma (EGEA). *Occup Env Med* 2000;**57**:635–41.
 - 16 Kauffmann F, Dizier MH, Annesi-Maesano I, *et al.* EGEA (Epidemiological study on the Genetics and Environment of Asthma, bronchial hyperresponsiveness and atopy)--descriptive characteristics. *Clin Exp Allergy* 1999;**29 Suppl 4**:17–21.
 - 17 Nadif R, Bouzigon E, Le Moual N, *et al.* EGEA Collection: a biobank devoted to asthma and asthma-related phenotypes. *Open J Bioresour* 2017;**322**:891–921.
 - 18 Peters U, Dixon A, Forno E. Obesity and Asthma. *J Allergy Clin Immunol* 2018;**141**:1169–79.
 - 19 Mamane A, Baldi I, Tessier JF, *et al.* Occupational exposure to pesticides and respiratory health. *Eur Respir Rev* 2015;**24**:306–19.
 - 20 LeVan TD, Koh W-P, Lee H-P, *et al.* Vapor, dust, and smoke exposure in relation to adult-onset asthma and chronic respiratory symptoms: the Singapore Chinese Health Study. *Am J Epidemiol* 2006;**163**:1118–28.
 - 21 Svanes Ø, Bertelsen RJ, Lygre S, *et al.* Cleaning at Home and at Work in Relation to Lung Function Decline and Airway Obstruction. *Am J Respir Crit Care Med* 2018;**197**:1157–63.
 - 22 Archangelidi O, Sathiyajit S, Consonni D, *et al.* Cleaning products and respiratory health outcomes in occupational cleaners: A systematic review and meta-analysis. *Occup Environ Med* 2020;**in press**. doi:10.1136/oemed-2020-106776
 - 23 Saito R, Virji MA, Henneberger PK, *et al.* Characterization of Cleaning and Disinfecting Tasks and Product Use Among Hospital Occupations. *Am J Ind Med* 2015;**58**:101–11.
 - 24 Vandenas O, Godet J, Hurdubaea L, *et al.* Are high- and low-molecular-weight sensitizing agents associated with different clinical phenotypes of occupational asthma? *Allergy* 2019;**74**:261–72.
 - 25 Maestrelli P, Boschetto P, Fabbri LM, *et al.* Mechanisms of occupational asthma. *J Allergy Clin Immunol* 2009;**123**:531–42;
 - 26 Ge CB, Friesen MC, Kromhout H, *et al.* Use and reliability of exposure assessment methods in occupational case–control studies in the general population: Past, present, and future. *Ann Work Expo Heal* 2018;**62**:1047–63.
 - 27 Bouyer J, Hémon D. Retrospective evaluation of occupational exposures in population-

- based case-control studies: general overview with special attention to job exposure matrices. *Int J Epidemiol* 1993;**22**:S57–64.
- 28 De Matteis S, Jarvis D, Hutchings S, *et al.* Occupations associated with COPD risk in the large population-based UK Biobank cohort study. *Occup Environ Med* 2016;**73**:378–84.
- 29 Dumas O, Le Moual N, Siroux V, *et al.* Work related asthma. A causal analysis controlling the healthy worker effect. *Occup Env Med* 2013;**70**:603–10.
- 30 Su FC, Friesen MC, Humann M, *et al.* Clustering asthma symptoms and cleaning and disinfecting activities and evaluating their associations among healthcare workers. *Int J Hyg Environ Health* 2019;**222**:873–83.
- 31 Lemiere C, NGuyen S, Sava F, *et al.* Occupational asthma phenotypes identified by increased fractional exhaled nitric oxide after exposure to causal agents. *J Allergy Clin Immunol* 2014;**134**:1063–7.
- 32 Mason P, Chiara Frigo A, Cristina Scarpa M, *et al.* Cluster analysis of occupational asthma caused by isocyanates. *J Allergy Clin Immunol* 2018;**142**:2011-2012.e2.
- 33 Le Moual N, Orłowski E, Schenker MB, *et al.* Occupational exposures estimated by means of job exposure matrices in relation to lung function in the PAARC survey. *Occup Env Med* 1995;**52**:634–43.
- 34 Omland O, Wurtz E, Aasen T, *et al.* Occupational chronic obstructive pulmonary disease : a systematic literature review. *Scand J Work Env Heal* 2014;**40**:19–35.
- 35 Casimirri E, Stendardo M, Bonci M, *et al.* Biomarkers of oxidative-stress and inflammation in exhaled breath condensate from hospital cleaners. *Biomarkers* 2016;**21**:115–22.
- 36 Matulonga B, Rava M, Siroux V, *et al.* Women using bleach for home cleaning are at increased risk of non-allergic asthma. *Respir Med* 2016;**117**:264–71.
- 37 McGovern TK, Goldberger M, Allard B, *et al.* Neutrophils mediate airway hyperresponsiveness after chlorine-induced airway injury in the mouse. *Am J Respir Cell Mol Biol* 2015;**52**:513–22.
- 38 Faruque MO, Vonk JM, Bültmann U, *et al.* Airborne occupational exposures and inflammatory biomarkers in the Lifelines cohort study. *Occup Env Med* 2021;**78**:82–5.
- 39 De Matteis S, Ronsmans S, Nemery B. Respiratory Health Effects of Exposure to Cleaning Products. *Clin Chest Med* 2020;**41**:641–50.
- 40 Sunyer J, Pekkanen J, Garcia-Esteban R, *et al.* Asthma score: predictive ability and risk factors. *Allergy* 2007;**62**:142–8.

Table 1: Description of the five respiratory endotypes

| Name | Description | n (%) |
|---------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|
| <i>Among non-asthmatics</i> | | |
| NA1 | Asymptomatic: absence of symptoms | 463 (46.3) |
| NA2 | Respiratory symptoms and dyspnea, use of Inhaled Corticosteroids, high neutrophil and eosinophil counts | 169 (16.9) |
| <i>Among current asthmatics</i> | | |
| CA1 | Active treated (active asthma and use of asthma treatments), adult-onset asthma, poor lung function, respiratory symptoms, highest FLOPs level and neutrophil count | 50 (5.0) |
| CA2 | Rhinitis and low IgE level | 203 (20.3) |
| CA3 | Childhood-onset asthma, allergic sensitization, and highest IgE level | 114 (11.4) |

FLOPs: fluorescent oxidation products, a biomarker of damages related to oxidative stress; IgE: immunoglobulin E.

Table 2: Description of personal, clinical and biological characteristics of the study population at EGEA2 (N=999) according to respiratory endotypes

| | All participants N=999 | Never asthmatics (n=632) | | Current asthmatics (n=367) | | |
|---------------------------------------------------------------|---------------------------|-----------------------------|-----------------|-------------------------------|-----------------|------------------|
| | | NA1 n=463 | NA2 n=169 | CA1 n=50 | CA2 n=203 | CA3 n=114 |
| Age, years | 44.8 ± 15.8 | 47.0 ± 14.8 | 48.5 ± 16.0 | 53.2 ± 15.5 | 41.9 ± 15.1 | 32.4 ± 13.2 |
| Sex, Women | 542 (54.3) | 255 (55.1) | 105 (62.1) | 26 (52.0) | 113 (55.7) | 43 (37.7) |
| Smoking status | | | | | | |
| Non-smokers | 487 (48.8) | 235 (50.8) | 76 (45.0) | 24 (48.0) | 88 (43.4) | 64 (56.1) |
| Ex-smokers | 290 (29.0) | 148 (32.0) | 42 (24.8) | 20 (40.0) | 68 (33.5) | 12 (10.5) |
| Current smokers | 222 (22.2) | 80 (17.3) | 51 (30.2) | 6 (12.0) | 47 (23.1) | 38 (33.3) |
| Body mass index, kg/m² * | | | | | | |
| < 20 | 95 (9.5) | 43 (9.3) | 9 (5.3) | 1 (2.00) | 25 (12.3) | 17 (14.9) |
| [20-25[| 501 (50.2) | 243 (52.5) | 77 (45.6) | 21 (42.0) | 89 (43.8) | 71 (62.3) |
| [25-30[| 295 (29.5) | 135 (29.2) | 59 (34.9) | 15 (30.0) | 67 (33.0) | 19 (16.7) |
| ≥ 30 | 108 (10.8) | 42 (9.1) | 24 (14.2) | 13 (26.0) | 22 (10.8) | 7 (6.1) |
| Age of asthma onset * | | | | | | |
| ≥ 16 years | 133 (36.2) | - | - | 31 (62.0) | 77 (37.9) | 25 (21.9) |
| Asthma symptom score[§], median (min-max) | 1 (0-5) | 0 (0-4) | 1 (0-4) | 4 (1-5) | 2 (0-5) | 1 (0-5) |
| Chronic cough | 81 (8.1) | 1 (0.2) | 31 (18.3) | 21 (42.0) | 24 (11.8) | 4 (3.5) |
| Chronic phlegm | 70 (7.0) | 0 | 27 (16.0) | 16 (32.0) | 22 (10.8) | 5 (4.4) |
| FEV₁ % predicted | 102 ± 18.5 | 108 ± 16.1 | 104 ± 17.7 | 76.2 ± 21.7 | 97.2 ± 17.3 | 97.0 ± 15.6 |
| Total IgE, IU/mL, GM (Q1-Q3) | 67.5 (23.2-195) | 40.3 (15.8-101) | 50.6 (16.1-135) | 144 (65.4-372) | 125 (46-279) | 203 (95.2-536) |
| Eosinophil counts, cells/mm³ | 198 ± 163 | 155 ± 108 | 177 ± 138 | 301 ± 320 | 255 ± 181 | 253 ± 187 |
| Neutrophil counts, cells/mm³ | 4012 ± 1410 | 3915 ± 1278 | 4177 ± 1487 | 4986 ± 1651 | 4130 ± 1584 | 3522 ± 1074 |
| FLOPs level, RFU/mL, GM (Q1-Q3) | 94.2 (80.7-107) | 94.5 (81.8-109) | 97.3 (82.6-111) | 106 (92.1-118) | 91.7 (79.3-105) | 88.9 (77.6-99.2) |

Data are presented as n (%) or mean ± SD unless otherwise stated. GM: geometric mean; Q1: 1st quartile; Q3: 3rd quartile.

* Included as continuous variables in the cluster analysis. FEV₁: forced expiratory volume in one second; IgE: immunoglobulin E; FLOPs: fluorescent oxidation products, a biomarker of damages related to oxidative stress; GM: geometric mean; §five-level of asthma symptom score based on the number of respiratory symptoms during the past 12 months [40], not included in the cluster analysis.

Table 3: Associations between current occupational exposure to any irritant agents and respiratory endotypes (N=999)

| | Occupational exposure to any irritant agents | | | | |
|----------------------|----------------------------------------------|------------------|----------------|-------------------------|----------------|
| | No exposure | Medium | p [§] | High | p [§] |
| NA2 vs. NA1 | | | | | |
| n [#] | 455 (122 vs. 333) | 132 (36 vs. 96) | | 44 (11 vs. 33) | |
| crudeOR (95% CI) | 1 | 1.03 (0.66-1.60) | 0.92 | 0.91 (0.46-1.81) | 0.80 |
| *adjustedOR (95% CI) | 1 | 0.98 (0.61-1.56) | 0.93 | 0.93 (0.46-1.89) | 0.85 |
| CA1 vs. NA1 | | | | | |
| n [#] | 367 (34 vs. 333) | 104 (8 vs. 96) | | 41 (8 vs. 33) | |
| crudeOR (95% CI) | 1 | 0.78 (0.35-1.72) | 0.62 | 2.53 (1.02-6.27) | 0.05 |
| *adjustedOR (95% CI) | 1 | 0.77 (0.34-1.74) | 0.52 | 2.70 (1.00-7.33) | 0.05 |
| CA2 vs. NA1 | | | | | |
| n [#] | 483 (150 vs. 333) | 134 (38 vs. 96) | | 48 (15 vs. 33) | |
| crudeOR (95% CI) | 1 | 0.85 (0.56-1.30) | 0.55 | 1.02 (0.51-1.98) | 0.98 |
| *adjustedOR (95% CI) | 1 | 0.87 (0.56-1.34) | 0.52 | 1.04 (0.52-2.10) | 0.91 |
| CA3 vs. NA1 | | | | | |
| n [#] | 408 (75 vs. 333) | 124 (28 vs. 96) | | 44 (11 vs. 33) | |
| crudeOR (95% CI) | 1 | 1.28 (0.78-2.12) | 0.30 | 1.49 (0.72-3.07) | 0.29 |
| *adjustedOR (95% CI) | 1 | 1.19 (0.69-2.07) | 0.53 | 1.28 (0.56-2.90) | 0.56 |

Significant results in bold.

#n total (n in NA2, CA1, CA2 or CA3 vs. n in NA1); OR: odds ratio for the association between occupational exposure and each endotype (NA2, CA1, CA2 or CA3 vs. NA1) *adjusted for age, sex and smoking status; §for the univariate model (crudeOR), p-values correspond to p-values of Chi-squared test or Fisher exact test.

Table 4: Associations between lifetime occupational exposure to any irritant agents and respiratory endotypes

| | Lifetime occupational exposure to any irritant agents | | |
|----------------------|-------------------------------------------------------|------------------|------------------|
| | No exposure | Medium exposure | High exposure |
| NA2 vs. NA1 | | | |
| n [#] | 303 (83 vs 220) | 177 (50 vs 127) | 151 (36 vs 115) |
| crudeOR (95% CI) | 1 | 1.04 (0.70-1.56) | 0.83 (0.53-1.30) |
| *adjustedOR (95% CI) | 1 | 1.09 (0.71-1.67) | 0.84 (0.52-1.35) |
| CA1 vs. NA1 | | | |
| n [#] | 240 (20 vs 220) | 141 (14 vs 127) | 131 (16 vs 115) |
| crudeOR (95% CI) | 1 | 1.23 (0.61-2.46) | 1.54 (0.76-3.12) |
| *adjustedOR (95% CI) | 1 | 1.28 (0.65-2.53) | 1.37 (0.60-3.14) |
| CA2 vs. NA1 | | | |
| n [#] | 324 (104 vs 220) | 186 (59 vs 127) | 155 (40 vs 115) |
| crudeOR (95% CI) | 1 | 0.97 (0.67-1.41) | 0.73 (0.48-1.12) |
| *adjustedOR (95% CI) | 1 | 0.92 (0.62-1.35) | 0.77 (0.49-1.21) |
| CA3 vs. NA1 | | | |
| n [#] | 276 (56 vs 220) | 166 (39 vs 127) | 134 (19 vs 115) |
| crudeOR (95% CI) | 1 | 1.19 (0.76-1.86) | 0.62 (0.36-1.09) |
| *adjustedOR (95% CI) | 1 | 1.04 (0.62-1.72) | 0.69 (0.36-1.33) |

OR : odds ratio for the association between occupational exposure and each endotype (NA2, CA1, CA2 or CA3 vs. NA1) *adjusted for age, sex and smoking status. #n total (n in NA2, CA1, CA2 or CA3 vs. n in NA1).